



ELSEVIER

Contents lists available at ScienceDirect

European Journal of Pharmacology

journal homepage: www.elsevier.com/locate/ejphar

Immunopharmacology and inflammation

Glucocorticoids decrease Treg cell numbers in lungs of allergic mice

P.C. Olsen^{a,b,*}, J.Z. Kitoko^a, T.P. Ferreira^a, C.T. de-Azevedo^a, A.C. Arantes^a, M.A. Martins^a^a Laboratory of Inflammation, Oswaldo Cruz Institute, Oswaldo Cruz Foundation, Rio de Janeiro, RJ, Brazil^b Laboratory of Clinical Bacteriology and Immunology, Federal University of Rio de Janeiro, Rio de Janeiro, RJ, Brazil

ARTICLE INFO

Article history:

Received 24 September 2014

Received in revised form

26 November 2014

Accepted 28 November 2014

Available online 8 December 2014

Keywords:

Asthma

Treg

Glucocorticoid

ABSTRACT

Glucocorticoids have been the hallmark anti-inflammatory drug used to treat asthma. It has been shown that glucocorticoids ameliorate asthma by increasing numbers and activity of Tregs, in contrast recent data show that glucocorticoid might have an opposite effect on Treg cells from normal mice. Since Tregs are target cells that act on the resolution of asthma, the aim of this study was to elucidate the effect of glucocorticoid treatment on lung Tregs in mouse models of asthma. Allergen challenged mice were treated with either oral dexamethasone or nebulized budesonide. Bronchoalveolar lavage and airway hyperresponsiveness were evaluated after allergenic challenge. Lung, thymic and lymph node cells were phenotyped on Treg through flow cytometry. Lung cytokine secretion was detected by ELISA. Although dexamethasone inhibited airway inflammation and hyperresponsiveness, improving resolution, we have found that both dexamethasone and budesonide induce a reduction of Treg numbers on lungs and lymphoid organs of allergen challenged mice. The reduction of lung Treg levels was independent of mice strain or type of allergen challenge. Our study also indicates that both glucocorticoids do not increase Treg activity through production of IL-10. Glucocorticoid systemic or localized treatment induced thymic atrophy. Taken together, our results demonstrate that glucocorticoids decrease Treg numbers and activity in different asthma mouse models, probably by reducing thymic production of T cells. Therefore, it is possible that glucocorticoids do not have beneficial effects on lung populations of Treg cells from asthmatic patients.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Asthma is a chronic lung inflammatory disease driven by a Th2 response to inhaled allergens in genetically pre-disposed individuals. Lung inflammation in asthma is characterized by infiltration of eosinophils, CD4 lymphocytes and neutrophils, as well as secretion of pro-inflammatory mediators such as Th2 cytokines (Busse and Rosenwasser, 2003). An important imbalance between pro-inflammatory cells and cells that regulate inflammation seems to occur in asthmatic patients, since they present quantitative and functional impairment of pulmonary regulatory T cells, termed Tregs (Hartl et al., 2007; Ling et al., 2004; Nguyen et al., 2009). Besides being constantly stimulated by aeroallergens, the reduced amount of Tregs in asthmatics might also limit the resolution of lung inflammation. Therefore therapeutic measures are of crucial importance in asthma.

Conventional therapy for asthma is generally based on the anti-inflammatory treatment with inhaled glucocorticoid combined preparations with a long-acting β_2 -agonist bronchodilator (Barnes, 2004). Asthmatic patients in general respond well to this therapy, although about 5% need additional and prolonged oral treatment with glucocorticoids, which leads to higher risk of adverse effects (Barnes, 2011a, 2011b; Montuschi and Barnes, 2011). The main mechanisms of action of glucocorticoids are thought to be the direct inhibition of expression of pro-inflammatory mediators and induction of expression of anti-inflammatory mediators, such as IL-10 (Barnes, 2011a).

Pieces of evidence in the literature state that glucocorticoid increases the number and activity of Tregs in asthmatic patients and experimental models of lung allergic disease (Hartl et al., 2007; John et al., 1998; Karagiannidis et al., 2004; Peek et al., 2005). However, most of these studies evaluated Treg numbers in peripheral blood, but not in the lungs. Moreover, detection of Tregs has been done for a long time by using markers that are also present in other lymphocytes such as CD4 and CD25 molecules. Most recently, other groups have shown that glucocorticoids can induce a reduction of Treg numbers in normal mice as well as in different inflammatory models (Sbiera et al., 2011; Stock et al., 2005;

* Corresponding author at: Laboratório de Bacteriologia e Imunologia Clínica, Faculdade de Farmácia, Universidade Federal do Rio de Janeiro, Av. Carlos Chagas Filho 373, Bloco A 2º Andar sala 07, Cidade Universitária, Ilha do Fundão, 21941-902 Rio de Janeiro, RJ, Brazil. Tel. +55 21 25626408.

E-mail address: priolsen@yahoo.com.br (P.C. Olsen).

Wust et al., 2008). Therefore, in this study, we sought to test the effect of oral or nebulized glucocorticoid treatment on numbers and activity of Tregs (CD4⁺CD25⁺Foxp3⁺ and CD4⁺IL10⁺) in the lungs of house dust mite challenged A/J mice.

2. Material and methods

2.1. Animals

Male A/J and BALB/c mice (18–20 g) were obtained from Oswaldo Cruz Foundation breeding facility (Rio de Janeiro, Brazil), and kept at a controlled room temperature (22–25 °C) with a 12 h light/dark cycle. Experimental procedures were approved by the Animal Ethics Committee of the Oswaldo Cruz Foundation (LW-23/10; CEUA-FIOCRUZ, Rio de Janeiro, Brazil). Animal sensitization, challenge and treatment protocol, as well as cell recovery from lungs and IL-10 detection are detailed in the online [Supporting information](#).

2.2. Sensitization, challenge and treatment protocol

A/J or BALB/c mice were exposed to 25 µl of HDM extract (Greer laboratories, Lenoir, NC, USA; *Dermatophagoides pteronyssinus*) intranasally (25 µg of protein), for 3 days/week during 3 weeks. Control mice were challenged only with 25 µl of saline. A group of A/J mice were immunized and challenged with ovalbumin as described previously (Serra et al., 2012).

Challenged mice were treated with dexamethasone or budesonide (Fig. 1). Distinct protocols of dexamethasone treatment were used. Protocol 1: dexamethasone oral administration (1 mg/kg) on the same days of HDM challenge. Protocol 2: dexamethasone oral administration (1 mg/kg) was performed once a day on the last week of challenge for 6 consecutive days. Treatment with nebulized budesonide (7.5 mg/ml) was performed on the same days as protocol 2 of dexamethasone using an aerosol nebulizer (Aeroneb[®] Lab Nebulizer System; Aerogen Ltd., Galway, Ireland) and Buxco-Nebulizer Control-5 (Buxco, Wilmington, USA) which performed 3 inhalation cycles of 10 min each. Dexamethasone was dissolved in sterile saline solution immediately before use. Budesonide was dissolved in a solution of Tween 80 (0.5%) and methanol (0.5%). A group of allergenic-challenged mice was treated with the budesonide vehicle (Tween 80 and methanol). Since this vehicle treated group did not show any difference when compared to the group of challenged mice, it was omitted from the graphs.

2.3. Respiratory mechanics

Airways responsiveness was determined as a change in airway function after the last allergen challenge with aerosolized methacholine in a FinePointe R/C Buxco Platform (Buxco Electronics, Sharon, CT, USA). Lung resistance and dynamic lung compliance

were assessed in anesthetized (nembutal 60 mg/kg, intraperitoneal), tracheostomized, and mechanically ventilated mice, as previously described (Olsen et al., 2011).

2.4. Cell recovery

BALF was retrieved by gentle aspiration and centrifuged (250 g, 10 min at 4 °C). Cell pellets were resuspended in PBS.

Lungs were perfused with saline containing 10 mM EDTA. The right lung was chopped and digested for 1 h with 0.15 mg/ml collagenase (Type D, Roche Diagnostics, Lewes, UK) and 0.025 mg/ml DNase (Type 1, Roche Diagnostics). Cells were retrieved, washed and resuspended in 1 ml RPMI media (10% FBS, 2 mM L-glutamine, 100 U/ml penicillin/streptomycin).

Cervical, inguinal and axillary lymph nodes were extracted, pooled in 1 ml PBS and mechanically homogenized. Thymus was excised and cells suspensions were obtained.

Total cells were counted in Neubauer chamber and differential cell count values were achieved after cyto centrifugation and staining with May-Grunwald-Giemsa. Eosinophils, neutrophils, macrophages and lymphocytes were determined by means of standard morphologic criteria under 1000 × magnification.

2.5. Flow cytometry analyses

CD4CD25Foxp3 characterization was performed with an eBiosciences kit (San Diego, CA, USA).

For characterizing CD4IL10 cells intracellular IL-10 was stained as described previously (Gregory et al., 2009). Samples were analyzed using a FACSCalibur flow cytometer (Becton Dickinson biosciences, San Jose, CA, USA). Analyses were performed on FlowJo 7.6.5 software (Tree Star Inc., Ashland, OR, USA).

2.6. Detection of IL-10

Lung tissue was homogenized as previously described (Serra et al., 2012). Commercial ELISA kits were used for the measurement of IL-10 (DuoSet R&D Systems, Minneapolis, MN, USA) in lung homogenates following instructions of the manufacturer.

2.7. Statistical analyses

Statistical analyses were done using one-way analysis of variance (ANOVA) followed by Newman–Keuls–Student test. Lung mechanics data were analyzed using two-way ANOVA with *post-hoc* Bonferroni correction. Data were expressed as mean ± Standard Error Median (S.E.M.). *P* < 0.05 (two-tailed test) was considered significant.

3. Results

3.1. Effects of dexamethasone on HDM induced lung inflammation and airway hyperreactivity

HDM challenges induced an increase in total leukocyte numbers detected in the BALF collected 1 day after the last challenge (Fig. 2A). The elevation in leukocyte counts was accounted for by an increase in the numbers of eosinophils, lymphocytes, macrophages and neutrophils. Eosinophil and lymphocyte changes were sensitive to treatment with dexamethasone (Fig. 2A). The leukocyte levels, including eosinophils and lymphocytes, naturally decreased in BALF samples during the resolution of lung inflammation, as assessed 7 and 14 days after the last HDM-challenge (Fig. 2C and E). Dexamethasone accelerates resolution, as evidenced by the significant reduction in the levels of these cells recovered in the BALF of treated animals (Fig. 2C and E).




Fig. 1. Schematic timeline of the study design. Mice were challenged with intranasal HDM three times a week for three weeks. Protocol 1 of dexamethasone treatment was performed by orally administering dexamethasone on the same days as the HDM challenge (1 h before each challenge). Protocol 2 of dexamethasone treatment was performed by orally administering dexamethasone for 6 consecutive days during the last week of HDM challenge. Budesonide nebulized treatment was always performed for 6 consecutive days during the last week of HDM challenge.




Fig. 2. Dexamethasone inhibits HDM-evoked lung inflammation and airway hyperreactivity in AJ mice during the peak of inflammation and through the resolution phase. Effects of oral dexamethasone on HDM-evoked leukocyte changes (total and differential counts) in samples of BALF obtained 1 (A), 7 (C) and 14 days (E) after the last challenge. Differential cell counts were identified as eosinophil (Eos), macrophage (Mac), neutrophil (Neut) and lymphocyte (Lymp). AHR in mice challenged with HDM, measured as resistance induced by provocation with increasing concentrations of methacholine, 1 day (B), 7 days (D) and 14 days (F) after the last challenge. Values represent mean \pm S.E.M. from 5 to 6 animals. + $P < 0.05$ as compared with PBS challenged group. * $P < 0.05$ as compared with HDM challenged group.




Fig. 3. Dexamethasone inhibits airway infiltration of Treg on allergen challenged AJ mice at the peak of lung inflammation and during the resolution. Treg cells were evaluated on the BALF of AJ mice 24 h (A), 7 (B) and 14 days (C) after the last HDM challenge. Values represent mean \pm S.E.M. from at least 6 to 8 animals. + $P < 0.05$ as compared with PBS challenged group. * $P < 0.05$ as compared with HDM challenged group.

Antigen challenge with HDM exacerbated airway resistance lung response to inhaled methacholine (3–27 mg/ml) as compared with control mice challenged with PBS, at 24 h after the last challenge (Fig. 2B). During the resolution phase there was also an increase in airway resistance response to methacholine in HDM-challenged mice, compared to PBS (Fig. 2D). By day 14 after the last challenge, there was no difference seen in airway resistance between groups challenged with HDM or PBS (Fig. 2F). Dexamethasone treatment reduced airway resistance in A/J mice HDM-challenged on the peak of inflammation (Fig. 2B), and during resolution (Fig. 2D and F). These results demonstrate that dexamethasone inhibits lung allergic inflammation in A/J mice challenged with HDM.

3.2. Dexamethasone inhibited Treg airway influx during the peak of inflammation as well as during the resolution phase

As shown in other lung inflammation models (Gregory et al., 2009), allergenic challenge increased the numbers of Tregs in the airways of A/J mice, when compared to PBS-challenged mice. Increase in Treg numbers in BALF from HDM-challenged A/J mice peaked at 1 day and decreased during resolution days (Fig. 3). Unexpectedly, dexamethasone treatment inhibited Treg numbers in BALF from HDM-challenged A/J mice during the peak of inflammation (Fig. 3A), as well as during the resolution phase (Fig. 3B and C).

3.3. Glucocorticoids inhibit lung infiltration of Tregs on HDM challenged A/J mice

As glucocorticoids are mostly thought to increase Treg cell numbers (John et al., 1998; Karagiannidis et al., 2004; Peek et al., 2005), we next sought to test a different glucocorticoid drug and a different treatment approach (protocol 1). HDM-challenged A/J mice showed increased numbers of CD4⁺CD25⁺Foxp3⁺ and CD4⁺IL10⁺ both in the airways (Fig. 4A and C) and lung tissue homogenates (Fig. 4B, D and Fig. S1). Dexamethasone pre-treatment (protocol 1) as well as dexamethasone (protocol 2) or budesonide treatment performed on the last week of challenge inhibited Treg numbers both in BALF and digested lung (Fig. 4). Budesonide vehicle had no effect on HDM challenged mice (data not shown).

3.4. Glucocorticoids fail to increase lung secretion of IL-10 on HDM challenged A/J mice

Studies have shown that glucocorticoid treatment increases Treg activity, by enhancing IL-10 secretion (John et al., 1998; Peek et al., 2005). On A/J mice challenged with HDM, neither dexamethasone nor budesonide treatments increased IL-10 production in lungs (Fig. 5). In fact, dexamethasone pre-treatment (protocol 1) actually




Fig. 4. Different glucocorticoids inhibit airway and lung infiltration of Tregs on allergen challenged AJ mice. CD4⁺CD25⁺Foxp3⁺ cells were evaluated on the BALF(A) and lungs (B) of AJ mice treated with either oral dexamethasone (1 mg/kg, protocols 1 and 2) or nebulized budesonide (7.5 mg/ml for 30 min), 24 h after the last HDM challenge. CD4⁺IL-10⁺ cells were also evaluated on the BALF(C) and lungs (D) of AJ mice, 24 h after the last HDM challenge. Values represent mean \pm S.E.M. from at least 4 to 6 animals. ⁺ $P < 0.05$ as compared with PBS challenged group. ^{*} $P < 0.05$ as compared with HDM challenged group.




Fig. 5. Different glucocorticoid treatments fail to increase lung secretion of IL-10 from HDM-challenged mice. Lung secretion of IL-10 induced by HDM was evaluated on HDM-challenged AJ mice treated with either 1 mg/kg oral dexamethasone treatment (protocols 1 or 2), or with nebulized budesonide (7.5 mg/ml for 30 min). Values represent mean \pm S.E.M. from at least 9 to 10 animals. ⁺ $P < 0.05$ as compared with PBS challenged group. ^{*} $P < 0.05$ as compared with HDM challenged group.

reduced secretion of IL-10 in lungs of HDM-challenged mice (Fig. 5), suggesting that glucocorticoids might reduce Treg cell activity.

3.5. Glucocorticoids decrease production and circulation of Treg on HDM- challenged AJ mice

Because glucocorticoids have been described to inhibit lymphocyte migration (Baschant and Tuckermann, 2010; Schweingruber et al., 2014), we sought to study if Treg numbers were decreased in the lungs because they were accumulated in lymphoid organs such as the thymus and lymph nodes. Allergic challenge did not change Treg counts in the thymus (Fig. 6A) or lymph nodes (Fig. 6B), compared to PBS, 1 day after the last challenge. Oral dexamethasone, as well as nebulized budesonide, led to a reduction in CD4⁺CD25⁺Foxp3⁺ Treg numbers in thymus (Fig. 6A) and lymph nodes (Fig. 6B). These results suggest that glucocorticoids can inhibit Treg cell production and circulation in peripheral lymphoid organs.

3.6. Effect of glucocorticoids on thymuses from HDM-challenged AJ mice

HDM challenge did not change size or cell count of thymuses, compared to PBS (Fig. 7). Both dexamethasone protocols, as well as the inhaled budesonide treatment, reduced thymuses weight (Fig. 7A) and cell counts (Fig. 7B), suggesting that different glucocorticoid treatment strategies might affect thymus structure in lung allergic inflammatory models.

3.7. Glucocorticoids inhibit lung infiltration of Treg cells post-HDM challenge on BALB/c mice

As expected, HDM challenge of BALB/c mice induced an increase in Treg cell numbers in the airways (Fig. 8A) and lungs (Fig. 8B), 1 day after the last challenge. Corroborating the results obtained with AJ mice, nebulized budesonide and oral dexamethasone treatment (protocol 2) also decreased Treg cell numbers in the airways (Fig. 8A) and in lung digests (Fig. 8B) of HDM-challenged BALB/c mice. These results suggest that glucocorticoids inhibit lung infiltration of Treg cells post-HDM challenge independent of the mouse strain studied.

3.8. Glucocorticoids inhibit lung infiltration of Treg on AJ mice challenged with OVA

OVA sensitization and challenge also induced an increase in Treg cell numbers in the airways (Fig. 8C) and lungs (Fig. 8D) of AJ mice, compared to PBS-challenged mice at 1 day after the last challenge. As seen with HDM-challenge, dexamethasone oral treatment (protocol 2) and nebulized budesonide decreased Treg cell numbers in the airways (Fig. 8C) and lung digests (Fig. 8D) of OVA-challenged AJ mice. These results suggest that glucocorticoid treatment inhibits lung infiltration of Treg on AJ mice independent of the allergen used to induce the lung inflammation.




Fig. 6. Different glucocorticoid treatments decrease production and circulation of Treg on HDM-challenged AJ mice. Both dexamethasone protocols (oral, 1 mg/kg), as well as budesonide nebulization (7.5 mg/ml 30 min), performed on HDM-challenged mice induce a reduction on number of Tregs in the thymus (A) and in the lymph nodes (B) 24 h after the last challenge. Values represent mean \pm S.E.M. from at least 4 to 6 animals. * $P < 0.05$ as compared with HDM challenged group.




Fig. 7. Different glucocorticoid treatments decrease thymus weight and cellularity. Both dexamethasone protocol treatments (1 mg/kg i.p.), as well as budesonide nebulization (7.5 mg/ml 30 min), performed on HDM-challenged mice induce a reduction on thymus size (A) and a decrease in thymus cellularity (B). Values represent mean \pm S.E.M. from at least 8 to 10 animals. * $P < 0.05$ as compared with HDM challenged group.




Fig. 8. Glucocorticoids inhibit lung infiltration of Treg post-allergen challenge independent of mouse strain and allergen tested. A and B: BALB/c mice were challenged with HDM and treated with either dexamethasone or budesonide using the same protocol described for AJ mice. Intraperitoneal dexamethasone treatment (1 mg/kg), as well as nebulized budesonide (7.5 mg/ml for 30 min), inhibit Treg infiltration to the airways (A) and the lungs (B) of HDM-challenged BALB/c mice. C and D: AJ mice were sensitized and challenged with OVA and treated with either dexamethasone or budesonide as described previously. Intraperitoneal dexamethasone treatment (1 mg/kg), as well as nebulized budesonide (7.5 mg/ml for 30 min), inhibit Treg infiltration to the airways (C) and the lungs (D) of OVA-challenged mice. Values represent mean \pm S.E.M. from at least 4 to 5 animals. + $P < 0.05$ as compared with PBS challenged group. * $P < 0.05$ as compared with HDM challenged group.

4. Discussion

Tregs have been arousing interest as target cells for the improvement of asthma symptoms; therefore it is crucial to understand the

effect of glucocorticoid treatment on these cells in mouse models of allergic lung disease. This study investigated the effect of oral dexamethasone or nebulized budesonide on lung Tregs from allergen-challenged mice. We reported that glucocorticoids inhibit

lung inflammation and airways hyperresponsiveness of A/J mice challenged with HDM at the peak of inflammation, 1 day after the last challenge, and during the resolution phase, especially 7 days after the last challenge. Glucocorticoid treatment, administered systemically or topically, decreases the levels of Tregs in the lungs of allergenic-challenged mice, independent of mouse strain or allergen type used. In the experimental models tested here, reduction on numbers of pulmonary Tregs induced by glucocorticoid might be due to a decrease in the production of lymphocytes in the thymus. These findings support the interpretation that glucocorticoid treatment may not favor the increase of Treg cell numbers or activity in asthma.

Different mice strains have genetic variations that influence respiratory mechanics and airway hyperresponsiveness (Ewart et al., 2000; Held and Uhlig, 2000; Reinhard et al., 2002; Tankersley et al., 1999). Genetic and functional studies have shown that A/J mice are prone to allergic lung inflammation and present naïve AHR to cholinergic stimulation (Antunes et al., 2009; De Sanctis et al., 1995; Ewart et al., 1996; Wills-Karp and Ewart, 1997). We studied a lung allergic inflammatory model in A/J mice by challenging these animals with one of the most relevant allergen in the clinical setting, house dust mite (Nelson et al., 1996). HDM A/J-challenged mice presented several features of severe asthma, such as lung inflammation and AHR, as well as mucus production and collagen deposition (data not shown), reinforcing the importance of this model for pre-clinical studies. Dexamethasone oral administration two weeks after the first HDM challenge, when inflammation was already present, inhibited lung inflammatory cell infiltration as well as the AHR at the peak of inflammation, as previously reported (Olsen et al., 2011; Ulrich et al., 2008).

GC had been known to accelerate the resolution of inflammation through diverse mechanisms (Brannan, 2010; Hutchinson et al., 2011). In our asthma mouse model, allergen induced inflammation and AHR decrease during the resolution phase, 7 and 14 days post-challenge. As expected, dexamethasone oral treatment administered during the last week of HDM challenge accelerated the natural resolution of inflammation, noted by a reduction of the eosinophil airway influx and AHR. Others found that GCs may improve resolution of inflammation by increasing Treg numbers and their anti-inflammatory activity (Chen et al., 2006; Karagiannidis et al., 2004). Leech et al. showed that Tregs migrate to lungs and draining lymph nodes after the allergenic challenge and reach their peak in number just before the resolution starting point (Leech et al., 2007). Tregs might play a pivotal role in resolution of lung allergic inflammation, since they can regulate immune responses through cell-to-cell contact or secretion of immunosuppressive cytokines, such as IL-10 (Kearley et al., 2005; Lewkowich et al., 2005; Zhang et al., 2014). Surprisingly, we have seen that dexamethasone oral treatment, although accelerating the resolution of inflammation, reduced Treg numbers in the lungs and lymph nodes of HDM-challenged mice, contradicting other studies (Hartl et al., 2007; Karagiannidis et al., 2004). Interestingly, there is a large heterogeneity in the characterization of Tregs, since most groups characterize these cells only by the expression of CD4 and higher expression of CD25, while other groups search only for the indirect presence of Tregs by testing IL-10 production or Foxp3 expression (Hartl et al., 2007; Stelmach et al., 2002). Although there are several additional surface markers used to identify Tregs, the best way to define natural Tregs is probably to analyze the expression of Foxp3 in CD4⁺CD25⁺ cells (Sakaguchi et al., 2010; Wieckiewicz et al., 2010). Furthermore, Treg cell populations have been analyzed on blood samples instead of relevant inflammatory sites for asthma, such as the lungs (Karagiannidis et al., 2004). The present study demonstrated that GCs decrease the numbers of Treg cells, characterized by expression of CD4, CD25 and Foxp3, in the BALF and lung digests of allergen challenged mice.

Aside from the thymus-derived CD4⁺CD25⁺Foxp3⁺ Treg cells, other T regulatory cells also play a role in maintaining peripheral tolerance to airway allergens. These cells are known as adaptive or induced Tregs (iTreg) because they are induced in the periphery. iTregs are capable of secreting high levels of anti-inflammatory IL-10 and TGF- β which might be important in the protection against asthma (Hawrylowicz and O'Garra, 2005). Some studies have shown that GCs increase IL-10 production (John et al., 1998; Stelmach et al., 2002), we observed that lung secretion of IL-10 was reduced by the dexamethasone pre-treatment and unaltered by dexamethasone and nebulized budesonide treatment that started two weeks after the first HDM challenge. These results suggest that dexamethasone treatment can impair IL-10 secreting Treg quantity and activity in murine models of asthma.

Although we have used two different steroid drugs through distinct administration routes, budesonide and dexamethasone were both able to decrease Treg levels in lungs of allergen challenged mice. In addition, dexamethasone was tested in two distinct protocols, one beginning at the same time as the first HDM challenge and the other starting on the third week of antigen challenge, when lung inflammation was already established. Notably, both dexamethasone protocols were able to decrease Treg lung infiltration. These results are in line with a prior study, which reported reduced Treg cell numbers in spleen and blood of C57/BL6 normal mice after dexamethasone intraperitoneal treatment (Sbiera et al., 2011). Sbiera et al. demonstrated that the decrease of Treg cell population induced by GC therapy is not dose dependent. It is noteworthy that the inhaled budesonide used in our study, which is used clinically for the treatment of asthma, had the same effects as oral dexamethasone. Nonetheless, the daily dose of budesonide or dexamethasone administered to mice in our study is lower than the amount generally used by asthmatic patients (Powell and Gibson, 2003; Szeffler et al., 2002). Therefore, it seems possible that continuous GC treatment in humans also reduce lung Treg cell numbers.

It is known that inhaled GC in asthma causes fewer side effects compared to oral GCs (Hagan et al., 2014). We showed that nebulized budesonide treatment, as well as oral dexamethasone treatment, reduced thymic size, thymic cell counts and amount of Tregs in the thymus and lymph nodes. These results indicate that GCs inhibit Treg thymic production regardless of the route of administration. It is likely that the decrease in Treg cell numbers observed in the lungs of allergenic challenged mice was due to the immunosuppressive effects of GCs in the thymus.

Our observations demonstrate that systemic or localized GC treatment can impair Treg quantity and activity in lungs of allergenic-challenged mice, independent of mouse strain or type of allergen tested. As Tregs are important target cells for the treatment of asthma, it is relevant to carefully analyze the effect of in vivo treatment with GCs on these cell populations in lungs of asthmatic patients.

Acknowledgments

We are indebted to Nathália de Andrade Guimarães for her technical support. Priscilla C. Olsen and Jamil Zola Kitoko acknowledge CNPq for their Fellowship (162606/2011-7). Priscilla C. Olsen acknowledges FAPERJ for their financial support (E-26/112.492/2012).

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.ejphar.2014.11.034>.

References

- Antunes, M.A., Abreu, S.C., Damaceno-Rodrigues, N.R., Parra, E.R., Capelozzi, V.L., Pinart, M., Romero, P.V., Silva, P.M., Martins, M.A., Rocco, P.R., 2009. Different strains of mice present distinct lung tissue mechanics and extracellular matrix composition in a model of chronic allergic asthma. *Respir. Physiol. Neurobiol.* 165, 202–207.
- Barnes, P.J., 2004. New drugs for asthma. *Nat. Rev. Drug Discov.* 3, 831–844.
- Barnes, P.J., 2011a. Biochemical basis of asthma therapy. *J. Biol. Chem.* 286, 32899–32905.
- Barnes, P.J., 2011b. Glucocorticosteroids: current and future directions. *Br. J. Pharmacol.* 163, 29–43.
- Baschant, U., Tuckermann, J., 2010. The role of the glucocorticoid receptor in inflammation and immunity. *J. Steroid Biochem. Mol. Biol.* 120, 69–75.
- Brannan, J.D., 2010. Bronchial hyperresponsiveness in the assessment of asthma control: Airway hyperresponsiveness in asthma: its measurement and clinical significance. *Chest* 138, 11S–17S.
- Busse, W.W., Rosenwasser, L.J., 2003. Mechanisms of asthma. *J. Allergy Clin. Immunol.* 111, S799–S804.
- Chen, X., Oppenheim, J.J., Winkler-Pickett, R.T., Ortaldo, J.R., Howard, O.M., 2006. Glucocorticoid amplifies IL-2-dependent expansion of functional FoxP3(+)CD4(+)CD25(+) T regulatory cells in vivo and enhances their capacity to suppress EAE. *Eur. J. Immunol.* 36, 2139–2149.
- De Sanctis, G.T., Merchants, M., Beier, D.R., Dredge, R.D., Grobholz, J.K., Martin, T.R., Lander, E.S., Drazen, J.M., 1995. Quantitative locus analysis of airway hyperresponsiveness in A/J and C57BL/6J mice. *Nat. Genet.* 11, 150–154.
- Ewart, S.L., Kuperman, D., Schadt, E., Tankersley, C., Grupe, A., Shubitowski, D.M., Peltz, G., Wills-Karp, M., 2000. Quantitative trait loci controlling allergen-induced airway hyperresponsiveness in inbred mice. *Am. J. Respir. Cell Mol. Biol.* 23, 537–545.
- Ewart, S.L., Mitzner, W., DiSilvestre, D.A., Meyers, D.A., Levitt, R.C., 1996. Airway hyperresponsiveness to acetylcholine: segregation analysis and evidence for linkage to murine chromosome 6. *Am. J. Respir. Cell Mol. Biol.* 14, 487–495.
- Gregory, L.G., Causton, B., Murdoch, J.R., Mathie, S.A., O'Donnell, V., Thomas, C.P., Priest, F.M., Quint, D.J., Lloyd, C.M., 2009. Inhaled house dust mite induces pulmonary T helper 2 cytokine production. *Clin. Exp. Allergy: J. Br. Soc. Allergy Clin. Immunol.* 39, 1597–1610.
- Hagan, J.B., Samant, S.A., Volcheck, G.W., Li, J.T., Hagan, C.R., Erwin, P.J., Rank, M.A., 2014. The risk of asthma exacerbation after reducing inhaled corticosteroids: a systematic review and meta-analysis of randomized controlled trials. *Allergy* 69, 510–516.
- Hartl, D., Koller, B., Mehlhorn, A.T., Reinhardt, D., Nicolai, T., Schendel, D.J., Griese, M., Krauss-Etschmann, S., 2007. Quantitative and functional impairment of pulmonary CD4+CD25hi regulatory T cells in pediatric asthma. *J. Allergy Clin. Immunol.* 119, 1258–1266.
- Hawrylowicz, C.M., O'Garra, A., 2005. Potential role of interleukin-10-secreting regulatory T cells in allergy and asthma. *Nat. Rev. Immunol.* 5, 271–283.
- Held, H.D., Uhlig, S., 2000. Basal lung mechanics and airway and pulmonary vascular responsiveness in different inbred mouse strains. *J. Appl. Physiol.* 88, 2192–2198.
- Hutchinson, J.L., Rajagopal, S.P., Sales, K.J., Jabbar, H.N., 2011. Molecular regulators of resolution of inflammation: potential therapeutic targets in the reproductive system. *Reproduction* 142, 15–28.
- John, M., Lim, S., Seybold, J., Jose, P., Robichaud, A., O'Connor, B., Barnes, P.J., Chung, K.F., 1998. Inhaled corticosteroids increase interleukin-10 but reduce macrophage inflammatory protein-1alpha, granulocyte-macrophage colony-stimulating factor, and interferon-gamma release from alveolar macrophages in asthma. *Am. J. Respir. Crit. Care Med.* 157, 256–262.
- Karagiannidis, C., Akdis, M., Holopainen, P., Woolley, N.J., Hense, G., Ruckert, B., Mantel, P.Y., Menz, G., Akdis, C.A., Blaser, K., Schmidt-Weber, C.B., 2004. Glucocorticoids upregulate FOXP3 expression and regulatory T cells in asthma. *J. Allergy Clin. Immunol.* 114, 1425–1433.
- Kearley, J., Barker, J.E., Robinson, D.S., Lloyd, C.M., 2005. Resolution of airway inflammation and hyperreactivity after in vivo transfer of CD4+CD25+ regulatory T cells is interleukin 10 dependent. *J. Exp. Med.* 202, 1539–1547.
- Leech, M.D., Benson, R.A., De Vries, A., Fitch, P.M., Howie, S.E., 2007. Resolution of Der p1-induced allergic airway inflammation is dependent on CD4+CD25+Foxp3+ regulatory cells. *J. Immunol.* 179, 7050–7058.
- Lewkowich, I.P., Herman, N.S., Schleifer, K.W., Dance, M.P., Chen, B.L., Dienger, K.M., Sproles, A.A., Shah, J.S., Kohl, J., Belkaid, Y., Wills-Karp, M., 2005. CD4+CD25+ T cells protect against experimentally induced asthma and alter pulmonary dendritic cell phenotype and function. *J. Exp. Med.* 202, 1549–1561.
- Ling, E.M., Smith, T., Nguyen, X.D., Pridgeon, C., Dallman, M., Arbery, J., Carr, V.A., Robinson, D.S., 2004. Relation of CD4+CD25+ regulatory T-cell suppression of allergen-driven T-cell activation to atopic status and expression of allergic disease. *Lancet* 363, 608–615.
- Montuschi, P., Barnes, P.J., 2011. New perspectives in pharmacological treatment of mild persistent asthma. *Drug Discov. Today* 16, 1084–1091.
- Nelson Jr., R.P., DiNicolo, R., Fernandez-Caldas, E., Seleznick, M.J., Lockey, R.F., Good, R.A., 1996. Allergen-specific IgE levels and mite allergen exposure in children with acute asthma first seen in an emergency department and in nonasthmatic control subjects. *J. Allergy Clin. Immunol.* 98, 258–263.
- Nguyen, K.D., Vanichsarn, C., Fohner, A., Nadeau, K.C., 2009. Selective deregulation in chemokine signaling pathways of CD4+CD25(hi)CD127(lo)(-/-) regulatory T cells in human allergic asthma. *J. Allergy Clin. Immunol.* 123, 933–939 (e910).
- Olsen, P.C., Ferreira, T.P., Serra, M.F., Farias-Filho, F.A., Fonseca, B.P., Viola, J.P., Cordeiro, R.S., Silva, P.M., Costa, J.C., Martins, M.A., 2011. Lidocaine-derivative JMF2-1 prevents ovalbumin-induced airway inflammation by regulating the function and survival of T cells. *Clin. Exp. Allergy: J. Br. Soc. Allergy Clin. Immunol.* 41, 250–259.
- Peek, E.J., Richards, D.F., Faith, A., Lavender, P., Lee, T.H., Corrigan, C.J., Hawrylowicz, C.M., 2005. Interleukin-10-secreting “regulatory” T cells induced by glucocorticoids and beta2-agonists. *Am. J. Respir. Cell Mol. Biol.* 33, 105–111.
- Powell, H., Gibson, P.G., 2003. Inhaled corticosteroid doses in asthma: an evidence-based approach. *Med. J. Aust.* 178, 223–225.
- Reinhardt, C., Eder, G., Fuchs, H., Ziesenis, A., Heyder, J., Schulz, H., 2002. Inbred strain variation in lung function. *Mamm. Genome* 13, 429–437.
- Sakaguchi, S., Miyara, M., Costantino, C.M., Hafler, D.A., 2010. FOXP3+ regulatory T cells in the human immune system. *Nat. Rev. Immunol.* 10, 490–500.
- Sbiera, S., Dexeit, T., Reichardt, S.D., Michel, K.D., van den Brandt, J., Schull, S., Kraus, L., Beyer, M., Mlynski, R., Wortmann, S., Allolio, B., Reichardt, H.M., Fassnacht, M., 2011. Influence of short-term glucocorticoid therapy on regulatory T cells in vivo. *PLoS One* 6, e24345.
- Schweingruber, N., Fischer, H.J., Fischer, L., van den Brandt, J., Karabinskaya, A., Labi, V., Villunger, A., Kretzschmar, B., Huppke, P., Simons, M., Tuckermann, J.P., Flügel, A., Lühder, F., Reichardt, H.M., 2014. *Acta Neuropathol.* 127 (5), 713–729, May.
- Serra, M.F., Anjos-Valotta, E.A., Olsen, P.C., Couto, G.C., Jurgilas, P.B., Cotias, A.C., Pao, C.R., Ferreira, T.P., Arantes, A.C., Pires, A.L., Cordeiro, R.S., Silva, P.M., Martins, M.A., 2012. Nebulized lidocaine prevents airway inflammation, peribronchial fibrosis, and mucus production in a murine model of asthma. *Anesthesiology* 117, 580–591.
- Stelmach, I., Jerzynska, J., Kuna, P., 2002. A randomized, double-blind trial of the effect of glucocorticoid, antileukotriene and beta-agonist treatment on IL-10 serum levels in children with asthma. *Clin. Exp. Allergy: J. Br. Soc. Allergy Clin. Immunol.* 32, 264–269.
- Stock, P., Akbari, O., DeKruyff, R.H., Umetsu, D.T., 2005. Respiratory tolerance is inhibited by the administration of corticosteroids. *J. Immunol.* 175, 7380–7387.
- Szefer, S.J., Martin, R.J., King, T.S., Boushey, H.A., Chmiack, R.M., Chinchilli, V.M., Craig, T.J., Dolovich, M., Drazen, J.M., Fagan, J.K., Fahy, J.V., Fish, J.E., Ford, J.G., Israel, E., Kiley, J., Kraft, M., Lazarus, S.C., Lemanske Jr., R.F., Mauger, E., Peters, S.P., Sorkness, C.A., Asthma Clinical Research Network of the National Heart, Lung and Blood Institute, 2002. Significant variability in response to inhaled corticosteroids for persistent asthma. *J. Allergy Clin. Immunol.* 109, 410–418.
- Tankersley, C.G., Rabold, R., Mitzner, W., 1999. Differential lung mechanics are genetically determined in inbred murine strains. *J. Appl. Physiol.* 86, 1764–1769.
- Ulrich, K., Hincks, J.S., Walsh, R., Wetterstrand, E.M., Fidock, M.D., Sreckovic, S., Lamb, D.J., Douglas, G.J., Yeadon, M., Perros-Huguet, C., Evans, S.M., 2008. Anti-inflammatory modulation of chronic airway inflammation in the murine house dust mite model. *Pulm. Pharmacol. Ther.* 21, 637–647.
- Wieckiewicz, J., Goto, R., Wood, K.J., 2010. T regulatory cells and the control of autoimmunity: from characterisation to clinical application. *Curr. Opin. Immunol.* 22, 662–668.
- Wills-Karp, M., Ewart, S.L., 1997. The genetics of allergen-induced airway hyperresponsiveness in mice. *Am. J. Respir. Crit. Care Med.* 156, S89–S96.
- Wust, S., van den Brandt, J., Tischner, D., Kleiman, A., Tuckermann, J.P., Gold, R., Lühder, F., Reichardt, H.M., 2008. Peripheral T cells are the therapeutic targets of glucocorticoids in experimental autoimmune encephalomyelitis. *J. Immunol.* 180, 8434–8443.
- Zhang, M., Qian, Y.Y., Chai, S.J., Liang, Z.Y., Xu, Q., Wu, Z.Q., Wang, K., 2014. Enhanced local Foxp3 expression in lung tissue attenuates airway inflammation in a mouse model of asthma. *J. Asthma* 51 (5), 451–458, Jun.