



Short communication

Detection and quantification of *Leishmania braziliensis* in ectoparasites from dogs



Rayana Carla Silva de Morais^{a,b}, Suênia da Cunha Gonçalves-de-Albuquerque^a, Rômulo Pessoa e Silva^a, Pietra Lemos Costa^a, Kamila Gaudêncio da Silva^b, Fernando José da Silva^a, Sinval Pinto Brandão-Filho^a, Filipe Dantas-Torres^{a,c}, Milena de Paiva-Cavalcanti^{a,*}

^a Department of Immunology, Aggeu Magalhães Research Center, Oswaldo Cruz Foundation, Av. Prof. Moraes Rego s/n, 50670-420 Recife, Pernambuco, Brazil

^b Federal Rural University of Pernambuco, R. Dom Manoel de Medeiros s/n, 52171-900 Recife, Pernambuco, Brazil

^c Department of Veterinary Medicine, University of Bari, Str. Provinciale per Casamassima km 3, 70010 Valenzano, Bari, Italy

ARTICLE INFO

Article history:

Received 10 January 2013

Received in revised form 18 March 2013

Accepted 21 March 2013

Keywords:

Ectoparasites

Dogs

Leishmania braziliensis

PCR

Real time PCR

ABSTRACT

American cutaneous leishmaniasis (ACL) is a disease caused by different species of *Leishmania* protozoa, *Leishmania braziliensis* being the main species found in Brazil. In this study, two rural areas in Pernambuco, northeastern Brazil, where ACL is endemic, were selected. Genomic DNA was extracted from canine ectoparasites (ticks, fleas, and lice) and tested using a conventional PCR and a quantitative real time PCR. A total of 117 ectoparasites were collected, being 50 (42.74%) of them positive for *L. braziliensis* (in at least one PCR protocol), with a mean parasite load of 14.14 fg/μL. Furthermore, 46 (92.00%) positive ectoparasites were collected from positive dogs and 4 (8.00%) from negative ones. This study reports the detection of *L. braziliensis* DNA in ectoparasites, but does not prove their vector competence. Certainly, experimental transmission studies are necessary to assess their role, if any, in the transmission of *Leishmania* parasites to dogs.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

American cutaneous leishmaniasis (ACL) constitutes an important public health problem, whose control is difficult to achieve. In Brazil, ACL is on the rise, with autochthonous cases of human infection by *Leishmania braziliensis* in all Brazilian regions (Brito et al., 2012). The difficulties in controlling ACL may be related to the complex epidemiology of this disease, which involves the participation of different vector species, different wild and domestic animal hosts (small rodents, marsupials, dogs, cats, and horses)

and humans as well, that interact in different kinds of environments (e.g., primary forests, human houses, and crop plantations).

In spite of the numerous reports of natural infection by *L. braziliensis* in dogs, their role in the life cycle of this parasite has not been clarified yet (Brandão-Filho et al., 2011). In recent years, there has been a growing scientific interest about the possible role of ectoparasites in the epidemiology of canine visceral leishmaniasis (Coutinho et al., 2005; Solano-Gallego et al., 2012), but their actual importance, if any, in the transmission cycle of *Leishmania infantum* remains unknown (Dantas-Torres, 2011).

In the present study, we assessed the presence of *L. braziliensis* in canine ectoparasites collected in the framework of a previous study (Morais et al., 2013) carried out in northeastern Brazil.

* Corresponding author. Tel.: +55 81 21012679; fax: +55 81 21012640.
E-mail addresses: milena.cavalcanti@bol.com.br, mp@cpqam.fiocruz.br (M.d. Paiva-Cavalcanti).

2. Material and methods

Ectoparasites were collected from dogs living in two rural areas in Pernambuco State. Site A and site B were in the municipalities of Igarassu (7°50'02" S, 34°54'21" W) and Passira (07°59'42" S, 35°34'51" W), respectively. Ectoparasites were collected from dogs by hand collection and maintained in 70% ethanol until identification using morphological taxonomic keys (Aragão and Fonseca, 1961; Guimarães et al., 2001). The Commission for Ethics in the Use of Animals (CEUA, FIOCRUZ, Brazil) of our institution approved the study protocol and owners gave their informed written consent before ectoparasite collection.

DNA from ectoparasites was extracted using Illustra tissue and cells genomicPrep Mini Spin kit (GE Healthcare), following the manufacturer's instructions. The extracted DNA was stored at -20°C for subsequent amplification by a conventional polymerase chain reaction (PCR) according to Brujin and Barker (1992) and by a quantitative real time PCR (qPCR), as described elsewhere (Paiva-Cavalcanti et al., in press). The level of agreement between results obtained with both PCR protocols was assessed using Kappa statistics. Statistical analysis was performed using BioEstat 5.0 (Ong Mamiraua, Belém, PA, Brazil).

3. Results

A total of 117 ectoparasites (21 ectoparasites from site A and 96 from site B) were collected from 99 dogs. Ectoparasites included fleas (*Ctenocephalides felis*), ticks (*Rhipicephalus sanguineus*), and lice (*Heterodoxus spiniger*).

A total of 50 (42.74%) ectoparasites were positive for *L. braziliensis* (Fig. 1), being 12 (41.38%) *C. felis*, 32 (42.67%) *R. sanguineus*, and 6 (46.15%) *H. spiniger* (Table 1). The mean parasite load detected in positive ectoparasites was 14.14 fg, ranging from 0.37 fg to 113.37 fg (Ct 38.60 and Ct 31.38, respectively). Furthermore, a total of 46 (92.00%) positive ectoparasites were collected from positive dogs and four from negative ones (Table 2).

Comparing the two PCR protocols used (i.e., PCR and qPCR), a low level of agreement was found (Table 1), being most qPCR-positive ectoparasites negative when tested by PCR.

4. Discussion

Again, this study reports the detection and quantification of *L. braziliensis* DNA in ectoparasites collected from dogs living in an area of ACL-endemicity. Remarkably, some positive ectoparasites were collected from negative dogs, indicating that they acquired the infection by feeding on




Fig. 1. Real time PCR amplification curve showing *L. braziliensis*-positive ectoparasites (PE₁ and PE₂) collected from dogs. Positive controls (100 pg/ μL and 100 fg/ μL = PC_{100pg} and PC_{100fg}, respectively) and no-template control (NTC) are also shown.

another infected dog. Dogs living in the study sites are highly exposed to *L. infantum* and *L. braziliensis* (Morais et al., 2013; Paiva-Cavalcanti et al., in press) and an outbreak of ACL in humans has recently been reported in site A (data not shown). Interestingly, human cases of ACL have not been reported in site B, where visceral leishmaniosis is endemic. This suggests that human cases of ACL may have been underreported in this area. On the other hand, it is important to clarify that the presence of dogs and ectoparasites positive to *L. braziliensis* in site A, does not imply that dogs or ectoparasites are involved in transmission cycle of this parasite.

Domestic animals are not considered important reservoirs of *L. braziliensis* (Brandão-Filho et al., 2011). However, some of these animals (e.g., chickens, cattle, and horses) are highly attractive to phlebotomine sand fly vectors and may play a part in the epidemiology of ACL (Brandão-Filho et al., 2011). The construction of human houses and the presence

Table 1

Positivity to *L. braziliensis* by conventional PCR (PCR) and real-time PCR (qPCR) in ectoparasites collected from dogs in Pernambuco, northeastern Brazil.

Ectoparasites	n	Positive (overall)		PCR	qPCR	Kappa statistics	
		n	%			Kappa	P value
<i>Ctenocephalides felis</i>	29	12	41.38	2	11	0.04	0.36
<i>Heterodoxus spiniger</i>	13	6	46.15	1	5	-0.15	0.21
<i>Rhipicephalus sanguineus</i>	75	32	42.67	1	31	-0.03	0.20
Total	117	50	42.73	4	47	-0.03	0.26

Table 2Positivity to *L. braziliensis* in ectoparasites collected from dogs either positive or negative by conventional PCR and/or real-time PCR.

Ectoparasites	n total	From positive dogs (n = 252)		From negative dogs (n = 23)	
		n	%	n	%
<i>Ctenocephalides felis</i>	12	10	83.33	2	16.67
<i>Heterodoxus spiniger</i>	6	5	83.33	1	16.67
<i>Rhipicephalus sanguineus</i>	32	31	96.87	1	3.13
Total	50	46	92.00	4	8.00

of domestic animals near forested areas, as commonly observed in rural areas in northeastern Brazil, may affect the epidemiology of ACL. Indeed, large populations of phlebotomine sand flies may establish in animal shelters, such as chicken coops and stables (Guimarães et al., 2012), thus increasing the risk of human infection (Brandão-Filho et al., 1999). Deforestation, changes in land use, construction of roads and climate changes may also alter the epidemiology of ACL, as these factors may directly impact on the ecology and distribution of vectors and reservoirs of *Leishmania* parasites (Dantas-Torres et al., 2012). However, it is imperative that populations in endemic areas are informed by public health authorities that dogs are not considered epidemiologically relevant reservoirs of *L. braziliensis* to avoid the unnecessary elimination of infected dogs.

Recent studies have reported the detection of *L. infantum* DNA or RNA fragments in ticks, fleas and lice (Coutinho et al., 2005; Colombo et al., 2011; Solano-Gallego et al., 2012; Morais et al., 2013). In this study, we reported for the first time the detection of *L. braziliensis* in ticks, fleas and lice collected from positive and negative dogs. Interestingly, most *L. braziliensis*-positive ectoparasites were also *L. infantum*-positive (data not shown). Furthermore, the mean parasite load found for *L. braziliensis* (14.14 fg/ μ L) was higher than to that previously reported for *L. infantum* (9.10 fg/ μ L) (Morais et al., 2013). Certainly, these findings should be interpreted with caution because blood-feeding arthropods may ingest whatever is in the blood of their hosts (Otranto and Dantas-Torres, 2010). This study reports the detection of *L. braziliensis* DNA in ectoparasites, but does not prove their vector competence. Certainly, the finding of *L. braziliensis*-positive chewing lice (*H. spiniger*) may indicate that even non-hematophagous arthropods may eventually ingest parasites that are present in the blood or lymph of dogs. However, experimental studies are necessary to achieve a definitive conclusion regarding the role of ectoparasites in the transmission of *Leishmania* parasites to dogs.

Conflict of interest

The authors declare there are no conflicts of interest.

Acknowledgments

Thanks to Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco. (FACEPE) (technical support), Centro de Pesquisas Aggeu Magalhães (CPqAM/FIOCRUZ-PE) (technical support, infrastructure and consumables) and

Program for Technical Development of Health Inputs-PDTIS FIOCRUZ for the use of its facilities.

References

- Aragão, H.B., Fonseca, F., 1961. Notas de Ixodologia VIII Lista e chave para representantes da fauna de Ixodológica Brasileira. Mem. Inst. Oswaldo Cruz. 59, 115–129.
- Brandão-Filho, S.P., Campbell-Lendrum, D., Brito, M.E.F., Shaw, J.J., Davies, C.R., 1999. Epidemiological surveys confirm an increasing burden of cutaneous leishmaniasis northeast Brazil. Trans. R. Soc. Trop. Med. Hyg. 93, 488–494.
- Brandão-Filho, S.P., Dantas-Torres, F., Tolezano, J.E., Shaw, J.J., 2011. Hospedeiros reservatórios de *Leishmania* spp. associados à leishmaniose tegumentar americana com especial ênfase no Brasil. In: Barral, A., Costa, J. (Eds.), *Leishmanias e a leishmaniose tegumentar nas Américas*. Cyted, CNPq Fiocruz-BA, Salvador, pp. 65–74.
- Brito, M.E.F., Andrade, M.S., Dantas-Torres, F., Rodrigues, E.H.G., Paiva-Cavalcanti, M., Almeida, A.M.P., Brandão-Filho, S.P., 2012. Cutaneous leishmaniasis in northeastern Brazil: a critical appraisal of studies conducted in State of Pernambuco. Rev. Soc. Bras. Med. Trop. 45, 1–4.
- Bruijn, M., Barker, D.C., 1992. Diagnosis of New World leishmaniasis: specific detection of species of the *Leishmania braziliensis* complex by amplification of kinetoplast DNA. Acta Trop. 52, 45–58.
- Colombo, F.A., Odorizzi, R.M.F.N., Laurenti, M.D., Galati, E.A.B., Canavez, F., Pereira-Chioccola, V.L., 2011. Detection of *Leishmania (Leishmania) infantum* RNA in fleas and ticks collected from naturally infected dogs. Parasitol. Res. 109, 267–274.
- Coutinho, M.T., Bueno, L.L., Sterzik, A., Fujiwara, R.T., Botelho, J.R., Maria, M., Genaro, O., Linardi, P.M., 2005. Participation of *Rhipicephalus sanguineus* (Acari: Ixodidae) in the epidemiology of canine visceral leishmaniasis. Vet. Parasitol. 128, 149–155.
- Dantas-Torres, F., 2011. Ticks as vectors of *Leishmania* parasites. Trends Parasitol. 27, 155–159.
- Dantas-Torres, F., Solano-Gallego, L., Baneth, G., Ribeiro, V.M., de Paiva-Cavalcanti, M., Otranto, D., 2012. Canine leishmaniasis in the Old and New Worlds: unveiled similarities and differences. Trends Parasitol. 28, 531–538.
- Guimarães, J.H., Tucci, E.C., Barros-Battesti, D.M. (Eds.), 2001. Ectoparasitas de importância veterinária. Plêiade/FAPESP, São Paulo, p. 218.
- Guimarães, V.C.F., Costa, P.L., Silva, J.F., Silva, K.T., Silva, K.G., Araújo, A.I.F., Rodrigues, E.H.G., Brandão-Filho, S.P., 2012. Phlebotomine sandflies (Diptera: Psychodidae) in São Vicente Férrer, a sympatric area to cutaneous and visceral leishmaniasis in the state of Pernambuco, Brazil. Rev. Soc. Bras. Med. 45, 66–70.
- Morais, R.C.S., Gonçalves, S.C., Costa, P.L., Silva, K.G., Silva, F.J., Silva, R.P., Brito, M.E.F., Brandão-Filho, S.P., Dantas-Torres, F., Paiva-Cavalcanti, M., 2013. Detection of *Leishmania infantum* in animals and their ectoparasites by conventional PCR and real time PCR. Exp. Appl. Acarol. 59, 473–481.
- Otranto, D., Dantas-Torres, F., 2010. Fleas and ticks as vectors of *Leishmania* spp. to dogs: Caution is needed. Vet. Parasitol. 168, 173–174.
- Paiva-Cavalcanti, M., Dantas-Torres, F., Gonçalves, S.C., Morais, R.C.S., Brito, M.E.F., Otranto, D., Brandão-Filho, S.P. Quantitative real time PCR assays for the detection of *Leishmania (Viannia) braziliensis* in animals and humans. Mol. Cel. Probes., in press.
- Solano-Gallego, L., Rossi, L., Scroccaro, A.M., Montarsi, F., Caldin, M., Furlanello, T., Trotta, M., 2012. Detection of *Leishmania infantum* DNA mainly in *Rhipicephalus sanguineus* male ticks removed from dogs living in endemic areas of canine leishmaniasis. Parasites Vectors. 5, 98.