

## Life course and work ability among older adults: ELSI-Brazil

Camila Menezes Sabino de Castro<sup>I</sup>, Maria Fernanda Lima-Costa<sup>II,IV</sup>, Cibele Comini César<sup>II,IV</sup>, Jorge Alexandre Barbosa Neves<sup>III</sup>, Fabíola Bof de Andrade<sup>II,IV</sup>, Paulo Roberto Borges de Souza Junior<sup>V</sup>, Rosana Ferreira Sampaio<sup>VI</sup>

<sup>I</sup> Fundação Oswaldo Cruz. Instituto René Rachou. Programa de Pós-graduação em Saúde Coletiva. Belo Horizonte, MG, Brasil

<sup>II</sup> Universidade Federal de Minas Gerais. Faculdade de Ciências Econômicas. Centro de Desenvolvimento e Planejamento Regional. Belo Horizonte, MG, Brasil

<sup>III</sup> Universidade Federal de Minas Gerais. Faculdade de Filosofia e Ciências Humanas. Departamento de Sociologia. Belo Horizonte, MG, Brasil

<sup>IV</sup> Fundação Oswaldo Cruz. Instituto René Rachou. Núcleo de Estudos em Saúde Pública e Envelhecimento. Belo Horizonte, MG, Brasil

<sup>V</sup> Fundação Oswaldo Cruz. Instituto de Comunicação e Informação Científica e Tecnológica em Saúde. Rio de Janeiro, RJ, Brasil

<sup>VI</sup> Universidade Federal de Minas Gerais. Escola de Educação Física, Fisioterapia e Terapia Ocupacional. Programa de Pós-graduação em Ciências da Reabilitação. Belo Horizonte, MG, Brasil

### ABSTRACT

**OBJECTIVE:** To examine factors associated with perception of work ability in a nationally representative sample of Brazilians aged 50 years and over.

**METHODS:** We used data from 8,903 participants of the baseline survey of the Brazilian Longitudinal Study of Aging (ELSI-Brazil). The dependent variable was self-rated work ability (good or very good *versus* fair, poor, or very poor). Independent variables included factors that operate at the beginning, middle, and current stage of life. Multivariate analysis was based on prevalence ratios (PR) and 95% confidence intervals (95%CI) estimated by Poisson regression.

**RESULTS:** Good work ability was reported by 49% of participants (49.4% among men and 48.6% among women). Results of the multivariate analysis showed that, for both men and women, good work ability showed positive and statistically significant associations ( $p < 0.05$ ) with good health up to 15 years of age (PR = 1.22 and 1.18, respectively), educational level  $\geq 8$  years (PR = 1.19 and 1.21, respectively), and current good self-rated health (PR = 1.88 and 1.94, respectively). Negative associations were observed for current age (PR = 0.99 for each increase of one year among men and women), medical diagnosis of depression (PR = 0.70 for men and PR = 0.87 for women), and having one or more at least chronic diseases (PR = 0.88 for men and 0.91 for women). Only for men, positive associations for the age at which they started working (PR = 1.14 and 1.12 for 11–17 and  $\geq 18$  years, respectively) and living with a spouse (PR = 1.09) were found.

**CONCLUSIONS:** Work ability in older ages is built over the life course, particularly by the health conditions in childhood and adolescence, age at which men begin working, educational level, and health conditions in older ages. Policies aimed at increasing longevity in the labor market must take these factors into account.

**DESCRIPTORS:** Aged. Work Capacity Evaluation. Health. Chronic Disease. Socioeconomic Factors. Epidemiologic Factors. Health Surveys.

#### Correspondence:

Maria Fernanda Lima-Costa  
Av. Augusto de Lima, 1715 Barro Preto  
30190-002 Belo Horizonte, MG, Brasil  
E-mail: lima-costa@cpqrr.fiocruz.br

Received: Jan 2, 2018

Approved: Apr 18, 2018

**How to cite:** Castro CMS, Lima-Costa MF, César CC, Neves JAB, Bof de Andrade F, Souza-Junior PRB, et al. Life course and work ability among older adults: ELSI-Brazil. Rev Saude Publica. 2018;52 Suppl 2:11s

**Copyright:** This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided that the original author and source are credited.



## INTRODUCTION

Aging of the workforce is a global phenomenon. World demographic estimates showed that there were seven persons of active age (20-64 years) for each individual aged 65 years and over in 2015; projections indicate that this ratio will fall to 3.5 in 2050<sup>1</sup>. In addition to this, increased life expectancy means that retired persons use the pension system for a longer time. In Brazil, these changes have been occurring rapidly and they tend to increase. In view of this, a measure adopted by different countries and under discussion in Brazil is the reformulation of social security policies, aiming at extending work life<sup>1-3</sup>. Thus, there is an increasing interest to know the determinants of staying in the work force in older ages.

Good work ability allows older persons to keep working<sup>4</sup>. The work ability demands a balance between personal resources (age, health, skills, values, and attitudes) and working conditions (work environment, demands, and organization)<sup>5</sup>. More recently, this concept has incorporated the family and the close community as factors that can affect work ability in different ways over the life course<sup>6,7</sup>. There is a consensus that the ideal balance is dynamic, and changes may occur at different stages of productive life<sup>5,8</sup>. A longitudinal study conducted in Finland<sup>9</sup> has shown that a perception of good or excellent work ability was associated with the age of retirement, which extended the stay in the workforce in approximately six years, and improved quality of life up to five years after leaving the labor market.

Chronological aging compromises the work ability<sup>6</sup>. The first decline occurs approximately at 45 years and the second, at 55<sup>10</sup>. A study conducted with workers in Finland has shown that work ability decreases by approximately 1.5% per year after 45 years of age<sup>11</sup>. Among persons of all ages, health, functional ability, and the characteristics of the work itself are the factors that most affect work ability<sup>4,7,8</sup>. Research on this subject also indicates that sex inequality worsens the level of work ability among women<sup>12,13</sup>.

Throughout life, work ability can vary from excellent to very poor<sup>10,14</sup>. Epidemiological studies from a life course perspective show that persons are constantly exposed to several factors at different stages of life. These factors can accumulate over time and impact the health and work ability at older ages<sup>15,16</sup>.

The few Brazilian studies that have examined the work ability of older persons were restricted to specific occupations<sup>12,17,18</sup>. The objective of this study was to examine factors associated with the perception of work ability, from a life course perspective, in the Brazilian population aged 50 years and over.

## METHODS

### Data Source and Sample

In this study, we used baseline data from the Brazilian Longitudinal Study of Aging (ELSI-Brazil), whose sample was designed to be representative of the non-institutionalized Brazilian population aged 50 years and over. The baseline data collection was conducted between 2015 and 2016. Sampling used a design with stages of selection, which combined stratification of primary units (municipalities), census tracts, and households. All residents of selected households aged 50 and over were eligible for interview and other procedures. The final sample was estimated at 10,000 persons residents in 70 municipalities from different Brazilian regions (9,412 participated). More details can be found on the homepage of the ELSI-Brazil<sup>a</sup> and in a previous publication<sup>19</sup>.

### Variables of the Study

The dependent variable of this study was the persons's perception of their work ability. The information was obtained by the following question: "How do you evaluate your current


<sup>a</sup>Fundação Oswaldo Cruz. Brazilian Longitudinal Study of Aging. Rio de Janeiro; c2015 [cited 2017 Nov 28]. Available from: <http://elsi.cpqrr.fiocruz.br>

work ability?” With five possible answers: very good, good, fair, poor, and very poor. For the analysis, the answers were categorized as good (good or very good) and poor (fair, poor, or very poor).

The independent variables were defined following a life course perspective, which considers exposure factors that operate at the beginning, middle, and current stage of life. The variables were temporally divided into distal (related to the characteristics at the beginning of life), intermediate, and proximal (current characteristics). We considered the following distal variables: (1) educational status at the age of 10, whose information was obtained by the question: “When you were 10 years old, did you go to school?”; (2) family financial situation up to 15 years of age, defined by the answer to the question: “Considering your childhood, since your birth to 15 years of age, you would say that your family...”, which had the following response options: above or within average, poor, variable financial situation; (3) health condition up to 15 years of age, which was measured by the question: “Would you say that your health, since your birth to 15 years of age, was...”, which had the following response options: good (good, very good, or excellent) or poor (fair or poor); and (4) age at which the person started working (10 years or less, 11 to 17 years, and 18 years and over).

Conditions in the middle stages of life were defined by the following variables: (1) physical demands of the work, defined by the answer to the question, “How would you describe the physical demands of the work you had for the most part of your life?”, which was categorized as: never worked or was seated most of the time; was standing or walking, or the work required some physical effort; and the work required intense physical effort; (2) number of complete years of education, categorized as 0–7 years *versus* 8 years or more.

Proximal or current conditions were: (1) age at the date of the interview, (2) position in the household (responsible for the household or not), (3) living arrangements (living or not with a spouse or partner), and (4) current health conditions. Health conditions included self-rated health, defined by the answer to the question, “In general, how would you evaluate your health?”, and chronic diseases, defined by the answer to the question “Has a doctor ever told you that you have “this disease?” In this analysis, self-rated health was categorized as good (good or very good) or poor (fair, poor, or very



Numbers of respondents = 8,903 (3,942 men and 4,961 women)

**Figure 1.** Prevalence of good self-rated work ability according to sex and age in the population aged 50 years and over. Brazilian Longitudinal Study of Aging (ELSI-Brazil), 2015–2016.

poor). The diseases included depression (analyzed separately) and number of chronic diseases, considering hypertension, diabetes, high cholesterol level, cerebrovascular disease, asthma, arthritis or rheumatism, osteoporosis, chronic back pain, cancer, and heart disease. In this analysis, we categorized the answers into none and one or more diseases.

### Data Analysis

In the description of the characteristics of the study participants, the analysis of the differences between men and women was based on Pearson's chi-square test and Student's t-test to examine the differences between frequencies and means, respectively. The multivariate analysis was based on prevalence ratios (PR) and respective 95% confidence intervals (95%CI), estimated by robust Poisson regression. We fitted the results to illustrate the prevalence ratios of the variables associated with work ability based on the final multivariate model (Figure 2). Collinearity between the study variables was analyzed using the variance inflation factor ( $VIF < 5$ ). All variables simultaneously entered the multivariate models, since no collinearity was identified.

The unadjusted and multivariate analyses were performed separately for men and women. In all estimates, we used the procedures for complex samples from the statistical program Stata 14.1 (College Station, Texas, USA) in order to consider the effect of the study design and the individuals sample weights.

**Table 1.** Distribution of good self-rated work ability according with lifecourse characteristics of the population aged 50 years and over, by sex. Brazilian Longitudinal Study of Aging (ELSI-Brazil), 2015–2016.

Variable	Total (n = 8,903)		Men (n = 3,942)		Women (n = 4,961)		p <sup>b</sup>
	% <sup>a</sup>	95%CI	% <sup>a</sup>	95%CI	% <sup>a</sup>	95%CI	
Good current work ability	49.0	46.6–51.4	49.4	46.4–52.5	48.6	46.2–51.0	0.511
Early life conditions							
Attended school at the age of 10	75.5	72.3–78.5	75.5	71.6–78.9	75.6	72.4–78.5	0.941
Financial situation of the family up to 15 years of age							
Above or within the average	27.7	25.4–30.0	26.8	24.3–29.4	28.4	25.9–31.1	< 0.001
Poor	57.3	54.9–59.6	56.0	53.2–58.8	58.4	55.7–61.1	
Variable	15.0	13.9–16.3	17.2	15.6–18.9	13.2	12.0–14.4	
Good health up to 15 years of age	78.0	76.2–79.6	78.8	76.8–80.7	77.2	75.2–79.1	0.138
Age at which started working (years)							
< 10	42.2	39.8–44.7	45.3	42.4–48.3	39.5	37.0–42.0	< 0.001
11–17	40.6	38.4–42.8	42.7	40.0–45.5	38.7	36.3–41.2	
≥ 18	17.2	15.2–19.5	12.0	10.4–13.8	21.8	19.2–24.6	
Conditions in the intermediate stages of life							
Physical demands of the previous work							
Never worked/Sitting	17.7	16.1–19.3	16.7	14.9–18.7	18.5	16.6–20.5	< 0.001
Some physical effort	63.4	60.8–66.0	55.9	52.6–59.2	70.1	67.2–72.8	
Intense physical effort	18.9	16.7–21.3	27.6	24.2–30.8	11.4	9.3–13.9	
Education level (full years)							
< 8	63.5	60.5–66.3	62.4	58.8–65.8	64.4	61.4–67.3	0.173
Current life conditions							
Age, mean (SD)	63.7 (9.9)	63.5–63.9	62.9 (9.8)	62.6–63.2	64.3 (10.0)	64.0–64.6	< 0.001
Responsible for the household	73.5	71.7–75.3	87.0	84.7–89.0	61.7	57.8–65.4	< 0.001
Living with spouse or partner	64.3	61.4–67.1	75.6	72.9–78.1	54.3	51.2–57.3	< 0.001
Good self-rated health	44.0	41.5–46.5	43.7	40.4–47.0	44.3	41.7–46.8	0.701
Medical diagnosis of depression	18.6	16.8–20.5	10.6	9.2–12.2	25.6	23.1–28.3	< 0.001
One or more chronic diseases <sup>c</sup>	82.7	81.5–83.9	76.7	75.0–78.3	88.1	86.7–89.4	< 0.001

95%CI: 95% confidence interval.

<sup>a</sup>Weighted percentage (or mean, when specified) for the sample parameters.

<sup>b</sup>Pearson's chi-square test or Student's t-test with significance level  $\leq 0.05$ , for differences between sexes.

<sup>c</sup>Medical diagnosis of hypertension, diabetes, cholesterol, cerebrovascular accident, asthma, arthritis or rheumatism, osteoporosis, chronic back pain, cancer, and heart disease.

## Ethical Issues

The ELSI-Brazil was approved by the Research Ethics Committee of the *Fundação Oswaldo Cruz*, Minas Gerais (CAAE 34649814.3.0000.5091). All participants signed the informed consent separately for each research procedure.

## RESULTS

Of the 9,412 participants of the baseline of the ELSI-Brazil, 8,903 had complete information for all variables and were included in this analysis. Most participants in this study were


**Table 2.** Results of the multivariate analysis of the association between lifecourse characteristics and good self-rated work, by sex. Brazilian Longitudinal Study of Aging (ELSI-Brazil), 2015–2016. (n = 8,903)

Variable	Good current work ability			
	Men (n = 3,942)		Women (n = 4,961)	
	% <sup>a</sup>	PR (95%CI) <sup>b</sup>	% <sup>a</sup>	PR (95%CI) <sup>b</sup>
Early life conditions				
Attended school at the age of 10				
No	39.1	1	38.5	1
Yes	52.8	1.04 (0.94–1.15)	51.8	1.06 (0.96–1.16)
Financial situation of the family up to 15 years of age				
Above/Within average	56.3	1	55.1	1
Poor	44.8	0.95 (0.88–1.02)	44.3	0.94 (0.87–1.02)
Variable	53.8	1.02 (0.93–1.12)	53.3	1.02 (0.94–1.11)
Good health up to 15 years of age				
No	36.0	1	37.8	1
Yes	53.1	1.22 (1.08–1.36)	51.8	1.18 (1.08–1.30)
Age at which started working				
≤ 10	41.7	1	42.8	1
11–17	55.4	1.14 (1.05–1.23)	48.8	1.00 (0.93–1.08)
≥ 18	57.5	1.12 (1.01–1.24)	58.7	1.07 (0.96–1.18)
Conditions in the middle stages of life				
Physical demands of the previous work				
Never worked/Sitting	58.1	1	54.4	1
Some physical effort	52.1	0.99 (0.92–1.07)	49.2	1.02 (0.94–1.10)
Intense physical effort	38.6	0.90 (0.81–1.01)	35.6	0.87 (0.73–1.04)
Education level (full years)				
0–7	40.6	1	41.2	1
≥ 8	64.0	1.19 (1.10–1.28)	61.9	1.21 (1.10–1.33)
Current life conditions				
Age (continuous)				
	61.0	0.99 (0.98–0.99)	61.7	0.99 (0.98–0.99)
Responsibility for the household				
No	46.9	1	50.2	1
Yes	49.8	1.09 (0.95–1.25)	47.6	0.97 (0.88–1.06)
Living with spouse or partner				
No	46.4	1	45.8	1
Yes	50.4	1.09 (1.02–1.17)	50.9	1.05 (0.96–1.15)
Self-rated health				
Poor	32.5	1	32.2	1
Good	71.3	1.88 (1.72–2.05)	69.2	1.94 (1.80–2.09)
Medical diagnosis of depression				
No	51.9	1	51.9	1
Yes	28.6	0.70 (0.58–0.84)	38.8	0.87 (0.80–0.95)
One or more chronic diseases <sup>c</sup>				
0	64.2	1	66.2	1
≥ 1	44.9	0.88 (0.82–0.94)	46.2	0.91 (0.85–0.97)

<sup>a</sup> Crude percentages considering all sample parameters.

<sup>b</sup> PR (95%CI): prevalence ratio (95% confidence interval) mutually adjusted for all variables listed in the table and estimated by the Poisson regression model, with robust variance.

<sup>c</sup> Medical diagnosis of hypertension, diabetes, cholesterol, cerebrovascular accident, asthma, arthritis or rheumatism, osteoporosis, chronic back pain, cancer, and heart disease.



\* Adjusted for all variables listed in the figure, in addition to school situation at the age of 10, family financial situation up to 15 years of age, physical demands of the work, and position in the household.

**Figure 2.** Prevalence ratios (PR) and 95% confidence intervals\* of the lifecourse characteristics associated with good self-rated work ability at 50 years and over. Brazilian Longitudinal Study of Aging (ELSI-Brazil), 2015–2016.

women ( $n = 4,961$ , 56%). Mean age of the sample was 63.7 years ( $SD = 9.9$ ), and it was higher among women [64.3 years ( $SD = 10.0$ )] than men [62.9 years ( $SD = 9.8$ )]. Work ability was evaluated as good by 49.4% of the men and 48.6% of the women. Perception of good work ability decreased with age (Figure 1) both among men (from 57.9% at 50–54 to 51.4% at 55–59 years, 48.5% at 60–64 years, 44.4% at 65–69 years, and 40% at 70 years and over) and among women (56.8%, 50.3%, 47.6%, 47.1%, and 39.4%, respectively).

Table 1 shows the characteristics of the study participants according to sex. Statistically significant differences between men and women were observed for the variables of family financial situation up to 15 years of age, age at which the individual began working, physical demands of the work, age, responsibility for the household, living with a spouse, medical diagnosis of depression, and medical diagnosis of one or more chronic diseases.

The results of the multivariate analysis of the association between the characteristics over the life course and work ability according to sex are shown in Table 2 and Figure 2. Among both men and women, good work ability showed positive and statistically significant associations with good health up to 15 years of age ( $PR = 1.22$  and  $1.18$ , respectively), educational level equal to or greater than eight years ( $PR = 1.19$  and  $1.21$ , respectively), and current good self-rated health ( $PR = 1.88$  and  $1.94$ , respectively). In both sexes, negative associations were observed for age ( $PR = 0.99$  for each increase of one year in age in both groups), medical diagnosis of depression ( $PR = 0.70$  for men and  $PR = 0.87$  for women), and having at least one chronic disease ( $PR = 0.88$  and  $0.91$ , respectively). Only among men, good work ability was positively associated with age at which they started working ( $PR = 1.14$  and  $1.12$  for those who started working at 11–17 years and  $^3$  18 years, respectively) and living with a spouse ( $PR = 1.09$ ).

## DISCUSSION

To our knowledge, this is the first nationally representative sample study to examine the influence of experiences over the life course on current work ability. The results show that, for both men and women, good health conditions in early life are associated with better conditions in work ability at older ages. The results also show that education level plays an important role in work ability among older adults, with better conditions among those with higher education. In relation to current conditions, the results highlight the importance of health conditions for work ability,

as those who evaluate their current health as poor, who had a medical diagnosis of depression, or who had a medical diagnosis of any other chronic disease had worse work ability.

Health conditions during childhood and adolescence are crucial for the health development of the adult and older adult<sup>15</sup>. Factors affecting health conditions in childhood and adolescence are associated with participation in school, family income, place of residence, parents' education and employment, cultural and religious aspects, among others<sup>20,21</sup>. Our results reinforce these observations by showing that, among the early life variables considered in this analysis, good health up to 15 years of age presented independent association with work ability at older ages and this association was consistent among men and women.

Another early life characteristic associated with work ability in this analysis was the age at which men began working. The Brazilian legislation allows the work of young persons from the age of 18 years and adolescents from the age of 14 years, allowing as it is carried out as employment, internship, and learning. According to a recent survey by the Brazilian Institute of Geography and Statistics (2017)<sup>22</sup>, at least 190,000 boys and girls up to 13 years of age were part of the illegal group of workers in 2016. Of these children, only 27% had paid work. In relation to the activities, 47.6% were included in agricultural work, 24.7% in the civil construction, transportation, and industry sectors, 21.4% in trade, and 6.3% in domestic services. Our results deepen these observations by showing that the higher the age a man starts working, the greater the tendency to have good work ability. This association was not observed among women. As shown by Kassouf<sup>23</sup>, the earlier the entry into the labor market, the lower the wage, the lower the education level achieved, and the worse the health condition in adult life. Women tend to enter the labor market later and this allows them to acquire a higher education level in relation to men. Mean age of entry into the labor market is lower for men than women, while variance is higher. Therefore, the association between the age when the individual started working and the tendency to have good work ability among older men (but not among women) may be due to both a statistical issue (the larger variance of the age men began working would lead to a greater association) and to the fact that men who enter the labor market earlier have lower income in adult life, and so the association is due to factors not considered in this analysis (such as income received throughout adult life).

The results of this analysis show that men and women with higher education ( $\geq 8$  years) are 20% more likely to evaluate their work ability as good. Workers with lower education levels experience more unfavorable working conditions and, consequently, are in jobs with greater physical demands, which restrict their work ability<sup>4</sup>. We emphasize that education level is the result of accumulated investments throughout life and it is related to personal factors and the characteristics of parents<sup>20</sup>. Thus, from a life-course perspective, we highlight the importance of improving the quantity and quality of education acquired by individuals, particularly among those whose parents have low education levels and who probably belong to the poorest strata of the population.

The decrease in work ability with increasing age, as observed in this study, is expected<sup>4,8,9</sup>, since older persons generally present worsening health conditions and declining functional ability, as well as difficulties in adapting to changes in work and technological advances<sup>6,24</sup>. According to Ilmarinen<sup>13</sup>, in a longitudinal study conducted in Finland, the mean score for the work ability index begins to decrease from 51 to 55 years, and this decrease is consistently observed in the different occupational groups studied. Our results indicate that, among Brazilian older adults, the assessment of work ability decreases by approximately 13%, for both sexes, between the age groups of 50–54 and 55–59 years, and it decreases progressively from the age of 60 years.

Work ability is not separated from life outside work. The family can also affect the work ability of an individual in different ways over the life course. It has become more important to balance work and family life<sup>5</sup>. Our analysis confirms the above observations by showing that living with a spouse or partner is associated with good work ability, but this association was observed only for men. Perhaps this result can be explained by the fact that men can

gain health benefits when living with women, since they are more health-oriented. Since the work of Gove<sup>25</sup>, researchers have shown that cohabitation with a spouse of the opposite sex tends to benefit the health of men more than women<sup>26</sup>.

Health is considered as one of the main determinants of work ability<sup>5,18</sup> and self-rated health is a strong predictor of early retirement<sup>27</sup>. All health conditions considered in this analysis presented statistically significant associations with work ability. Among them, self-rated health showed the strongest association. Self-rated health is a multidimensional indicator that expresses the individual's physical, mental, and social health, and it is a robust predictor of mortality among older adults<sup>28</sup>. We observed negative associations with work ability for the medical diagnosis of chronic diseases and depression, which indicates that persons with these conditions have a worse evaluation of their work ability. Mental disorders are the first cause of disability in the world and the third cause of absenteeism and disability retirement in Brazil<sup>29</sup>. Mental disorders are also significantly associated with unemployment and early retirement because of permanent or temporary incapacity<sup>30</sup>. A longitudinal study conducted for a year in the Netherlands<sup>8</sup> among individuals aged 45–64 years has shown that workers with chronic diseases have worse work ability compared to their peers without health problems. The major differences between the two groups were found for mental health problems (9.4%) and musculoskeletal problems (4.2%), while heart disease and diabetes mellitus had less impact (2.7% and 2.0% respectively). Our results are in line with these findings as they show a stronger association between depressive symptoms and work ability compared to other chronic diseases. In this case, we cannot rule out reverse causality, since depression may be the cause or the consequence of work incapacity.

This study has advantages and limitations. Among the advantages we can mention the national representative sample of Brazilian older adults. As for the limitations, the collection of self-reported data of the variables related to past experiences may have been influenced by the desire to transmit a socially and culturally acceptable image (bias of social desirability), besides being subject to information bias. Another limitation of the study refers to its cross-sectional nature, which does not allow us to establish temporal associations for the variables related to current conditions. However, it is important to note that longitudinal studies show that age, education level, and current health conditions are associated with current work ability<sup>8,9</sup>. Our results are consistent with these observations.

In summary, the results of this study show that work ability in older ages is built over the life course, particularly by the health conditions in childhood and adolescence, age at which men began working, educational level, and health conditions in older ages. Policies aimed at increasing longevity in the labor market must take these factors into account, which profoundly affect the quality of life and work.

## REFERENCES

1. United Nations, Department of Economic and Social Affairs, Population Division. World Population Ageing 2015. New York; 2015 [cited 2016 Jul 29]. Available from: <http://www.un.org/en/development/desa/population/theme/ageing/WPA2015.shtml>
2. Marques RM. Envelhecimento da população e previdência sócia: a dinâmica demográfica está diretamente relacionada às políticas para aposentadoria. *Rev Pre Univesp*. 2016-2017 [cited 2017 Nov 25];(61). Available from: <http://pre.univesp.br/previdencia-social#.WzPsU9JKjIV>
3. Zacher H, Griffin B. Work, aging, and retirement in Australia: introduction to the special issue. *Work Aging Retire*. 2015;1(2):129-32. <https://doi.org/10.1093/workar/wau011>
4. Hasselhorn HM, Apt W, editors. Understanding employment participation of older workers: creating a knowledge base for future labour market challenges. Berlin: Federal Ministry of Labour and Social Affairs; 2015 [cited 2016 Jul 29]. Available from: [https://www.baua.de/EN/Service/Publications/Cooperation/Gd81.pdf?\\_\\_blob=publicationFile&v=5](https://www.baua.de/EN/Service/Publications/Cooperation/Gd81.pdf?__blob=publicationFile&v=5)
5. Ilmarinen J. Work ability: a comprehensive concept for occupational health research and prevention [editorial]. *Scand J Work Environ Health*. 2009;35(1):1-5. <https://doi.org/10.5271/sjweh.1304>



6. Ilmarinen J. Promoting active ageing in the workplace. Bilbao: European Agency for Safety and Health at Work; 2012 [cited 2017 Nov 1]. Available from: <https://osha.europa.eu/en/tools-and-publications/publications/articles/promoting-active-ageing-in-the-workplace>
7. Gould R, Ilmarinen J, Järvisalo J, Koskinen S, editors. Dimension of work ability: results of the Health 2000 Survey. Helsinki: Finnish Centre of Pensions, The Social Insurance Institution, National Public Health Institute, Finnish Institute of Occupational Health; 2008 [cited 2017 Nov 1]. Available from: [https://www.julkari.fi/bitstream/handle/10024/78055/dimensions\\_of\\_work\\_ability\\_7.pdf?sequence=1&isAllowed=y](https://www.julkari.fi/bitstream/handle/10024/78055/dimensions_of_work_ability_7.pdf?sequence=1&isAllowed=y)
8. Leijten FRM, Heuvel SG, Ybema JF, Beek AJ, Robroek SJW, Burdorf A. The influence of chronic health problems on work ability and productivity at work: a longitudinal study among older employees. *Scand J Work Environ Health*. 2014;40(5):473-82. <https://doi.org/10.5271/sjweh.3444>
9. Feldt T, Hyvönen K, Mäkilangas A, Kinnunen U, Kokko K. Development trajectories of Finnish manager's work ability over a 10-year follow-up period. *Scand J Work Environ Health*. 2009;35(1):37-47. <https://doi.org/10.5271/sjweh.1301>
10. Sampaio RF, Augusto VG. Envelhecimento e trabalho: um desafio para a agenda da reabilitação. *Rev Bras Fisioter*. 2012;16(2):94-101. <https://doi.org/10.1590/S1413-35552012000200003>
11. Tuomi K, Ilmarinen J, Martikainen R, Aalto L, Klockars M. Aging, work, life-style and work ability among Finnish municipal workers in 1981-1992. *Scand J Work Environ Health*. 1997 [cited 2017 Nov 1];23 Suppl 1:58-65. Available from: <http://www.jstor.org/stable/4096692>
12. Godinho MR, Greco RM, Teixeira MTB, Teixeira LR, Guerra MR, Chaoubah A. Work ability and associated factors of Brazilian technical-administrative workers in education. *BMC Res Notes*. 2016;9:1. <https://doi.org/10.1186/s13104-015-1837-x>
13. Ilmarinen J, Tuomi K, Klockars M. Changes in the work ability of active employees over an 11-year period. *Scand J Work Environ Health*. 1997 [cited 2017 Nov 1];23 Suppl 1:49-57. Available from: <http://www.jstor.org/stable/4096691>
14. Lindberg P, Josephson M, Alfredsson L, Vingård E. Promoting excellent work ability and preventing poor work ability: the same determinants? Results from the Swedish HAKuL study. *Occup Environ Med*. 2006;63(2):113-20. <https://doi.org/10.1136/oem.2005.022129>
15. Ben-Shlomo Y, Kuh D. A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. *Int JEpidemiol*. 2002;31(2):285-93. <https://doi.org/10.1093/ije/31.2.285>
16. Blane D, Higgs P, Hyde M, Wiggins RD. Life course influences on quality of life in early old age. *Soc Sci Med*. 2004;58(11):2171-9. <https://doi.org/10.1016/j.socscimed.2003.08.028>
17. Martinez MC, Latorre MRDO, Fischer FM. Capacidade para o trabalho: revisão de literatura. *Cienc Saude Coletiva*. 2010;15 Supl 1:1553-61. <https://doi.org/10.1590/S1413-81232010000700067>
18. Martinez MC, Latorre MRDO. Saúde e capacidade para o trabalho em trabalhadores de área administrativa. *Rev Saude Publica* 2006;40(5):851-8. <https://doi.org/10.1590/S0034-89102006000600015>
19. Lima-Costa MF, Andrade FB, Souza Jr PRB, Neri AL, Oliveira Duarte YA, Castro-Costa E, et al. The Brazilian Longitudinal Study of Aging (ELSI-Brazil): objectives and design. *Am JEpidemiol*. 2018;187(7):1345-53. <https://doi.org/10.1093/aje/kwx387>
20. Reis MC, Ramos L. Escolaridade dos pais, desempenho no mercado de trabalho e desigualdade de rendimentos. *Rev Bras Econ*. 2011;65(2):177-205. <https://doi.org/10.1590/S0034-71402011000200004>
21. Sucupira ACSL, Andrade LOM, Barreto ICHC, Lima JW, Santiago AV, Santiago AX. Determinantes sociais da saúde de crianças de 5 a 9 anos da zona urbana de Sobral, Ceará, Brasil. *Rev Bras Epidemiol*. 2014;17 Supl 2:160-77. <https://doi.org/10.1590/1809-4503201400060014>
22. Quase um milhão de crianças são vítimas do trabalho infantil no País, diz IBGE. *Bras Econ*. 29 nov 2017 [cited 2017 Nov 30]. Available from: <http://economia.ig.com.br/2017-11-29/ibge-trabalho-infantil.html>
23. Kassouf AL. O que conhecemos sobre o trabalho infantil? *Nova Econ*. 2007;17(2):323-50. <https://doi.org/10.1590/S0103-63512007000200005>
24. Camarano AA, Kanso S, Fernandes D. Menos jovens e mais idosos no mercado de trabalho. In: Camarano AA, organizadora. Novo regime demográfico: uma nova relação entre população e desenvolvimento? Rio de Janeiro: IPEA; 2014. Cap 12, p. 377-406.

25. Gove WR. The relationship between sex roles, marital status and mental illness. *Soc Forces*. 1972;51(1):34-44. <https://doi.org/10.2307/2576129>
26. Ploubidis GB, Silverwood RJ, DeStavola B, Grundy E. Life-course partnership status and biomarkers in midlife: evidence from the 1958 British Birth Cohort. *Am J Public Health*. 2015;105(8):1596-603. <https://doi.org/10.2105/AJPH.2015.302644>
27. Karpansalo M, Manninen P, Kauhanen J, Lakka TA, Salonen JT. Perceived health as a predictor of early retirement. *Scand J Work Environ Health*. 2004;30(4):287-92. <https://doi.org/10.5271/sjweh.796>
28. Lima-Costa MF, Cesar CC, Chor D, Proietti FA. Self-rated health compared with objectively measured health status as a tool for mortality risk screening in older adults: 10-year follow-up of the Bambuí Cohort Study of Aging. *Am J Epidemiol*. 2012;175(3):228-35. <https://doi.org/10.1093/aje/kwr290>
29. França MH, Barreto SM, Pereira FG, Andrade LHSG, Paiva MCA, Viana MC. Mental disorders and employment status in the São Paulo Metropolitan Area, Brazil: gender differences and use of health services. *Cad Saude Publica*. 2017;33(9):e00154116. <https://doi.org/10.1590/0102-311X00154116>
30. Segel-Karpas D. Number of illnesses, self-perceived health, and depressive symptoms: the moderating role of employment in older adulthood and old age. *Work, Aging Retire*. 2015;1(4):382-92. <https://doi.org/10.1093/workar/wav009>

---

**Funding:** The ELSI-Brazil baseline study was supported by the Brazilian Ministry of Health (DECIT/SCTIE – Department of Science and Technology from the Secretariat of Science, Technology, and Strategic Inputs (Grant 404965/2012-1); COSAPI/DAPES/SAS – Healthcare Coordination of Older Adults, Department of Strategic and Programmatic Actions from the Secretariat of Health Care (Grants 20836, 22566, and 23700) and the Brazilian Ministry of Science, Technology, Innovation, and Communication.

**Authors' Contribution:** All authors contributed equally to all phases of the study.

**Conflict of Interest:** The authors declare no conflict of interest.