

RESEARCH

Open Access

# Clinical outcomes of patients requiring ventilatory support in Brazilian intensive care units: a multicenter, prospective, cohort study

Luciano CP Azevedo<sup>1,2\*</sup>, Marcelo Park<sup>1,2</sup>, Jorge IF Salluh<sup>3</sup>, Alvaro Rea-Neto<sup>4</sup>, Vicente C Souza-Dantas<sup>5,6</sup>, Pedro Varaschin<sup>6</sup>, Mirella C Oliveira<sup>4</sup>, Paulo Fernando GMM Tierno<sup>7</sup>, Felipe dal-Pizzol<sup>8</sup>, Ulysses VA Silva<sup>9</sup>, Marcos Knibel<sup>10</sup>, Antonio P Nassar Jr<sup>11</sup>, Rossine A Alves<sup>12</sup>, Juliana C Ferreira<sup>13</sup>, Cassiano Teixeira<sup>14</sup>, Valeria Rezende<sup>15</sup>, Amadeu Martinez<sup>16</sup>, Paula M Luciano<sup>17</sup>, Guilherme Schettino<sup>1</sup> and Marcio Soares<sup>3</sup>, for The ERICC (Epidemiology of Respiratory Insufficiency in Critical Care) investigators

## Abstract

**Introduction:** Contemporary information on mechanical ventilation (MV) use in emerging countries is limited. Moreover, most epidemiological studies on ventilatory support were carried out before significant developments, such as lung protective ventilation or broader application of non-invasive ventilation (NIV). We aimed to evaluate the clinical characteristics, outcomes and risk factors for hospital mortality and failure of NIV in patients requiring ventilatory support in Brazilian intensive care units (ICU).

**Methods:** In a multicenter, prospective, cohort study, a total of 773 adult patients admitted to 45 ICUs over a two-month period requiring invasive ventilation or NIV for more than 24 hours were evaluated. Causes of ventilatory support, prior chronic health status and physiological data were assessed. Multivariate analysis was used to identify variables associated with hospital mortality and NIV failure.

**Results:** Invasive MV and NIV were used as initial ventilatory support in 622 (80%) and 151 (20%) patients. Failure with subsequent intubation occurred in 54% of NIV patients. The main reasons for ventilatory support were pneumonia (27%), neurologic disorders (19%) and non-pulmonary sepsis (12%). ICU and hospital mortality rates were 34% and 42%. Using the Berlin definition, acute respiratory distress syndrome (ARDS) was diagnosed in 31% of the patients with a hospital mortality of 52%. In the multivariate analysis, age (odds ratio (OR), 1.03; 95% confidence interval (CI), 1.01 to 1.03), comorbidities (OR, 2.30; 95% CI, 1.28 to 3.17), associated organ failures (OR, 1.12; 95% CI, 1.05 to 1.20), moderate (OR, 1.92; 95% CI, 1.10 to 3.35) to severe ARDS (OR, 2.12; 95% CI, 1.01 to 4.41), cumulative fluid balance over the first 72 h of ICU (OR, 2.44; 95% CI, 1.39 to 4.28), higher lactate (OR, 1.78; 95% CI, 1.27 to 2.50), invasive MV (OR, 2.67; 95% CI, 1.32 to 5.39) and NIV failure (OR, 3.95; 95% CI, 1.74 to 8.99) were independently associated with hospital mortality. The predictors of NIV failure were the severity of associated organ dysfunctions (OR, 1.20; 95% CI, 1.05 to 1.34), ARDS (OR, 2.31; 95% CI, 1.10 to 4.82) and positive fluid balance (OR, 2.09; 95% CI, 1.02 to 4.30).

**Conclusions:** Current mortality of ventilated patients in Brazil is elevated. Implementation of judicious fluid therapy and a watchful use and monitoring of NIV patients are potential targets to improve outcomes in this setting.

**Trial registration:** ClinicalTrials.gov NCT01268410.

\* Correspondence: luciano.azevedo@hsl.org.br

<sup>1</sup>Research and Education Institute, Hospital Sírio-Libanês, Rua Cel. Nicolau dos Santos 69 - São Paulo, 01308-060, Brazil

Full list of author information is available at the end of the article

## Introduction

Acute respiratory failure is frequent and commonly a severe organ dysfunction occurring in the intensive care unit (ICU) [1]. Under this circumstance, invasive or non-invasive mechanical ventilation (MV) are life-sustaining interventions [2]. However, despite significant advances in ventilatory support [3], it remains associated with elevated mortality [4] and a significant impairment in the patients' quality of life in the post-ICU setting [5]. Therefore, information about the epidemiological aspects of patients under MV is important from both clinical and health policy perspectives. However, most studies on the epidemiology of ventilatory support are outdated or were carried out before significant developments in the field, such as lung protective ventilation [6] or the widespread application of non-invasive mechanical ventilation (NIV) [7-9]. Moreover, these studies were usually carried out in high-income countries and very few contemporary data from emerging countries are available [10-12]. Specifically from Brazil, a previous trial evaluated the mortality of patients with acute respiratory failure. However, this study was a single center trial carried out only in a tertiary hospital and just included individuals with invasive mechanical ventilation and not patients under NIV [11]. Comprehensive information about the clinical characteristics, outcomes and risk factors for mortality of these patients is essential to assist clinicians in the decision-making process and to allow better resource allocation. Therefore, we carried out a multicenter, observational cohort study in Brazilian ICUs to describe the clinical outcomes of patients submitted to ventilatory support as well as to identify variables associated with hospital mortality.

## Materials and methods

### Design and setting

The Epidemiology of Respiratory Insufficiency in Critical Care (ERICC) study was a multicenter prospective cohort study conducted in 45 Brazilian ICUs between 1 June 2011 and 31 July 2011. The study was coordinated by the Research and Education Institute from Hospital Sírio-Libanês, São Paulo and D'Or Institute for Research and Education, Rio de Janeiro. Invitations were sent to ICUs registered at the Brazilian Research in Intensive Care Network (BRICNet) database and 45 ICUs from 12 Brazilian states agreed to participate. Investigators and centers are listed at the acknowledgements section. The study was strictly observational and decisions related to patients' care were at the discretion of the attending ICU team. The study was approved by the institutional review board (IRB) at the coordinating center (Comitê de Ética em Pesquisa - CEPesq - approval number HSL 2010/51) and, subsequently, by the National Ethics Committee and local review boards at each participating

site. The need for informed consent was waived or requested in some sites according to the local IRBs directives.

### Selection of participants and definitions

Patients aged  $\geq 18$  years old requiring ventilatory support for  $> 24$  h during the first 48 h of ICU admission at the participating ICUs were included in the study. In the subgroup of patients undergoing NIV, only those that used this modality for at least 6 h/day were included. Patients with a previous tracheostomy, admitted for routine uncomplicated postoperative care (ICU stay  $< 48$  h), readmissions and those with terminal conditions were not considered.

Demographic, clinical and laboratory data were collected during the ICU stay, including the main diagnosis for ICU admission, the reasons for and modality of ventilatory support (conventional MV or NIV), chronic health status, the Charlson Comorbidity Index [13], the need for vasopressors, dialysis, tracheostomy, the Simplified Acute Physiology Score 3 (SAPS 3) [14] and the Sequential Organ Failure Assessment (SOFA) score [15]. Patients who first received NIV, irrespective of its duration, and subsequently required endotracheal intubation were considered as NIV failure. The cumulative fluid balance over the first 72 h of ICU stay was also calculated. Sepsis was diagnosed using the current definitions [16]. The patient was considered to have an infection when there were clinical, laboratory, radiological and microbiological findings suggesting the presence of infection that justified the administration of antibiotics (excluding prophylaxis) [17]. Acute respiratory distress syndrome (ARDS) was defined and classified according to the Berlin definition [18]. The main outcome of interest in the current study was all cause in-hospital mortality.

### Data entry and processing

Data were collected using a web-based specific and standardized electronic case report form. Each investigator and research coordinator was provided access to the website, where all study documentation, including a comprehensive manual describing data collection requirements and variable definitions, was available. A central office was accessible through telephone and email contact to provide support to investigators. Local investigators were responsible for training local staff for data collection, supervising data collection, controlling for data completeness and quality.

Data consistency was assessed through a rechecking procedure of a 5% random sample of patients. Data were screened in detail by three investigators (LCA, MS, MP) for missing information, implausible and outlying values, logical errors and insufficient details. In case of

unconformity, local investigators were contacted to provide the requested information.

### Statistical analysis

Standard descriptive statistics were used to describe the study population. Continuous variables were reported as median (25% to 75% interquartile range, IQR). Univariate and multivariate analysis using a binary logistic regression were used to identify factors associated with the dependent variables (hospital mortality or NIV failure) [19]. We also carried out analysis of SOFA score excluding the respiratory component to reduce interaction with ARDS in the multivariate analysis and to assess the severity of associated organ failures. Linearity between each continuous variable and the dependent variable was demonstrated using locally weighted scatterplot smoothing (LOWESS) [19]. In case of nonlinearity, the variable was transformed or stratified according to the analysis of the functional form and clinical significance. For categorical variables with multiple levels, the reference level was attributed to the one with the lowest probability of the dependent variable. Variables yielding  $P$ -values  $< 0.2$  by univariate analysis were entered in the multivariate analysis to estimate the independent association of each covariate with the dependent variable. Results were summarized as odds ratios (OR) and respective 95% confidence intervals (CI). Possible interactions were tested. Two-tailed  $P$ -values  $< 0.05$  were considered statistically significant.

## Results

### Characteristics of study population

The study flowchart is shown in Figure 1. A total of 773 patients fulfilled the eligibility criteria of the study and were therefore evaluated. Their main characteristics are depicted in Table 1. The most frequent diagnoses at ICU admission were pneumonia (27%), neurological diseases (19%), non-pulmonary sepsis (12%) and obstructive pulmonary disease (6%). Patients were admitted to the ICU at a median of one (zero to three) day after hospital admission. Median SAPS 3 score was 62 (52 to 72) points and the probability of death estimated by the global equation was  $40 \pm 24\%$ . Using the customized equation for countries from Central and South America, the probability of death estimated by SAPS 3 was  $52 \pm 26\%$ . Most patients (67%) used vasopressors during their stay in the ICU and 19% required renal replacement therapy (RRT).

### Ventilatory support

Invasive MV was initially used in 80% ( $n = 622$ ) of the patients and NIV was used in the remaining 20% ( $n = 151$ ) of the patients as the initial ventilatory support (Table 1 and Figure 2). Of the later, 81 (54%) patients

failed NIV support and were subsequently intubated for invasive MV. Ventilatory modes used initially in patients who received invasive MV were pressure-controlled ventilation ( $n = 371$ , 60%), volume-controlled ventilation ( $n = 186$ , 30%), pressure-support ventilation ( $n = 54$ , 9%) and others ( $n = 11$ , 1%). Median tidal volume was 7.5 (6.1 to 8.7) mL/kg of predicted body weight and plateau pressures were below 30 cmH<sub>2</sub>O in the vast majority of the patients.

### Outcome analysis


The overall ICU and hospital mortality rates were 34% and 42%, respectively (Figure 2 and Table 1). In the univariate analysis, age, ideal body weight, SOFA score at day 1, SAPS 3 score, Charlson comorbidity index, hospital length of stay before ICU, admission from the emergency room and from the operating room were associated with hospital mortality. Additionally, NIV failure, lower PaO<sub>2</sub>/FiO<sub>2</sub> ratio, ARDS diagnosis, tracheostomy, duration of ventilatory support, need for vasopressors and renal replacement therapy (RRT), cumulative fluid balance and maximal blood lactate concentrations were also associated with hospital mortality (Table 1). In multivariate analysis, older age, higher SOFA scores (without respiratory component at Day 1), Charlson comorbidity index  $> 2$ , moderate to severe ARDS, NIV failure, use of invasive MV, higher lactate concentrations and both very negative or positive cumulative fluid balance over the first 72 hours of ICU stay were independently associated with increased hospital mortality (Table 2).

### ARDS diagnosis according to the Berlin definition

ARDS was diagnosed in 242 (31%) patients (Figure 2). Of these, 77% were supported with invasive MV and 23% received NIV as the initial ventilatory support. The rate of NIV failure in ARDS patients was 69%, as compared to 45% in non-ARDS patients ( $P = 0.007$ ). ICU and hospital mortality in the ARDS population was 46% and 52%, respectively (Figure 2). In Figure 3, we depicted the ICU and hospital mortality rates for each category of ARDS. The combined ICU mortality for ARDS moderate and severe (the former definition of ARDS [20]) was 55% and the hospital mortality was 60%.

### Non-invasive ventilation characteristics and failure

The characteristics of patients that initially received NIV are shown in Table 3. The most common diagnoses were pneumonia (23%), neurologic disorders (21%) and non-pulmonary sepsis (12%). Classical indications for NIV, such as obstructive pulmonary disease and congestive heart failure, were present in only 5% and 8% of the cases, respectively. NIV failure occurred in 54% (81/151) of patients receiving NIV initially. Factors related to NIV failure in univariate analysis were total SOFA score,



SOFA score excluding respiratory component, ARDS diagnosis, length of NIV, tracheostomy, use of vasopressors and a positive cumulative fluid balance. As expected, ICU and hospital lengths of stay and mortality were higher in patients who experienced NIV failure (Table 3).

In multivariate analysis, a SOFA score without the respiratory component  $\geq 4$  points, a diagnosis of ARDS and a cumulative fluid balance higher than 2 L in the first 72 hours of ICU stay were associated with NIV failure (Table 3). The frequency of NIV failure as well as

**Table 1 Patients' characteristics and univariate analysis of factors associated with hospital mortality**

Characteristic	All patients (n = 773)	Survivors (n = 451)	Non-survivors (n = 322)	OR (CI 95%)	P-value
<b>General</b>					
Age - yo	62 (43 to 76)	58 (35 to 72)	69 (53 to 80)	1.03 (1.02 to 1.04)	< 0.001
Male gender - n (%)	435 (56)	260 (58)	175 (54)	0.87 (0.66 to 1.17)	0.362
Ideal weight - Kg	61 (52 to 69)	62 (54 to 71)	61 (52 to 67)	0.98 (0.97 to 0.99)	0.034
Admission SAPS 3 score (points)	62 (52 to 72)	58 (49 to 67)	68 (57 to 77)	1.04 (1.03 to 1.06)	< 0.001
SOFA score on Day 1 (points)	8 (5 to 10)	7 (4 to 9)	8 (7 to 11)	1.17 (1.12 to 1.23)	< 0.001
SOFA score excluding respiratory component (points)	6 (4 to 8)	5 (3 to 8)	7 (5 to 9)	1.18 (1.13 to 1.24)	< 0.001
Charlson Comorbidity Index (points)					
0	314 (41)	227 (50)	87 (27)	Ref.	
1 to 2	288 (37)	160 (36)	128 (40)	0.23 (0.15 to 0.34)	< 0.001
> 2	171 (22)	64 (14)	107 (33)	0.48 (0.33 to 0.70)	
LOS prior ICU - days	1 (0 to 3)	1 (0 to 2)	1 (0 to 4)	1.03 (1.01 to 1.05)	0.009
<b>Admission source</b>					
Ward - n (%)	242 (31)	119 (26)	122 (38)	Ref.	
Emergency room - n (%)	367 (47)	225 (50)	142 (44)	0.62 (0.45 to 0.86)	0.004
Operation room - n (%)	164 (22)	107 (24)	58 (18)	0.54 (0.36 to 0.81)	0.003
<b>Admission Diagnosis</b>					
Pneumonia - n (%)	207 (27)	111 (25)	96 (30)	0.63 (0.32 to 1.23)	0.177
Neurological - n (%)	146 (19)	73 (16)	73 (23)	0.70 (0.41 to 1.22)	0.210
Non-pulmonary sepsis - n (%)	90 (12)	41 (9)	49 (15)	0.77 (0.49 to 1.19)	0.238
Asthma/COPD - n (%)	50 (6)	33 (7)	17 (5)	Ref.	
Cardiogenic pulmonary edema - n (%)	43 (6)	24 (5)	19 (6)	0.86 (0.42 to 1.75)	0.681
Extracranial trauma - n (%)	41 (5)	24 (5)	17 (5)	1.22 (0.76 to 1.95)	0.415
Hypovolemic/cardiogenic shock - n (%)	37 (5)	23 (5)	14 (4)	0.64 (0.27 to 1.55)	0.326
Aspirative syndromes - n (%)	26 (3)	17 (4)	9 (4)	0.96 (0.48 to 1.92)	0.915
Others - n (%)	133 (17)	73 (16)	60 (18)	0.74 (0.34 to 1.56)	0.431
<b>Comorbidities</b>					
Hypertension - n (%)	328 (42)	167 (37)	161 (50)	1.71 (1.28 to 2.29)	< 0.001
Diabetes - n (%)	177 (23)	81 (18)	96 (30)	1.95 (1.39 to 2.73)	< 0.001
Heart failure - n (%)	67 (9)	27 (6)	40 (12)	2.23 (1.34 to 3.73)	0.002
Chronic renal failure - n (%)	51 (7)	16 (4)	35 (11)	3.32 (1.80 to 6.10)	< 0.001
Chronic renal failure in dialysis n (%)	15 (2)	7 (2)	8 (2)	1.62 (0.58 to 4.50)	0.359
Neoplasm - n (%)	162 (21)	66 (15)	96 (30)	1.91 (1.38 to 2.65)	< 0.001
AIDS - n (%)	13 (2)	7 (2)	6 (2)	1.21 (0.40 to 3.63)	0.736
COPD - n (%)	80 (10)	47 (10)	33 (10)	0.98 (0.61 to 1.57)	0.938
<b>Ventilatory support category</b>					
Successful NIV - n (%)	70 (9)	57 (13)	13 (4)	Ref.	
NIV failure - n (%)	81 (10)	40 (8)	41 (13)	4.49 (2.14 to 9.45)	< 0.001
Invasive MV - n (%)	622 (80)	354 (78)	268 (82)	3.32 (1.78 to 6.19)	< 0.001
<b>Respiratory Data at first day</b>					
Pressure controlled mode - n (%)	371 (48)	208 (34)	163 (27)	1.09 (0.79 to 1.51)	0.603
Volume controlled mode - n (%)	186 (24)	104 (17)	82 (13)	1.06 (0.75 to 1.49)	0.742
Pressure support mode - n (%)	54 (7)	34 (6)	20 (3)	0.76 (0.42 to 1.35)	0.349
Tidal volume (ideal weight) - mL/kg	7.5 (6.1 to 8.7)	7.3 (6.0 to 8.7)	7.5 (6.6 to 8.9)	1.00 (0.90 to 1.11)	0.992
Plateau pressure - cmH <sub>2</sub> O	20 (17 to 24)	20 (17 to 24)	20 (16 to 24)	0.99 (0.95 to 1.04)	0.746
PEEP - cmH <sub>2</sub> O	6 (5 to 8)	6 (5,8)	6 (5 to 8)	0.95 (0.87 to 1.03)	0.206

**Table 1 Patients' characteristics and univariate analysis of factors associated with hospital mortality (Continued)**

PaO <sub>2</sub> /FiO <sub>2</sub> ratio	257 (185 to 330)	260 (196 to 346)	250 (167 to 316)	0.99 (0.99 to 1.00)	0.014
ARDS - n (%)	242 (31)	116 (26)	126 (39)	1.50 (1.06 to 2.12)	0.021
Length of ventilatory support - days	5 (3 to 10)	5 (2 to 8)	6 (3 to 12)	1.02 (1.01 to 1.04)	0.005
<b>Support during ICU stay</b>					
RRT - n (%)	148 (19)	47 (10)	101 (31)	3.91 (2.67 to 5.75)	< 0.001
Vasopressors - n (%)	521 (67)	265 (60)	256 (79)	2.83 (2.02 to 3.98)	< 0.001
Cumulative fluid balance (72 h) - L	2.9 (0.8 to 5.4)	2.6 (0.8 to 4.6)	3.7 (1.1 to 6.2)	1.05 (1.01 to 1.10)	0.011
<b>Outcomes</b>					
ICU mortality - n (%)	260 (34)	-----	-----	-----	-----
Hospital mortality - n (%)	322 (42)	-----	-----	-----	-----
ICU LOS - days	10 (6 to 18)	10 (6 to 17)	10 (5 to 19)	1.01 (1.00 to 1.02)	0.047
Hospital LOS - days	20 (11 to 34)	23 (14 to 36)	16 (7 to 30)	0.99 (0.98 to 0.99)	0.009

Abbreviations: AIDS, Acquired Immunodeficiency Syndrome; ARDS, Acute Respiratory Distress Syndrome; COPD, Chronic Obstructive Pulmonary Disease; ICU, Intensive care unit; LOS, Length-of-stay; MV, Mechanical ventilation; NIV, Non-invasive ventilation; PEEP, Positive End-Expiratory Pressure; RRT, Renal Replacement Therapy; SAPS, Simplified Acute Physiology Score; SOFA, Sequential Organ Failure Assessment.

hospital mortality increased significantly with the number of these risk factors presented by the patients (Figure 4).

#### Weaning and tracheostomy

Table 4 depicts the variables related to weaning in our population. Carrying out a spontaneous breathing trial and a successful extubation were protective factors for mortality in the univariate analysis. Additionally, 30% of the patients that were successfully extubated received non-invasive ventilation after extubation. Weaning failure with subsequent reintubation occurred in 15% of the patients. Tracheostomy was carried out in 182 (29%) patients under invasive mechanical ventilation 7 (5 to 11) days after endotracheal intubation.

#### Discussion


In the present study, mortality rates of patients in Brazilian ICUs requiring ventilatory support were elevated, regardless of the underlying condition. Factors such as age, comorbidities, ARDS, disease severity and variables related to ICU support like positive fluid balance and NIV failure are independently associated with hospital mortality. We also observed that more than half of the patients receiving NIV as the primary modality of ventilatory support failed and required invasive mechanical ventilation subsequently. Variables independently associated with NIV failure were the severity of organ dysfunctions, the presence of ARDS and a positive fluid balance.

The mortality rate of critically ill patients under ventilatory support and patients with ARDS is elevated in both observational and interventional studies [6,21,22]. In the last decade, however, important ventilatory interventions, such as lung protective strategies with reduction in tidal volumes [3] and widespread use of NIV [23], were more

frequently incorporated in the clinical practice [24] and could have resulted in different mortality rates. Nevertheless, some recent studies showed very modest or no changes in these outcomes [2,4,9,25]. In a systematic review by Phua *et al.*, the pooled mortality rate of ARDS in observational studies was 48% and did not decrease significantly in the last years [26]. More recently, Villar *et al.* reported a hospital mortality rate of 48% for ARDS patients under low tidal volume ventilation [4]. We observed a higher mortality rate for the entire cohort and a more prominent rate for ARDS patients. However, our results are within the predicted mortality range of SAPS 3 and comparable to those reported in similar countries, such as Argentina [27]. Possible explanations for our findings may include unequal access to healthcare [28,29] as well as unmeasured factors related to the process of caring for these patients.

There is a significant gap between the recommendations of low tidal volumes for ARDS patients and their adoption in practice. Several observational studies demonstrated the lack of adherence to this strategy [9,30] and our study confirms these findings. Moreover, a recent meta-analysis suggests that even ventilated patients without ARDS may benefit from low tidal volumes [31]. However, it is important also to emphasize that in this trial and similar to other studies, the majority of patients were ventilated with plateau pressures below the limit of 30 cmH<sub>2</sub>O, which may partially compensate the harmful effects of high tidal volumes.

In our study, more than 60% of the patients under invasive MV for more than 24 hours were submitted to a spontaneous breathing trial and 54% were extubated, which is similar to previous reports [32]. Seventeen percent of the patients used NIV after extubation, an incidence also



comparable to other studies [33]. Tracheostomy was done in 29% percent of the patients in a median period of one week after initiation of MV. There is significant heterogeneity in the rates of tracheostomy in patients under MV as well as at the time of the procedure [9,34-36]. A previous one-day point-prevalence study of MV that included patients from Brazil showed that tracheostomy was done in 27% of the individuals in a median of 8 (1 to 15) days after ventilatory support [12]. These numbers are similar to our report and to comparable countries such as Argentina, Chile and Uruguay [12].

NIV was used as the first line of treatment for respiratory failure in 20% of the patients in our population, with a 54% failure rate. The failure rates for NIV are quite variable in the literature and seem to be related to the cause of respiratory failure and disease severity [23,37,38]. Elevated failure rates are worrisome since NIV failure has been previously associated with increased mortality risk [39] and, in this study, was an independent risk factor for mortality. We could speculate that misperception of

disease severity by the multidisciplinary team may have contributed to over-utilization of NIV for high-risk patients, delaying invasive mechanical ventilation and contributing to the poor outcome of these patients.


Another example of a potentially modifiable risk factor for mortality is related to the fluid strategy. A positive fluid balance is consistently associated with adverse outcomes in the ICU setting, mainly for patients with ARDS [40,41] and acute kidney injury [42]. We found that the extremes of cumulative fluid balance in the first three days are independently associated with hospital mortality. Interestingly, we also found that a positive fluid balance in the first days was associated with NIV failure. Our data suggest that, taking into account the hemodynamic status, a judicious fluid balance in the first days of ICU stay may be a safer goal in patients under ventilatory support.

The present study has several shortcomings. First, it was carried out during the winter period in the Southern hemisphere, and this may have influenced the incidence of respiratory infections and also the occurrence of ARDS.

**Table 2 Factors associated with hospital mortality in a multivariate analysis**

Parameter	OR	CI 95%	P-value
Age	1.03	1.01 to 1.03	< 0.001
SOFA Score excluding respiratory component	1.12	1.05 to 1.19	< 0.001
Charlson comorbidity index			0.002
0	ref	ref	
1 to 2	1.37	0.92 to 2.04	0.124
> 2	2.30	1.28 to 3.17	< 0.001
ARDS			0.037
Absence ARDS	ref	ref	
ARDS mild	1.09	0.68 to 1.73	0.721
ARDS moderate	1.92	1.11 to 3.35	0.02
ARDS severe	2.12	1.02 to 4.41	0.045
Mechanical ventilation			0.004
Successful NIV	ref	ref	
NIV failure	3.96	1.74 to 8.99	0.001
Invasive MV	2.67	1.32 to 5.39	0.006
Cumulative fluid balance (72 h)			0.005
< -1.5 L	3.08	1.47 to 6.48	0.003
-1.5 L to +1.5 L	ref	ref	
> +1.5 L to +5.0 L	1.84	1.11 to 3.04	0.018
> 5.0 L	2.44	1.39 to 4.28	0.002
Lactate levels (Ln + 0.5)	1.78	1.27 to 2.51	0.001

Abbreviations: ARDS, Acute Respiratory Distress Syndrome; MV, Mechanical ventilation; NIV, Non-invasive ventilation; SOFA, Sequential Organ Failure Assessment Score. Area under receiver operating curve for predicted mortality (95% CI): 0.77 (0.73 to 0.81),  $P < 0.001$ . Hosmer-Lemeshow  $\chi^2$ : 8.535,  $P = 0.383$ .






**Table 3 Factors associated with NIV failure on univariate and multivariate analysis**

Characteristic	Univariate comparison			Logistic regression	
	NIV success (n = 70)	NIV failure (n = 81)	P-value (univariate)	OR (CI 95%)	P-value (multivariate)
<b>General</b>					
Age - yo	63 (48 to 80)	67 (43 to 78)	0.826		
Male gender - n (%)	34 (49)	45 (56)	0.392		
Admission SAPS 3 Score (points)	59 (50 to 68)	62 (55 to 69)	0.180		
SOFA Score on Day 1 (points)	5 (3 to 7)	7 (4 to 8)	< 0.001		
SOFA Score excluding respiratory component (points)	3 (1 to 5)	4 (4 to 7)	0.001		
SOFA excluding respiratory component ≥ 4 points	34 (48)	65 (80)	< 0.001	1.20 (1.05 to 1.34)	0.009
Charlson Comorbidity Index (points)					
0	21 (30)	25 (31)	Ref		
1 to 2	32 (46)	38 (47)	0.794		
> 2	17 (24)	18 (22)	0.782		
<b>Admission diagnoses</b>					
Pneumonia - n (%)	17 (24)	18 (22)	0.562		
Neurological - n (%)	13 (19)	19 (23)			
Non-pulmonary sepsis - n (%)	9 (13)	10 (12)			
Asthma/COPD - n (%)	4 (6)	3 (4)			
Cardiogenic pulmonary edema - n (%)	5 (7)	7 (9)			
Extracranial trauma - n (%)	4 (6)	2 (3)			
Hypovolemic/cardiogenic shock - n (%)	5 (7)	5 (6)			
Aspirative syndromes - n (%)	4 (7)	3 (4)			
Others - n (%)	9 (13)	14 (17)			
<b>Respiratory data</b>					
Inspiratory pressure first day - cmH <sub>2</sub> O	12 (10 to 15)	12 (11 to 15)	0.956		
Expiratory pressure first day - cmH <sub>2</sub> O	8 (8 to 10)	8 (6 to 10)	0.891		
PaO <sub>2</sub> /FiO <sub>2</sub> ratio first day	230 (185 to 300)	200 (150 to 300)	0.271		
ARDS - n (%)	17 (31)	38 (69)	0.004	2.31 (1.10 to 4.82)	0.026
Length of NIV - days	3 (2 to 4)	1 (0 to 2)	< 0.001		
Day of NIV failure	—	1 (0 to 2)	—		
Tracheostomy - n (%)	2 (3)	16 (20)	0.007		
<b>ICU variables</b>					
Vasopressors - n (%)	16 (23)	50 (62)	< 0.001		
Cumulative fluid balance ≥ 2 L (72 h) - n (%)	36 (51)	59 (73)	0.007	2.09 (1.02 to 4.30)	0.045
<b>Outcome</b>					
ICU mortality - n (%)	7 (10)	38 (47)	< 0.001		
Hospital Mortality - n (%)	13 (19)	41 (51)	< 0.001		
ICU LOS - days	7 (4 to 8)	13 (7 to 23)	< 0.001		
Hospital LOS - days	14 (9 to 26)	21 (13 to 30)	0.071		

Data are expressed as median (p25 to p75) or number (%). Abbreviations: ARDS, Acute Respiratory Distress Syndrome; COPD, Chronic Obstructive Pulmonary Disease; LOS, Length-of-stay; NIV, Non-invasive ventilation; RRT, Renal Replacement Therapy; SAPS, Simplified Acute Physiology Score; SOFA, Sequential Organ Failure Assessment. Area under receiver operating curve for NIV failure (95% CI): 0.72 (0.64 to 0.80),  $P < 0.001$ . Hosmer-Lemeshow  $\chi^2$ : 2.555,  $P = 0.768$ .

However, the study was conducted between early June and the end of July and epidemiologic data demonstrate that both Influenza and other lower respiratory infections present a different seasonality varying according to the region of the country [43]. We did not collect data on the origin and on the time of ARDS development, which precludes a more detailed evaluation of these patients. In addition, we used a convenience sample of ICUs that usually participate

in clinical studies in Brazil and the sites that included patients are predominantly from the Southeast and Southern regions of the country, thus our data may not be representative of the entire nation. However, the concentration of hospitals from these regions is in accordance with a higher concentration of ICU beds and the health-care system in Brazil [44]. Considering that academic institutions and those participating in clinical studies usually



**Table 4 Weaning variables of patients under invasive mechanical ventilation**

Characteristic	All patients (n = 622)	Survivors (n = 354)	Non-survivors (n = 268)	OR (CI 95%)	P-value
<b>General variables - n (%)</b>					
Successful spontaneous breathing trial	378 (61)	304 (86)	74 (28)	0.06 (0.04 to 0.09)	< 0.001
Extubation	338 (54)	270 (76)	68 (25)	0.11 (0.08 to 0.16)	< 0.001
Use of NIV after extubation	108 (17)	77 (22)	31 (12)	0.52 (0.34 to 0.83)	0.006
Reintubation	94 (15)	57 (16)	37 (14)	0.94 (0.59 to 1.47)	0.792
Time to reintubation	1 (0 to 3)	1 (0 to 3)	1 (0 to 3)	1.00 (0.94 to 1.07)	0.837
Time < 48 hours to reintubation	57 (9)	30 (8)	27 (10)	0.74 (0.38 to 1.47)	0.400
Tracheostomy	182 (29)	93 (26)	89 (33)	1.37 (0.97 to 1.39)	0.073
Days to tracheostomy	7 (5 to 11)	7 (5 to 10)	9 (6 to 12)	1.04 (0.99 to 1.10)	0.057
<b>Cause of extubation failure (more than one may apply) - n (%)</b>					
Coma	41 (6)	14 (4)	27 (10)	3.06 (1.57 to 5.97)	< 0.001
Agitation/delirium	11 (2)	8 (2)	3 (1)	0.54 (0.14 to 2.06)	0.371

**Table 4 Weaning variables of patients under invasive mechanical ventilation (Continued)**

Shock	20 (3)	3 (1)	17 (6)	8.86 (2.56 to 30.6)	< 0.001
Ventilator-associated pneumonia	13 (2)	6 (2)	7 (3)	1.73 (0.57 to 5.21)	0.330
Dyspnea	50 (8)	31 (9)	19 (7)	0.88 (0.49 to 1.61)	0.697
Cardiac dysfunction	7 (1)	2 (0.8)	5 (2)	3.71 (0.71 to 19.3)	0.118
Hypoxemia/Hypercapnia	36 (6)	16 (5)	20 (7)	1.91 (0.97 to 3.76)	0.063
Excessive respiratory secretions	24 (4)	14 (4)	10 (4)	1.05 (0.45 to 2.40)	0.912

have better organization and standards of care, it is possible that the actual mortality of mechanically ventilated patients may be even higher.

## Conclusions

Current mortality of mechanically ventilated patients in Brazil remains elevated. Implementation of judicious fluid therapy and a watchful use and monitoring of NIV patients are potential targets to improve outcomes in this setting.

## Key messages

- Contemporary information on mechanical ventilation use in emerging countries is limited. Moreover, most epidemiological studies on ventilatory support were carried out before significant developments, such as lung protective ventilation or widespread application of non-invasive ventilation.
- In mechanically ventilated patients in Brazil, factors such as age, comorbidities, ARDS, disease severity and variables related to ICU support, such as positive fluid balance and NIV failure, are independently related to hospital mortality.
- NIV failure occurred in 54% of the patients and was associated with the severity of organ dysfunctions, presence of ARDS and positive fluid balance.
- Current mortality of ventilated patients in Brazil is exceedingly high. Implementation of judicious fluid therapy and a watchful use and monitoring of NIV patients are potential targets to improve outcomes in this setting.

## Abbreviations

AIDS: Acquired immunodeficiency syndrome; ARDS: Acute respiratory distress syndrome; BRICNet: Brazilian Research in Intensive Care Network; CI: Confidence interval; COPD: Chronic Obstructive Pulmonary Disease; ERICC: Epidemiology of Respiratory Insufficiency in Critical Care; ICU: Intensive care unit; IRB: Institutional Review Board; LOS: Length-of-stay; LOWESS: Locally weighted scatterplot smoothing; MV: Mechanical ventilation; NIV: Non-invasive mechanical ventilation; OR: Odds ratio; PEEP: Positive end-expiratory pressure; RRT: Renal replacement therapy; SAPS 3: Simplified Acute Physiology Score 3; SOFA: Sequential Organ Failure Assessment

## Authors' contributions

All authors contributed significantly to this manuscript, including study conception (LCPA, MP, JIFS, GS, MS), data acquisition (all authors), data analysis and interpretation (LCPA, MP, MS, JIFS), drafting manuscript (LCPA, MS, JIFS), revising the manuscript for important intellectual content (all authors), and approval of the final copy (all authors).

## Competing interests

The authors declare they have no competing interests regarding the topic of this manuscript.

## Acknowledgements

This study was funded by the Research and Education Institute from Hospital Sírio-Libanês, São Paulo and the D'Or Institute for Research and Education, Rio de Janeiro. The study was supported by the Brazilian Research in Intensive Care Network (BRICNet). Dr. Azevedo had full access to all data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The authors would like to thank the following sites and investigators: **Bahia:** Hospital Espanhol - Salvador (Amadeu Martinez, Livia Leal, Antonio Jorge Pereira). **Distrito Federal:** Hospital Santa Luzia - Brasília (Marcelo de Oliveira Maia, José Aires Neto). **Espírito Santo:** Vitória Apart Hospital - Vitória (Claudio Piras), Centro Integrado de Atenção à Saude (CIAS) Unimed Vitória - Vitória (Eliana Bernadete Caser, Cora Lavigne Moreira), Hospital Meridional - Cariacica (Pablo Braga Gusman, Dyanne Moysés Dalcomune). **Maranhão:** UDI Hospital - São Luís (Alexandre Guilherme Ribeiro de Carvalho, Louise Aline Romão Gondim, Livia Mariane Castelo Branco Reis). **Minas Gerais:** Hospital Madre Tereza - Belo Horizonte (Daniel da Cunha Ribeiro, Leonardo de Assis Simões, Rafaela Siqueira Campos, José Carlos Fernandez Versiani dos Anjos), Hospital Mater Dei - Belo Horizonte (Frederico Bruzzi Carvalho). **Pará:** Hospital Regional Público do Araguaia - Redenção (Rossine Ambrosio Alves, Lilian Batista Nunes). **Paraná:** Hospital do Trabalhador - Curitiba (Álvaro Réa-Neto, Mirella Cristine de Oliveira), Hospital Vita Batel - Curitiba (Álvaro Réa-Neto, Mirella Cristine de Oliveira), Hospital Universitario Cajuru - Curitiba (Álvaro Réa-Neto, Luana Tannous), Instituto de Neurologia de Curitiba (INC) - Curitiba (Álvaro Réa-Neto, Brenno Cardoso Gomes). **Rio de Janeiro:** Instituto Nacional de Câncer - Hospital do Câncer I - Rio de Janeiro (Vicente Cés de Souza Dantas), Hospital de Clínicas de Niterói - Niterói (Fernando Borges Rodriguez, Priscila Abelha); Hospital de Clínicas Mario Lioni - Duque de Caxias (Marcelo E. Lugarinho); Instituto de Pesquisa Clínica Evandro Chagas (IPEC) - Rio de Janeiro (Andre Japiassu), Hospital da Mulher Heloneida Studart - Rio de Janeiro (Hélder Konrad de Melo, Elton Afonso Lopes), Hospital Pasteur - Rio de Janeiro (Pedro Varaschin, Vicente Cés de Souza Dantas), Hospital São Lucas - Rio de Janeiro (Marcos Freitas Knibel, Micheli Ponte, Pedro Mendes de Azambuja Rodrigues), Hospital Pro-Cardiaco - Rio de Janeiro (Rubens Carmo Costa Filho, Felipe Saddy, Thêia Forny Wanderley Castellões, Suzana Alves Silva), Nortecor Hospital de Clínica - Rio de Janeiro (Luiz Antonio Gomes Osorio, Dora Mannarino), Hospital Copa D'Or - Rio de Janeiro (Rodolfo Espinoza, Cassia Righy, Marcio Soares, Jorge Salluh, Lilian Tanaka, Daniel Aragão, Maria Eduarda Tavares, Maura Gonçalves Pereira Kehdi). **Roraima:** Hospital Geral de Roraima - Boa Vista (Valéria Maria Campos Rezende, Roberto Carlos Cruz Carbonell). **Rio Grande do Sul:** Hospital Moinhos de Vento - Porto Alegre (Cassiano Teixeira, Roselaine Pinheiro de Oliveira, Juçara Gasparetto Maccari, Priscylla Souza Castro), Santa Casa de Misericórdia de Porto Alegre - Pavilhão Pereira Filho (Paula Berto, Patricia Schwarz); Santa Casa de Misericórdia de Porto Alegre - Hospital Santa Rita (André Peretti Torelly, Thiago Lisboa, Paula Berto, Edison Moraes). **Santa Catarina:** Hospital São José - Criciúma (Felipe Dal-Pizzol, Cristiane Tomasi Damiani, Cristiane Ritter). **São Paulo:** Hospital A. C. Camargo - São Paulo (Juliana Carvalho Ferreira, Ramon Teixeira Costa, Pedro Caruso); Fundação Pio XII - Hospital de Câncer de Barretos - Barretos (Cristina Prata Amendola, Amanda Maria R. R. de Oliveira, Ulysses V. A. Silva, Luciana Coelho Sanches, Rosana D. S. Almeida); Hospital Sírio Libanês - São Paulo (Luciano Cesar Azevedo, Marcelo Park, Guilherme Schettino), Hospital Israelita Albert Einstein - São Paulo (Murillo Santucci Assunção, Eliezer Silva), Hospital São Camilo Santana - São Paulo (Carlos Eduardo Barboza, Antonio Paulo Nassar Junior), Hospital São Camilo Pompéia - São Paulo (Antonio

Paulo Nassar Junior), Hospital das Clínicas da Faculdade de Medicina da USP - UTI Disciplina Emergências Clínicas - São Paulo (Luciano Cesar Azevedo, Marcelo Park), Hospital das Clínicas da Faculdade de Medicina da USP - UTI Disciplina Emergências Cirúrgicas - São Paulo (Paulo Fernando Guimarães Morando Marzocchi Tierno, Luis Marcelo Malbouisson, Lucas Oliveira), Hospital das Clínicas da Faculdade de Medicina da USP - UTI Disciplina Anestesiologia - São Paulo (Davi Cristovao), Hospital Ipiranga - Rede Amil - São Paulo (Manoel Leitão Neto, Énio Rego, Fernanda Eugênia Fernandes), Hospital Do Coração - São Paulo (Marcelo Luz Pereira Romano, Alexandre Biasi Cavalcanti, Dalton de Souza Barros, Érica Aranha Suzumura, Karla Loureiro Meira, Gustavo Affonso de Oliveira), Hospital Estadual de Américo Brasiliense - Américo Brasiliense (Paula Menezes Luciano, Evelin Drociunas Pacheco), Hospital São Paulo da Universidade Federal de São Paulo - São Paulo (Bruno Franco Mazza, Flavia Ribeiro Machado, Elaine Ferreira), Hospital Universitário da Universidade de São Paulo - São Paulo (Ronaldo Batista dos Santos, Alexandra Siqueira Colombo, Antonio Carlos Nogueira, Juliana Baroni Fernandes, Raquel Siqueira Nóbrega, Barbara do C.S. Martins, Francisco Soriano), Hospital São Luiz Jardim Anália Franco - São Paulo (Rafaela Deczka Morsch, Andre Luiz Baptiston Nunes), Instituto do Câncer do Estado de São Paulo (ICESP) - São Paulo (Juliano Pinheiro de Almeida, Ludhmila Hajjar, Sílvia Moulin), Hospital e Maternidade São Luiz - Unidade Itaim - São Paulo (Fábio Poianas Giannini, Andre Luiz Baptiston Nunes).

#### Author details

<sup>1</sup>Research and Education Institute, Hospital Sírio-Libanês, Rua Cel. Nicolau dos Santos 69 - São Paulo, 01308-060, Brazil. <sup>2</sup>ICU, Emergency Medicine Department, Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, Av. Eneas Carvalho Aguiar 255, São Paulo, 05403-000, Brazil. <sup>3</sup>D'Or Institute for Research and Education, Rua Dinis Cordeiro 30, Rio de Janeiro, 22281-100, Brazil. <sup>4</sup>CEPETI - Centro de Estudos e Pesquisas em Terapia Intensiva, Rua Monte Castelo 366, Curitiba, 82530-200, Brazil. <sup>5</sup>ICU, Instituto Nacional de Câncer - Hospital do Câncer I, Praça Cruz Vermelha 23, Rio de Janeiro, 20230-130, Brazil. <sup>6</sup>ICU, Hospital Pasteur, Av. Amaro Cavalcanti 495, Rio de Janeiro, 20735-040, Brazil. <sup>7</sup>ICU, Surgical Emergency Department, Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, Av. Eneas Carvalho Aguiar 255, São Paulo, 05403-000, Brazil. <sup>8</sup>ICU, Hospital São José, Rua Cel. Pedro Benedet 630, Criciúma, 88801-250, Brazil. <sup>9</sup>ICU, Fundação Pio XII, Hospital de Câncer de Barretos, Rua Antenor Duarte Vilela 1331, Barretos, 14780-000, Brazil. <sup>10</sup>ICU, Hospital São Lucas, Travessa Frederico Pamplona 32, Rio de Janeiro, 22061-080, Brazil. <sup>11</sup>ICU, Hospital São Camilo Pompéia, Av. Pompeia 1178, São Paulo, 05024-000, Brazil. <sup>12</sup>ICU, Hospital Regional Público do Araguaia - Av. Brasil, s/n Qd. 30, Redenção, 68552-735, Brazil. <sup>13</sup>ICU, Hospital A. C. Camargo, R. Professor Antônio Prudente 109, São Paulo, 01509-010, Brazil. <sup>14</sup>ICU, Hospital Moinhos de Vento, Rua Ramiro Barcelos 910, Porto Alegre, 90035-001, Brazil. <sup>15</sup>ICU, Hospital Geral de Roraima, Av. Brg. Eduardo Gomes 3388, Boa Vista, 69305-284, Brazil. <sup>16</sup>ICU, Hospital Espanhol, Av. Sete de Setembro 4161, Salvador, 40140-110, Brazil. <sup>17</sup>ICU, Hospital Estadual Américo Brasiliense, Alameda Dr. Aldo Lupo 1260, Américo Brasiliense, 14802-520, Brazil.

Received: 7 January 2013 Revised: 12 March 2013

Accepted: 26 March 2013 Published: 4 April 2013

#### References

1. Estenssoro E: **The FINNALI study on acute respiratory failure: not the final cut.** *Intensive Care Med* 2009, **35**:1328-1330.
2. Wunsch H, Linde-Zwirble WT, Angus DC, Hartman ME, Milbrandt EB, Kahn JM: **The epidemiology of mechanical ventilation use in the United States.** *Crit Care Med* 2010, **38**:1947-1953.
3. **Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The Acute Respiratory Distress Syndrome Network.** *N Engl J Med* 2000, **342**:1301-1308.
4. Villar J, Blanco J, Añón JM, Santos-Bouza A, Blanch L, Ambrós A, Gandía F, Carriedo D, Mosteiro F, Basaldúa S, Fernández RL, Kacmarek RM, ALIEN Network: **The ALIEN study: incidence and outcome of acute respiratory distress syndrome in the era of lung protective ventilation.** *Intensive Care Med* 2011, **37**:1932-1941.
5. Herridge MS, Tansey CM, Matté A, Tomlinson G, Diaz-Granados N, Cooper A, Guest CB, Mazer CD, Mehta S, Stewart TE, Kudlow P, Cook D, Slutsky AS, Cheung AM, Canadian Critical Care Trials Group: **Functional**

6. **disability 5 years after acute respiratory distress syndrome.** *N Engl J Med* 2011, **364**:1293-1304.
7. Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, Kairalla RA, Deheinzelin D, Munoz C, Oliveira R, Takagaki TY, Carvalho CR: **Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome.** *N Engl J Med* 1998, **338**:347-354.
8. Demoule A, Girou E, Richard JC, Taille S, Brochard L: **Increased use of noninvasive ventilation in French intensive care units.** *Intensive Care Med* 2006, **32**:1747-1755.
9. Demoule A, Girou E, Richard JC, Taille S, Brochard L: **Benefits and risks of success or failure of noninvasive ventilation.** *Intensive Care Med* 2006, **32**:1756-1765.
10. Linko R, Okkonen M, Pettila V, Perttinen J, Parviainen I, Ruokonen E, Tenhunen J, Ala-Kokko T, Varpula T: **Acute respiratory failure in intensive care units. FINNALI: a prospective cohort study.** *Intensive Care Med* 2009, **35**:1352-1361.
11. Tomicic V, Espinoza M, Andresen M, Molina J, Calvo M, Ugarte H, Godoy J, Gálvez S, Maurelia JC, Delgado I, Esteban A, Grupo Chileno para el Estudio de la Ventilación Mecánica: **[Characteristics and factors associated with mortality in patients receiving mechanical ventilation: first Chilean multicenter study].** *Rev Med Chil* 2008, **136**:959-967.
12. Franca SA, Toufen C Jr, Hovnanian AL, Albuquerque AL, Borges ER, Pizzo VR, Carvalho CR: **The epidemiology of acute respiratory failure in hospitalized patients: a Brazilian prospective cohort study.** *J Crit Care* 2011, **26**:330, e1-8.
13. Esteban A, Anzueto A, Alía I, Gordo F, Apezteguía C, Pálizas F, Cide D, Goldwaser R, Soto L, Bugedo G, Rodrigo C, Pimentel J, Raimondi G, Tobin MJ: **How is mechanical ventilation employed in the intensive care unit? An international utilization review.** *Am J Respir Crit Care Med* 2000, **161**:1450-1458.
14. Charlson ME, Pompei P, Ales KL, MacKenzie CR: **A new method of classifying prognostic comorbidity in longitudinal studies: development and validation.** *J Chronic Dis* 1987, **40**:373-383.
15. Moreno RP, Metnitz PG, Almeida E, Jordan B, Bauer P, Campos RA, Iapichino G, Edbrooke D, Capuzzo M, Le Gall JR: **SAPS 3-From evaluation of the patient to evaluation of the intensive care unit. Part 2: Development of a prognostic model for hospital mortality at ICU admission.** *Intensive Care Med* 2005, **31**:1345-1355.
16. Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, Reinhart CK, Suter PM, Thijs LG: **The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine.** *Intensive Care Med* 1996, **22**:707-710.
17. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, Cohen J, Opal SM, Vincent JL, Ramsay G: **2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference.** *Crit Care Med* 2003, **31**:1250-1256.
18. Rosolem MM, Rabello LS, Lisboa T, Caruso P, Costa RT, Leal JV, Salluh JI, Soares M: **Critically ill patients with cancer and sepsis: clinical course and prognostic factors.** *J Crit Care* 2012, **27**:301-307.
19. Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, Camporota L, Slutsky AS: **Acute respiratory distress syndrome: the Berlin Definition.** *JAMA* 2012, **307**:2526-2533.
20. Hosmer DW, Lemeshow S: *Applied logistic regression. (Wiley Series in Probability and Statistics).* 2 edition. New York, NY: Wiley-Interscience; 2000.
21. Bernard GR, Artigas A, Brigham KL, Carlet J, Falke K, Hudson L, Lamy M, Legall JR, Morris A, Spragg R: **The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination.** *Am J Respir Crit Care Med* 1994, **149**:818-824.
22. Vincent JL, Akca S, De Mendonca A, Haji-Michael P, Sprung C, Moreno R, Antonelli M, Suter PM, SOFA Working Group: **The epidemiology of acute respiratory failure in critically ill patients(\*).** *Chest* 2002, **121**:1602-1609.
23. Behrendt CE: **Acute respiratory failure in the United States: incidence and 31-day survival.** *Chest* 2000, **118**:1100-1105.
24. Schettino G, Altobelli N, Kacmarek RM: **Noninvasive positive-pressure ventilation in acute respiratory failure outside clinical trials: experience at the Massachusetts General Hospital.** *Crit Care Med* 2008, **36**:441-447.
25. Esteban A, Ferguson ND, Meade MO, Frutos-Vivar F, Apezteguia C, Brochard L, Raymonds K, Nin N, Hurtado J, Tomicic V, González M, Elizalde J, Nightingale P, Abroug F, Pelosi P, Arabi Y, Moreno R, Jibaja M, D'Empaire G, Sandi F, Matamis D, Montañez AM, Anzueto A, VENTILA

- Group: Evolution of mechanical ventilation in response to clinical research. *Am J Respir Crit Care Med* 2008, **177**:170-177.
25. Metnitz PG, Metnitz B, Moreno RP, Bauer P, Del Sorbo L, Hoermann C, de Carvalho SA, Ranieri VM: **Epidemiology of mechanical ventilation: analysis of the SAPS 3 database.** *Intensive Care Med* 2009, **35**:816-825.
  26. Phua J, Badia JR, Adhikari NK, Friedrich JO, Fowler RA, Singh JM, Scales DC, Stather DR, Li A, Jones A, Gattas DJ, Hallett D, Tomlinson G, Stewart TE, Ferguson ND: **Has mortality from acute respiratory distress syndrome decreased over time? A systematic review.** *Am J Respir Crit Care Med* 2009, **179**:220-227.
  27. Estenssoro E, Dubin A, Laffaire E, Canales H, Sáenz G, Moseinco M, Pozo M, Gómez A, Baredes N, Jannello G, Osatnik J: **Incidence, clinical course, and outcome in 217 patients with acute respiratory distress syndrome.** *Crit Care Med* 2002, **30**:2450-2456.
  28. Cardoso LT, Grion CM, Matsuo T, Anami EH, Kauss IA, Seko L, Bonametti AM: **Impact of delayed admission to intensive care units on mortality of critically ill patients: a cohort study.** *Crit Care* 2011, **15**:R28.
  29. Bozza FA, Salluh JI: **An urban perspective on sepsis in developing countries.** *Lancet Infect Dis* 2010, **10**:290-291.
  30. Young MP, Manning HL, Wilson DL, Mette SA, Riker RR, Leiter JC, Liu SK, Bates JT, Parsons PE: **Ventilation of patients with acute lung injury and acute respiratory distress syndrome: has new evidence changed clinical practice?** *Crit Care Med* 2004, **32**:1260-1265.
  31. Serpa Neto A, Cardoso SO, Manetta JA, Pereira VG, Espósito DC, Pasqualucci Mde O, Damasceno MC, Schultz MJ: **Association between use of lung-protective ventilation with lower tidal volumes and clinical outcomes among patients without acute respiratory distress syndrome: a meta-analysis.** *JAMA* 2012, **308**:1651-1659.
  32. Esteban A, Anzueto A, Frutos F, Alía I, Brochard L, Stewart TE, Benito S, Epstein SK, Apezteguía C, Nightingale P, Arroliga AC, Tobin MJ, Mechanical Ventilation International Study Group: **Characteristics and outcomes in adult patients receiving mechanical ventilation: a 28-day international study.** *JAMA* 2002, **287**:345-355.
  33. Frutos-Vivar F, Esteban A, Apezteguía C, González M, Arabi Y, Restrepo MI, Gordo F, Santos C, Alhashemi JA, Pérez F, Peñuelas O, Anzueto A: **Outcome of reintubated patients after scheduled extubation.** *J Crit Care* 2011, **26**:502-509.
  34. Zagli G, Linden M, Spina R, Bonizzoli M, Cianchi G, Anichini V, Matano S, Benemei S, Nicoletti P, Peris A: **Early tracheostomy in intensive care unit: a retrospective study of 506 cases of video-guided Ciaglia Blue Rhino tracheostomies.** *J Trauma* 2010, **68**:367-372.
  35. Nathens AB, Rivara FP, Mack CD, Rubenfeld GD, Wang J, Jurkovich GJ, Maier RV: **Variations in rates of tracheostomy in the critically ill trauma patient.** *Crit Care Med* 2006, **34**:2919-2924.
  36. Flaatten H, Gjerde S, Heimdal JH, Aardal S: **The effect of tracheostomy on outcome in intensive care unit patients.** *Acta Anaesthesiol Scand* 2006, **50**:92-98.
  37. Antonelli M, Conti G, Moro ML, Esquinas A, Gonzalez-Diaz G, Confalonieri M, Pelaia P, Principi T, Gregoretti C, Beltrame F, Pennisi MA, Arcangeli A, Proietti R, Passariello M, Meduri GU: **Predictors of failure of noninvasive positive pressure ventilation in patients with acute hypoxemic respiratory failure: a multi-center study.** *Intensive Care Med* 2001, **27**:1718-1728.
  38. Carrillo A, Gonzalez-Diaz G, Ferrer M, Martinez-Quintana ME, Lopez-Martinez A, Llamas N, Alcazar M, Torres A: **Non-invasive ventilation in community-acquired pneumonia and severe acute respiratory failure.** *Intensive Care Med* 2012, **38**:458-466.
  39. Chandra D, Stamm JA, Taylor B, Ramos RM, Satterwhite L, Krishnan JA, Mannino D, Sciruba FC, Holguin F: **Outcomes of noninvasive ventilation for acute exacerbations of chronic obstructive pulmonary disease in the United States, 1998-2008.** *Am J Respir Crit Care Med* 2012, **185**:152-159.
  40. Sakr Y, Vincent JL, Reinhart K, Groeneveld J, Michalopoulos A, Sprung CL, Artigas A, Ranieri VM: **High tidal volume and positive fluid balance are associated with worse outcome in acute lung injury.** *Chest* 2005, **128**:3098-3108.
  41. National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network, Wiedemann HP, Wheeler AP, Bernard GR, Thompson BT, Hayden D, deBoisblanc B, Connors AF Jr, Hite RD, Harabin AL: **Comparison of two fluid-management strategies in acute lung injury.** *N Engl J Med* 2006, **354**:2564-2575.
  42. Payen D, de Pont AC, Sakr Y, Spies C, Reinhart K, Vincent JL: **A positive fluid balance is associated with a worse outcome in patients with acute renal failure.** *Crit Care* 2008, **12**:R74.
  43. Alonso WJ, Viboud C, Simonsen L, Hirano EW, Daufenbach LZ, Miller MA: **Seasonality of influenza in Brazil: a traveling wave from the Amazon to the subtropics.** *Am J Epidemiol* 2007, **165**:1434-1442.
  44. Ministerio-da-Saude, Brasil: **Cadastro Nacional de Estabelecimentos de Saude.** 2012 [<http://cnes.datasus.gov.br>].

doi:10.1186/cc12594

**Cite this article as:** Azevedo et al.: Clinical outcomes of patients requiring ventilatory support in Brazilian intensive care units: a multicenter, prospective, cohort study. *Critical Care* 2013 **17**:R63.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

