

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/345851556>

Novel Stepwise Approach to Assess Representativeness of a Large Multicenter Observational Cohort of Tuberculosis Patients: The Example of RePORT Brazil

Article in *International Journal of Infectious Diseases* · November 2020

DOI: 10.1016/j.ijid.2020.11.140

CITATIONS

0

READS

52

22 authors, including:



Maria Arriaga

Fundação Oswaldo Cruz

55 PUBLICATIONS 86 CITATIONS

[SEE PROFILE](#)



Artur Trancoso Lopo de Queiroz

Fundação Oswaldo Cruz

160 PUBLICATIONS 468 CITATIONS

[SEE PROFILE](#)



Moreno S. Rodrigues

Fundação Oswaldo Cruz

51 PUBLICATIONS 538 CITATIONS

[SEE PROFILE](#)



Mariana Araújo Pereira

Fundação Oswaldo Cruz

9 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Viral Infections [View project](#)



Respiratory pathogens, immune response and pathology [View project](#)

Journal Pre-proof

Novel Stepwise Approach to Assess Representativeness of a Large Multicenter Observational Cohort of Tuberculosis Patients: The Example of RePORT Brazil

María B. Arriaga, Gustavo Amorim, Artur T.L. Queiroz, Moreno M.S. Rodrigues, Mariana Araújo-Pereira, Betania M.F. Nogueira, Alexandra Brito Souza, Michael S. Rocha, Aline Benjamin, Adriana S.R. Moreira, Jamile G. de Oliveira, Marina C. Figueiredo, Megan M. Turner, Kleydson Alves, Betina Durovni, José R. Lapa-e-Silva, Afrânio L. Kritski, Solange Cavalcante, Valeria C. Rolla, Marcelo Cordeiro-Santos, Timothy R. Sterling, Bruno B. Andrade, for the RePORT Brazil consortium



PII: S1201-9712(20)32452-8

DOI: <https://doi.org/10.1016/j.ijid.2020.11.140>

Reference: IJID 4850

To appear in: *International Journal of Infectious Diseases*

Received Date: 5 October 2020

Revised Date: 28 October 2020

Accepted Date: 8 November 2020

Please cite this article as: Arriaga MB, Amorim G, Queiroz ATL, Rodrigues MMS, Araújo-Pereira M, Nogueira BMF, Souza AB, Rocha MS, Benjamin A, Moreira ASR, de Oliveira JG, Figueiredo MC, Turner MM, Alves K, Durovni B, Lapa-e-Silva JR, Kritski AL, Cavalcante S, Rolla VC, Cordeiro-Santos M, Sterling TR, Andrade BB, Novel Stepwise Approach to Assess Representativeness of a Large Multicenter Observational Cohort of Tuberculosis Patients: The Example of RePORT Brazil, *International Journal of Infectious Diseases* (2020), doi: <https://doi.org/10.1016/j.ijid.2020.11.140>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Novel Stepwise Approach to Assess Representativeness of a Large Multicenter Observational Cohort of Tuberculosis Patients: The Example of RePORT Brazil

Short title: Assessing sample representativeness of TB cohorts

María B. Arriaga^{1,2,3}, Gustavo Amorim⁴, Artur T.L. Queiroz^{2,5}, Moreno M.S. Rodrigues⁶, Mariana Araújo-Pereira^{1,2,3}, Betania M. F. Nogueira^{2,3,7,17}, Alexandra Brito Souza^{8,9}, Michael S. Rocha^{2,7}, Aline Benjamin¹⁰, Adriana S. R. Moreira¹¹, Jamile G. de Oliveira¹², Marina C. Figueiredo⁴, Megan M. Turner⁴, Kleydson Alves¹³, Betina Durovni^{11,14}, José R. Lapa-e-Silva¹¹, Afrânio L. Kritski¹¹, Solange Cavalcante^{10,12}, Valeria C. Rolla¹⁰, Marcelo Cordeiro-Santos^{8,9,15}, Timothy R. Sterling^{4*}, Bruno B. Andrade^{1,2,3,4,8,16,17,18*} for the RePORT Brazil consortium

¹Laboratório de Inflamação e Biomarcadores, Instituto Gonçalo Moniz, Fundação Oswaldo Cruz, Salvador, Brazil

²Multinational Organization Network Sponsoring Translational and Epidemiological Research (MONSTER) Initiative, Salvador, Brazil

³Faculdade de Medicina, Universidade Federal da Bahia, Salvador, Brazil

⁴Division of Infectious Diseases, Department of Medicine, Vanderbilt University School of Medicine, Nashville, Tennessee, USA.

⁵Center of Data and Knowledge Integration for Health (CIDACS), Instituto Gonçalo Moniz, Fundação Oswaldo Cruz, Salvador, Brazil

⁶Laboratório de Análise e Visualização de Dados, Fundação Oswaldo Cruz, Porto Velho, Brazil

⁷Instituto Brasileiro para Investigação da Tuberculose, Fundação José Silveira, Salvador, Brazil

⁸Fundação Medicina Tropical Doutor Heitor Vieira Dourado, Manaus, Brazil

⁹Programa de Pós-Graduação em Medicina Tropical, Universidade do Estado do Amazonas, Manaus, Brazil

¹⁰Instituto Nacional de Infectologia Evandro Chagas, Fiocruz, Rio de Janeiro, Brazil

¹¹Programa Acadêmico de Tuberculose. Faculdade de Medicina, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

¹²Secretaria Municipal de Saúde do Rio de Janeiro (Clínica da Família Rinaldo Delamare), Rocinha, Rio de Janeiro, Brazil

¹³Ministério da Saúde, National Tuberculosis Control Program, Brasília, Brazil

¹⁴Secretaria Municipal de Saúde do Rio de Janeiro, Rio de Janeiro, Brazil

¹⁵Universidade Nilton Lins, Manaus, Brazil

¹⁶Curso de Medicina, Universidade Salvador (UNIFACS), Laureate University, Salvador, Brazil

¹⁷Curso de Medicina, Centro Universitário Faculdade de Tecnologia e Ciências (UniFTC), Salvador, Brazil

¹⁸Programa de Pós-graduação em Ciências da Saúde, Faculdade de Medicina da Bahia, Universidade Federal da Bahia, Salvador, Bahia, Brazil.

* T.R.S. and B.B.A. equally contributed to the work

Corresponding author: Bruno B. Andrade, MD, PhD, Laboratório de Inflamação e Biomarcadores, Instituto Gonçalo Moniz, Fundação Oswaldo Cruz, Rua Waldemar Falcão, 121, Candeal, Salvador, Bahia 40296-710, Brazil. E-mail: bruno.andrade@fiocruz.br

40-word summary: We compared a multicenter tuberculosis cohort enrolled from three regions of Brazil with the Brazilian National Tuberculosis Program registry, using innovative statistical approaches. The cohort was representative of TB patients overall, facilitating generalizability of study outcomes.

Highlights

- Representativeness of a large TB cohort was tested using the Brazilian National TB databank
- Novel analysis of second-generation p-values uncovered similarities between the cohorts
- There is a need to improve epidemiological and operational indicators in the Brazilian TB program

Journal Pre-proof

ABSTRACT

Background: A major goal of tuberculosis (TB) epidemiologic studies is to obtain results that can be generalized to the larger population with TB. The ability to extrapolate findings on determinants of TB treatment outcomes is also important.

Methods: We compared baseline clinical and demographic characteristics and determinants of anti-TB treatment outcomes between persons enrolled into the Regional Prospective Observational Research in Tuberculosis (RePORT)-Brazil cohort between June 2015 and June 2019, and the registry of TB cases reported to the Brazilian National TB Program (Information System for Notifiable Diseases [SINAN]) during the same time period. Multivariable regression models adjusted for study site were performed using *second-generation* p-values, a novel statistical approach. Associations with unfavorable treatment outcomes were tested for both RePORT-Brazil and SINAN cohorts.

Findings: 1,060 culture-confirmed TB patients were enrolled in RePORT-Brazil and 455,873 TB cases were reported to SINAN. *Second-generation* p-value analyses revealed that the cohorts were strikingly similar with regard to sex, age, use of antiretroviral therapy and positive initial smear sputum microscopy. However, diabetes, HIV infection and smoking were more frequently documented in RePORT-Brazil. Illicit drug use, presence of diabetes and history of prior TB were associated with unfavorable TB treatment outcomes; illicit drug use was associated with such outcomes in both cohorts.

Conclusions: There were important similarities in demographic characteristics and determinants of clinical outcomes between the RePORT-Brazil cohort and the Brazilian National registry of TB cases.

Keywords: tuberculosis; cohort study; sample representativeness; epidemiology; treatment outcome

INTRODUCTION

Tuberculosis (TB) is a major cause of death worldwide, especially in people living with HIV (PLWH) (World Health Organization, 2019). The World Health Organization (WHO) estimated that in 2018, ten million people developed TB, and of those, 1.5 million died (including 251,000 PLWH) (World Health Organization, 2019). The WHO, through the End TB Strategy, (World Health Organization, 2017) has proposed two essential pillars to improve TB control worldwide: (i) robust, coordinated data systems, prioritizing existing systems where possible, and (ii) support systems that promote research and innovation (World Health Organization, 2017). Routinely collected TB data allow for monitoring and calculating epidemiological and operational indicators, which are important for TB control (World Health Organization, 2018). However, to be useful and effective, frequent data updates and quality control are essential for accurate data (Agency for Healthcare Research and Quality (US), 2014).

TB research has a vital role in generating knowledge that guides decision-making strategies (World Health Organization, 2017). However, different components of study design (e.g., eligibility criteria, recruitment and enrollment, retention) can include confounders and biases, which may affect the representativeness of the study population and limit generalizability of the results (He et al., 2016). This can result in erroneous estimates, and consequently lead to adoption of inappropriate actions or strategies (Kukull and Ganguli, 2012). Thus, it is of critical importance to evaluate the representativeness of a study population, and whether results from the study population reflect what occurs in the larger population (Jaehn et al., 2020, Kukull and Ganguli, 2012).

In Brazil, the National TB Program (NTBP) implemented in 1993 collection of detailed demographic and clinical data on all TB patients. Importantly, notification of all TB cases and reporting such information is mandatory, and occurs through an electronic system referred as the Notifiable Disease Information System (SINAN) (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2020b). This system is critical for monitoring the epidemiology of TB in all Brazilian regions, guiding governmental investments

and policy changes to optimize patient care, minimizing the occurrence of drug-resistance and leading to better TB outcomes and control. SINAN is also a unique source of data for scientists interested in testing the generalizability of their research populations and results.

Although national TB datasets are useful, often the amount and detail of clinical and demographic data that can be collected is limited, and patient follow-up is passive rather than active. Prospective observational cohorts with active participant follow-up for TB treatment outcomes are valuable, but their cost usually limits cohort size. Regional Prospective Observational Research in Tuberculosis (RePORT) Brazil is a multicenter cohort study with enrollment sites in 3 major regions in Brazil (Hamilton et al., 2015, Regional Prospective Observational Research for Tuberculosis, 2020). However, it is unclear whether the RePORT-Brazil cohort reflects TB epidemiology and treatment outcomes nationally. We therefore evaluated the representativeness of RePORT-Brazil by comparing epidemiological variables and operational indicators with those in the SINAN dataset. In addition, clinical and epidemiologic determinants of unfavorable TB treatment outcomes were also compared between the two cohorts. We also assessed whether data and variables collected in RePORT-Brazil could improve the utility of SINAN.

MATERIALS AND METHODS

Overall Study Design

We compared baseline clinical and sociodemographic characteristics, and determinants of anti-TB treatment outcomes among persons with TB enrolled into RePORT-Brazil and those reported to SINAN, both between June 2015 and June 2019. The primary hypothesis was that the overall characteristics of TB patients enrolled into RePORT-Brazil and reported to SINAN were similar.

The Regional Prospective Observational Research in Tuberculosis (RePORT) - Brazil

Description of RePORT sites is presented in **Supplementary Methods** and in (Regional Prospective Observational Research for Tuberculosis, 2020). Between June 2015 and June 2019, 1060 persons ≥ 18 years old with new or recurrent pulmonary TB (with or without extrapulmonary disease) and culture-positive sputum (Lowenstein-Jensen medium or BD BACTEC MGIT) were enrolled. Epidemiologic information was collected during study visits at baseline, during, and at the end of treatment, and up to 24 months after enrollment. This included sex, age, self-reported race, weight, height, education level, consumption of alcohol, use of illicit drugs and smoking, presence of comorbidities and HIV status. Additionally, glycosylated hemoglobin testing (HbA1c), chest X-ray, drug susceptibility testing for anti-TB drugs and CD4 count (if HIV positive) were performed. In RePORT-Brazil the treatment outcome was recorded at the last study visit (24 months after the start of treatment). For this study, 434 participants had not yet completed their 24-month visit, so the analysis of treatment outcomes did not include these participants. All data were entered into a REDCap database (Harris et al., 2009) and stored at the RePORT-Brazil data coordination center at Vanderbilt University Medical Center. The database was continuously monitored for data quality and completeness and updated as appropriate.

Notifiable Diseases Information System (SINAN), Brazilian Ministry of Health

SINAN is a system for the notification and investigation of transmissible diseases that has been implemented, supported and maintained by the Brazilian Ministry of Health (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2007). Description of SINAN is presented in **Supplementary Methods**. Data from 455,873 TB patients were reported to SINAN between 2015 and 2019. TB cases were diagnosed by one or more of the following criteria: (a) clinical and epidemiologic factors (presumptive diagnosis), (b) bacteriology (sputum smear positive) or positive culture (solid or liquid), (c) GeneXpert MTB RIF, (d) chest radiography or (e) in the case of extrapulmonary TB by histopathology (the details regarding the diagnosis are described in the Manual of Recommendations for the Control of TB in Brazil) (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2013). After TB diagnosis, the information collected in the evaluation interview and the laboratory results were recorded on a standardized form that included the clinical form of TB, individual characteristics (sex, age, race, education, alcohol consumption, illicit drug use, smoking habits, associated conditions and diseases), presence of TB-HIV coinfection and test results, among others (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2013). Information regarding the anti-TB treatment outcome was also provided.

Outcome definition

For this study, a favorable treatment outcome was defined as cure or completed treatment and an unfavorable outcome was defined as treatment failure, lost to follow-up or death during treatment. The definitions for clinical and bacteriological cure, failure, lost to follow-up and death for both cohorts corresponded with those in the Manual of Recommendations for the Control of TB of Brazil (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2013). The definitions of treatment outcomes for RePORT-Brazil and SINAN are described in **Supplementary Table 1**.

Operational Indicators

We also evaluated operational indicators for TB, which monitor the performance of TB Control Programs (Arakawa et al., 2015). The list of indicators is shown in **Supplementary Figure 1**.

Data analysis

All analyses were pre-specified. Categorical variables were compared using a two-sided Pearson's chi-square test (with Yates correction) or the Fisher's two-tailed test in 2x3 or 2x2 tables, respectively. Quantitative variables were compared using the Mann Whitney U test. LASSO (Least Absolute Shrinkage and Selection Operator) (Trevor Hastie et al., 2009) regression analysis and mixed effects models (Gelman and Hill, 2006) were performed, in RePORT-Brazil and SINAN cohorts, to identify independent associations between clinical characteristics of TB patients and anti-TB treatment unfavorable outcomes. In LASSO regression, the optimal parameters were found by a cross-validation step, which was repeated 100 times to stabilize the results. The mixed effects model included the variable "Brazilian states" as a random effect (to avoid possible selection bias, because RePORT-Brazil participants were also reported to and part of the SINAN dataset). Results from both regressions approaches were presented in terms of point estimates and 95% confidence intervals (95%CI). P-values <0.05 were considered statistically significant. In addition to the p-value, the *second-generation* p-value ($p\delta$ -value) and the *delta-gap* (Δ) (when applicable) were calculated (Blume et al., 2018) as described in **Supplementary Methods** and **Supplementary Figure 2**.

RESULTS

Tuberculosis Epidemiology in Brazil and RePORT-Brazil sites

Brazil has 26 states and a federal district, all allocated to five macro-regions. RePORT-Brazil has sites located in the cities of Manaus, Salvador, Rio de Janeiro and Duque de Caxias (**Figure 1A**). Between 2015 and 2019, Sao Paulo was the state with the highest number of notified TB cases, followed by Rio de Janeiro and Rio Grande do Sul; the states of Bahia and Amazonas were in the top 10 (**Figure 1B**). The top 10 cities in number of TB cases documented during the same period included all of the cities that had RePORT sites, including Rio de Janeiro (1), Manaus (3), Salvador (4), and Duque de Caxias (10) (**Figure 1B**). When evaluating TB incidence per 100,000 persons in the Brazilian states where RePORT-Brazil has sites, different patterns were detected. Bahia had a trend in TB incidence that was similar to that observed in the whole country, whereas Amazonas had a substantially higher incidence over time, followed by Rio de Janeiro (**Figure 1C, left panel**). Similarly, Manaus city displayed the highest TB incidence among the cities containing RePORT-Brazil sites, followed by Rio de Janeiro, Duque de Caxias and Salvador. (**Figure 1C, right panel**). **Figure 1D** compares the number of cases notified to SINAN and the number of cases in RePORT Brazil in each year of the study period. The number of cases enrolled into RePORT-Brazil (1060 TB cases) represented 0.23% of the total registries in SINAN (455,873 TB cases). These analyses indicate that RePORT-Brazil sites are located in regions of Brazil that account for a significant TB burden of the country.

Characteristics of the TB cases in RePORT-Brazil and SINAN

Population characteristics of TB cases in both datasets were then compared (**Table 1**). In both datasets, the majority of patients were male, and the age distributions were similar (second-generation p-values not significant). The frequencies of self-reported race were somewhat similar between the two cohorts (second-generation p-value indeterminate: $p\delta$ -value =0.5). The RePORT cohort had a higher proportion of literate persons (95.3%, compared to 83% in SINAN; ($p<0.001$, $p\delta=0$; $\Delta=14.9$). Compared to the national dataset, the RePORT cohort also had a higher proportion of health care workers (4.3% vs. 1.3%), proportion with

no comorbidities (82.9% vs. 74.1%), prevalence of diabetes (24% vs. 8%) and HIV infection (21% vs. 13.5%), alcohol consumption (83.9% vs. 19.5%), use of illicit drugs (34.2% vs. 15.6%), smoking (52.4% vs. 24.3%), abnormal chest x-ray (97% vs. 93.3%), presence of extrapulmonary TB (11% vs. 3%) and drug resistant *Mycobacterium tuberculosis* (17.3% vs. 9.7%), all with $p < 0.001$ and $p\delta = 0$; **Table 1**).

In both cohorts, the most frequent treatment outcome was cure (69.5% in RePORT and 66.9% in SINAN). Treatment failure was more frequently documented in the RePORT cohort (4.2% vs. 1.9%), whereas patients who were transferred out to health care facilities of higher complexity were more commonly observed in the SINAN dataset (7.3% vs. 1.9%). When all categories of treatment outcomes were compared between the datasets, a statistically significant difference was observed between RePORT and SINAN ($p < 0.001$ and a $p\delta = 0$; **Table 1**).

Additional comparisons are shown in **Figure 2**. The analyses of the second-generation p-values detected in the comparisons of the age distribution between patients stratified by sex ($p\delta = 0.135$), diabetes comorbidity ($p\delta = 0$, with a delta-gap of 14.4) and HIV status ($p\delta = 0.245$), indicated that there were marginal differences (except for diabetes) in age distributions within each of these subpopulations between the datasets (**Figure 2A**). Of note, the majority of the TB patients were new cases (treatment naïve), with a similar profile between the datasets ($p\delta = 0.5$; **Table 1, Figure 2B**). The populations from the two distinct datasets exhibited similar proportions of positive acid-fast bacilli (AFB) in sputum smears ($p\delta = 0.5$; **Figure 2C**). In contrast, the proportions of TB cases with abnormal chest radiographs and positive sputum culture results were substantially higher in the RePORT cohort than SINAN (all $p\delta = 0$; **Figure 2C**); a positive sputum culture was required for enrollment into RePORT-Brazil. TB patients from RePORT-Brazil more frequently reported smoking and alcohol consumption, and more commonly had diabetes, than patients from SINAN ($p\delta = 0$; **Figure 2D**).

Characteristics associated with unfavorable tuberculosis treatment outcomes

In RePORT-Brazil, the patients who experienced an unfavorable treatment outcome were more frequently male (75.4%), *pardo* (57.6%), and had diabetes (31.1%) and/or HIV infection (44%). In addition, alcohol consumption (92.1%), the use of illicit drugs (47.6%) and smoking (63.4%) were also associated ($p < 0.001$ in all) with unfavorable anti-TB treatment outcomes (**Supplementary Table 3**). Similarly, in the SINAN cohort, male sex (72.3%), *pardo* race (53%), HIV infection (24.9%), alcohol consumption (27.6%), illicit drug use (23.3%) and smoking (30.2%) were all associated with unfavorable outcomes. In addition, increasing age, previous TB (30.5%) and positive sputum culture (69.9%) also characterized individuals who had unfavorable outcomes (**Supplementary Table 4**). Comparisons of clinical characteristics and treatment outcomes between Report-Brazil and SINAN patients in the cities of Duque de Caxias, Manaus, Salvador and Rio de Janeiro are shown in **Supplemental tables 5, 6, 7 and 8**.

To define the specific associations with unfavorable treatment outcomes in each cohort, LASSO regression (model 1) and mixed effects (model 2) were utilized. The LASSO regression model showed that prior TB, diabetes, illicit drug use and HIV coinfection were all independently associated with the occurrence of unfavorable treatment outcomes in the RePORT-Brazil cohort (**Figure 3, left panel**), whereas increased age, male sex, alcohol consumption, illicit drug use, prior TB and HIV coinfection were associated with an increased odds of such outcomes in patients from the SINAN cohort (**Figure 3, right panel**). In the mixed effects model, diabetes comorbidity (OR:1.81; 95%CI:1.11-2.94, $p\delta=0$; $\Delta=1.0$) and the use of illicit drugs (OR: 2.28; 95%CI:1.36-3.82; $p\delta=0$; $\Delta=2.2$) were associated with an unfavorable outcome in RePORT-Brazil (**Figure 3, left panel**). Likewise, in the SINAN cohort, the use of illicit drugs (OR:1.85; 95%CI:1.80-1.91; $p\delta=0$; $\Delta=5.1$) was also associated with an unfavorable result, along with prior TB (OR: 1.96; 95%CI:1.95-1.97; $p\delta=7.5$), HIV positive (OR: 2.81; 95%CI: 2.74-2.90; $p\delta=0$; $\Delta=9.6$) and alcohol consumption (OR: 1.42; 95%CI:1.38-2.53; $p\delta=0$; $\Delta=2.4$) (**Figure 3, right panel**).

Epidemiological and operational indicators of tuberculosis cases in RePORT-Brazil and SINAN

Finally, the epidemiological and operational indicators were compared between the cohorts (**Supplementary Figure 1A-L**). Among the 11 operational indicators, significant differences were identified between RePORT-Brazil and SINAN, during each year in the study period. The frequency of new cases of pulmonary TB diagnosed through Xpert -MTB/-RIF, smear, culture or drug susceptibility testing among the total pulmonary TB cases was lower in each year in the SINAN cohort, likely due to RePORT inclusion criteria ($p < 0.01$, in each year) (**Supplementary Figure 1B**). In turn, this difference was also reflected by higher proportions of each type of TB screening test compared (**Supplementary Figure 1, 1F, 1H**). Coverage of HIV testing was also different between the datasets, with 70% in SINAN and 95% in RePORT ($p < 0.001$), also likely due to RePORT inclusion criteria. The ART initiation and cases receiving DOT was also significantly higher in RePORT-Brazil.

DISCUSSION

In this study, demographic characteristics such as age, sex, and race were similar in both the RePORT-Brazil and SINAN cohorts of TB patients, supporting our hypothesis that the RePORT-Brazil cohort is representative of individuals with pulmonary TB in Brazil (Fernandes et al., 2018, Zhang et al., 2011). However, certain characteristics varied in the RePORT-Brazil cohort compared to the general TB population. Literacy rate was higher among patients in RePORT, possibly because study sites are located in large cities, where access to school and overall literacy are higher than in rural areas (Ministério da Educação, 2008). Alcohol consumption, smoking and use of illicit drugs were significantly more commonly reported in RePORT patients. The RePORT study may have a higher detection rate of substance use as it was included in structured questionnaires and validated tools for screening. Conversely, SINAN notification templates include simple yes/no questions to evaluate these characteristics, which may be more prone to reporting or detection bias (Rocha et al., 2020). Moreover, SINAN templates are completed by health care workers who collect this information from the patient or medical charts, which often lack the detailed information needed (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2007). These factors may have led to reduced substance use identification in the national database, as noted previously (Rocha et al., 2020, Silva et al., 2010).

The proportions of patients reporting co-morbidities such as cancer, chronic obstructive pulmonary disease, kidney disease, and hypertension were similar in both cohorts. The proportion reporting “other” co-morbidities was higher in SINAN, and the proportions reporting diabetes or HIV were higher in RePORT-Brazil. This may be due to differences in ascertainment of these co-morbid conditions in the two cohorts (Althubaiti, 2016). Abnormal chest x-rays were reported more often in RePORT cases, possibly reflecting care provided by trained TB specialists, which is not as common in most health care facilities in Brazil (Shazzadur Rahman et al., 2019). The increased frequency of extrapulmonary manifestations of TB may also reflect treatment by experienced TB specialists and greater availability of diagnostic tests at reference centers from RePORT.

Interestingly, our study identified differences in relevant clinical characteristics such as an increased prevalence of DM, HIV, and multidrug resistance in RePORT compared to SINAN. These disparities may not necessarily reflect differences in the prevalence of these conditions in the study populations, but rather more frequent diagnosis and notification in RePORT because of uniform HbA1c and HIV testing. These comorbidities are known to be more frequent in patients with TB (Saraceni et al., 2018, Tankeu et al., 2017).

Similarly, all participants in RePORT-Brazil were required to have positive *M. tuberculosis* sputum cultures and drug susceptibility testing, in contrast to SINAN, resulting in a selection bias. When compared to the national data, we found that in 2019 only 76.1% of patients with TB were tested for HIV, despite the fact that the national guidelines recommend HIV testing in all patients with a TB diagnosis and that low-cost HIV testing is available nationally. Testing for DM is not routinely recommended by national guidelines; patients may have blood glucose testing if they exhibit risk factors or classical symptoms of DM (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2013). In SINAN, the prevalence of DM is likely only self-reported DM and may therefore underrepresent the actual prevalence.

Regarding the detection of drug resistant TB (DR-TB), the national statistics also indicate possible underdiagnosis. In 2019 nationally, 37.7% of patients with newly diagnosed TB had Xpert-MTB/RIF performed, and only 24% had sputum culture performed, suggesting that, except for rifampicin, drug sensitivity was unknown for many cases. Among cases of re-treatment, 30.4% had sputum culture performed and only 50% of those completed the recommended diagnostic workup including drug sensitivity tests. This reflects the ongoing limited access to Xpert-MTB-RIF technology nationally (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2020a, Silva et al., 2018), and lack of routine solicitation from physicians or capacity to perform sputum cultures and drug sensitivity tests in cases of presumed TB. The comparison of these two datasets indicates that the broad testing strategy adopted in the RePORT-Brazil protocol more accurately detected the presence of important comorbidities, exposures and drug resistance in TB cases.

RePORT-Brazil and SINAN had similar proportions of cases with cure, death and loss to follow-up. Reporting on these outcomes is less difficult than other TB outcomes possibly due to the simplicity of their definitions. Nevertheless, RePORT had a higher proportion of treatment failures, and lower frequency of individuals who were transferred out, than SINAN. These discrepancies may be because patients enrolled in RePORT-Brazil were actively followed up for 24 months from treatment initiation. In such scenarios, the treatment outcome is likely obtained in a more reliable fashion, especially in cases of treatment failure, loss to follow-up and transferred out. In SINAN, information on outcome is usually recorded at the end of treatment. In public health care centers, it is not recommended that patients return after finishing anti-TB treatment. Indeed, other studies (Baldez do Canto and Borges Nedel, 2020, Rocha et al., 2020, Silva et al., 2010) have described limitations related to underreporting of treatment outcome results and lack of follow-up for patients who were transferred out or discontinued treatment.

This study had some limitations. RePORT-Brazil had more active follow-up than SINAN, and the ability to follow-up prospectively in the latter cohort was mainly because of the large number of patients. In addition, routine follow-up for clinical care usually does not extend beyond of the end of TB treatment. Another limitation was that follow-up for study endpoints in RePORT-Brazil is ongoing. However, for this analysis we truncated follow-up so that it was similar to SINAN. Nonetheless, the present study adds to current knowledge in the field by demonstrating the close similarities of the RePORT-Brazil and SINAN cohorts regarding patient characteristics and treatment outcomes. This is beneficial for both cohorts. For the smaller RePORT-Brazil cohort, this supports the generalizability of findings from current and future translational studies (Jaehn et al., 2020). For the larger SINAN cohort, it means that smaller, less-expensive population-based studies in RePORT-Brazil may provide important data that can inform TB treatment strategies throughout all of Brazil.

In this study, we found similarities between both databases, though significant differences were identified regarding TB diagnosis, due primarily to RePORT-Brazil inclusion criteria. This study highlights critical gaps in comprehensive TB care, and identifies data that could be collected in SINAN that would facilitate

improvements in the care and outcomes of TB patients in Brazil (Ministério da Saúde do Brasil and Secretaria de Vigilância em Saúde, 2013).

NOTES

Acknowledgments: The authors thank the study participants. We also thank the teams of clinical and laboratory platforms of RePORT Brazil. A special thanks to Elze Leite (FIOCRUZ, Salvador, Brazil), Eduardo Gama (FIOCRUZ, Rio de Janeiro, Brazil), Elcimar Junior (FMT-HVD, Manaus, Brazil) and Hilary Vansell (VUMC, Nashville, USA), for administrative and logistical support.

Ethical approval: All clinical investigations were conducted according to the principles of the Declaration of Helsinki. The RePORT-Brazil protocol, informed consent, and study documents were approved by the institutional review boards at each study site and at Vanderbilt University Medical Center. Participation in RePORT-Brazil was voluntary, and written informed consent was obtained from all such participants. For the data extracted from SINAN, the anonymity of study subjects was preserved, and all data were de-identified.

Funding: The study was supported by the Intramural Research Program of the Fundação Oswaldo Cruz (B.B.A.), Intramural Research Program of the Fundação José Silveira (B.B.A., M.S.R., B.M.F.N.), Departamento de Ciência e Tecnologia (DECIT) - Secretaria de Ciência e Tecnologia (SCTIE) – Ministério da Saúde (MS), Brazil [25029.000507/2013-07 to V.C.R.] and the National Institutes of Allergy and Infectious Diseases [U01-AI069923 to T.R.S. and U01-AI115940 to B.B.A.]. M.B.A. received a fellowship from the Fundação de Amparo à Pesquisa da Bahia (FAPESB). M.A.-P. received a fellowship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Finance code: 001). B.B.A., J.R.L.S. and A.K. are senior investigators of the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil.

Disclaimer: The funder of the study had no role in the study design, data collection, or data analysis; however, a representative of the Brazilian Ministry of Health (K.A.) was involved in data interpretation and writing the report.

Conflicts of interest: All authors: No reported conflicts of interest.

Author contributions: Conceptualization, T.R.S., M.C.F., M.C.S., V.C.R., and B.B.A.; Data curation, M.B.A., G.A., A.T.L.Q., M.M.S.R., M.M.T., and B.B.A.; Investigation, M.B.A., G.A., A.T.L.Q., B.N., A.B.S., M.S.R., A.B., A.S.R.M., J.G.d.O., M.C.F., K.A., B.D., J.R.L.S., A.L.K., S.C., V.C.R., T.R.S., M.C.S., and B.B.A.; Formal analysis, M.B.A., G.A., A.T.L.Q., M.M.S.R., M.A.P., M.C.F., and B.B.A.; Funding acquisition, B.D., J.R.L.S., A.L.K., S.C., V.C.R., T.R.S., M.C.S., M.C.F., and B.B.A.; Methodology, M.B.A., G.A., A.T.L.Q., and B.B.A.; Project administration, M.C.F., T.R.S., and B.B.A.; Resources, M.B.A., G.A., T.R.S., and B.B.A.; Software, M.B.A., G.A., A.T.L.Q., M.M.S.R., M.A.P., M.C.F., T.R.S., and B.B.A.; Supervision, T.R.S., and B.B.A.; Writing—original draft, M.B.A., B.N., M.A.P., and B.B.A.; Writing—review and editing, all authors. All authors have read and agreed to the submitted version of the manuscript.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

- Agency for Healthcare Research and Quality (US). Registries for Evaluating Patient Outcomes. 2014.
- Althubaiti A. Information bias in health research: definition, pitfalls, and adjustment methods. *J Multidiscip Healthc* 2016;9:211-7.
- Arakawa T, Magnabosco GT, Lopes LM, Arnaez MA, Gavin MA, Gallardo Mdel P, et al. Evaluation of the performance of Tuberculosis Control Programs in Brazil and Spain: an integrative review of the literature. *Cien Saude Colet* 2015;20(12):3877-89.
- Baldez do Canto V, Borges Nedel F. Completude dos registros de tuberculose no Sistema de Informação de Agravos de Notificação (Sinan) em Santa Catarina, Brasil, 2007-2016. *Epidemiol Serv Saúde* 2020;29(3).
- Blume JD, D'Agostino McGowan L, Dupont WD, Greevy RA, Jr. Second-generation p-values: Improved rigor, reproducibility, & transparency in statistical analyses. *PLoS One* 2018;13(3):e0188299.
- Fernandes P, Ma Y, Gaeddert M, Tsacogianis T, Marques-Rodrigues P, Fregona G, et al. Sex and age differences in *Mycobacterium tuberculosis* infection in Brazil. *Epidemiol Infect* 2018;146(12):1503-10.
- Gelman A, Hill J. *Data Analysis Using Regression and Multilevel/Hierarchical Models* 2006.
- Hamilton CD, Swaminathan S, Christopher DJ, Ellner J, Gupta A, Sterling TR, et al. RePORT International: Advancing Tuberculosis Biomarker Research Through Global Collaboration. *Clin Infect Dis* 2015;61Suppl 3:S155-9.
- Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42(2):377-81.
- He Z, Ryan P, Hoxha J, Wang S, Carini S, Sim I, et al. Multivariate analysis of the population representativeness of related clinical studies. *J Biomed Inform* 2016;60:66-76.
- Jaehn P, Rehling J, Klawunn R, Merz S, Holmberg C, group AGs. Practice of reporting social characteristics when describing representativeness of epidemiological cohort studies - A rationale for an intersectional perspective. *SSM Popul Health* 2020;11:100617.
- Kukull WA, Ganguli M. Generalizability: the trees, the forest, and the low-hanging fruit. *Neurology* 2012;78(23):1886-91.
- Ministério da Educação. National Report from Brazil. Brazil2008.
- Ministério da Saúde do Brasil, Secretaria de Vigilância em Saúde. Sistema de Informação de Agravos de Notificação. Brazil2007.
- Ministério da Saúde do Brasil, Secretaria de Vigilância em Saúde. Manual de Recomendações para o Controle da Tuberculose no Brasil. 2013. p. 288.
- Ministério da Saúde do Brasil, Secretaria de Vigilância em Saúde. Boletim Epidemiológico Tuberculose. 2020a.

Ministério da Saúde do Brasil, Secretaria de Vigilância em Saúde. Sistema de Informação de Agravos de Notificação; 2020b. Available from: <http://portalsinan.saude.gov.br/>. [Accessed 08-14-2020].

Ministério da Saúde do Brasil SdVeS. Plano Nacional pelo Fim da Tuberculose como Problema de Saúde Pública. 2017. p. 54.

Regional Prospective Observational Research for Tuberculosis. RePORT-Brazil; 2020. Available from: www.reportbrazil.org. [Accessed 08-08-2020].

Rocha MS, Bartholomay P, Cavalcante MV, Medeiros FC, Codenotti SB, Pelissari DM, et al. Notifiable Diseases Information System (SINAN): main features of tuberculosis notification and data analysis. *Epidemiol Serv Saude* 2020;29(1):e2019017.

Saraceni V, Benzaken AS, Pereira GFM, Andrade KB, Oliveira PB, Arakaki-Sanchez D, et al. Tuberculosis burden on AIDS in Brazil: A study using linked databases. *PLoS One* 2018;13(11):e0207859.

Shazzadur Rahman AAM, Langley I, Galliez R, Kritski A, Tomeny E, Squire SB. Modelling the impact of chest X-ray and alternative triage approaches prior to seeking a tuberculosis diagnosis. *BMC Infect Dis* 2019;19(1):93.

Silva DR, Sotgiu G, D'Ambrosio L, Pereira GR, Barbosa MS, Dias NJD, et al. Diagnostic performances of the Xpert MTB/RIF in Brazil. *Respir Med* 2018;134:12-5.

Silva GDMd, Bartholomay P, Cruz OG, Garcia LP. Assessment of the tuberculosis surveillance system in Rio de Janeiro/RJ, Brazil, 2001 to 2006. *Cad saúde colet* 2010;18(3).

Tankeu AT, Bigna JJ, Nansseu JR, Endomba FTA, Wafeu GS, Kaze AD, et al. Global prevalence of diabetes mellitus in patients with tuberculosis: a systematic review and meta-analysis protocol. *BMJ Open* 2017;7(6):e015170.

Trevor Hastie, Robert Tibshirani, Friedman J. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*, Second Edition Springer Series in Statistics. 2, editor2009.

World Health Organization. Framework for implementing the “End TB Strategy” in the African Region 2016-2020. World Health Organization; 2017.

World Health Organization. Guidance for tuberculosis programme managers, analysis and use of health facility data. 2018. p. 1-44.

World Health Organization. Global Tuberculosis Report 2019. 2019. p. 283.

Zhang X, Andersen AB, Lillebaek T, Kamper-Jorgensen Z, Thomsen VO, Ladefoged K, et al. Effect of sex, age, and race on the clinical presentation of tuberculosis: a 15-year population-based study. *Am J Trop Med Hyg* 2011;85(2):285-90.

Table 1. Characteristics of the study participants

Characteristics	RePORT-Brazil (n=1,060)	SINAN (n=455,873)	p-value	p δ -value
Sex – no. (%)			0.027	0.744
Male	702 (66.3)	316693 (69.5)		
Female	358 (33.7)	139149 (30.5)		
Age – median (IQR)	36 (25-49)	37 (26-52)	0.001	1.0
Race/Ethnicity – no. (%)			0.002	0.5
White	214 (20.2)	136446 (32.4)		
Black	271 (25.6)	58061 (13.8)		
Asian	6 (0.6)	3314 (0.8)		
Pardo	551 (52.1)	218312 (51.9)		
Indigenous	16 (1.5)	4817 (1.1)		
Literate– no. (%)	1008 (95.3)	254989 (83.0)	<0.001	0 (Δ =14.9)
Health worker – no. (%)	45 (4.3)	5572 (1.3)	<0.001	0 (Δ =7.0)
Co-morbidities^a – no. (%)			<0.001	0 (Δ =0.38)
Cancer	10 (1.0)	3933 (0.9)		
Chronic Obstructive Pulmonary Disease/Emphysema	6 (0.6)	650 (0.1)		
Kidney disease	5 (0.5)	328 (0.1)		
Hypertension	81 (8.2)	46668 (10.2)		
Others	66 (6.7)	66568 (14.6)		
No comorbidity	817 (82.9)	337715 (74.1)		
Diabetes – no. (%)	250 (24.0)	33961 (8.0)	<0.001	0 (Δ =11.3)
HIV infection – no. (%)	220 (21.0)	49046 (13.5)	<0.001	0 (Δ =11.3)
Antiretroviral therapy (ART)^b – no. (%)	193 (87.7)	16592 (33.8)	<0.001	0.5
Alcohol consumption – no. (%)	889 (83.9)	82842 (19.5)	<0.001	0 (Δ =31.2)
Illicit drug use – no. (%)	361 (34.2)	64919 (15.6)	<0.001	0 (Δ =12.5)
Smoking – no. (%)	555 (52.4)	102030 (24.3)	<0.001	0 (Δ =14.7)
Prior TB – no. (%)	170 (16.2)	88266 (19.4)	0.010	0.5
Abnormal chest x-ray – no. (%)	1027 (97.0)	323940 (93.3)	<0.001	0 (Δ =4.8)
Type of TB^c – no. (%)			0.001	0 (Δ =13.6)
Pulmonary	936 (88.4)	384709 (84.4)		
Extrapulmonary	0 (0.0)	57301 (12.6)		
Pulmonary and Extrapulmonary	123 (11.6)	13629 (3.0)		
Positive AFB – no. (%)	852 (80.8)	225900 (67.9)	<0.001	0.5
Positive culture – no. (%)	1053 (99.9)	93668 (69.1)	<0.001	0 (Δ =47.8)
Drug-susceptibility testing (DST) – no. (%)			<0.001	0 (Δ =14.5)
Rifampicin resistance	5 (0.5)	674 (1.3)		
Isoniazid resistance	51 (4.9)	1774 (3.4)		
Rifampicin-Isoniazid resistance	26 (2.5)	1264 (2.4)		
Any drug resistance ^d	98 (9.4)	1444 (2.7)		
Sensitive	859 (82.7)	47785 (90.3)		
Directly observed treatment (DOT) – no. (%)	740 (70.3)	150016 (48.7)	<0.001	0 (Δ =10.7)
Treatment Outcome^e– no. (%)			<0.001	0 (Δ =13.6)
Cure	435 (69.5)	258355 (66.9)		
Failure	26 (4.2)	7343 (1.9)		
Relapse	9 (1.4)	9039 (2.3)		
Death	52 (8.3)	32972 (8.5)		
Lost to follow-up	92 (14.7)	50228 (13.0)		
Transferred out	12 (1.9)	28198 (7.3)		

Table note: Data represent no. (%) or median with Interquartile range (IQR). $p\delta$ -value: second-generation p-value. Δ = *delta-gap*. See the **Supplemental Figure 2** for interpretation of the $p\delta$ -value. Details of the total data available in the re Report-Brazil and SINAN bases in the **Supplementary Table 2**.

Alcohol consumption: Past or current any consumption of alcohol. Smoking: Past or current cigarette smoker. Illicit drug use: Past or current illicit drug use (marijuana, cocaine, heroin or crack)

^a It did not include DM and HIV. ^bART frequency was calculated among the persons living with HIV. ^cAll individuals from the RePORT cohort had a diagnosis of pulmonary tuberculosis, in some cases with presence in other anatomical sites. ^dAny drug (anti-TB) resistance except rifampicin and isoniazid: Pyrazinamide, ethambutol, streptomycin, kanamycin, ethionamide. ^eIn RePORT-Brazil study, the results of anti-TB treatment are recorded at the last study visit (24 months after the start of treatment). By the time the present analyses were performed, 434 participants had not yet completed the last visit, and analyzes of treatment outcomes did not include such participants.

Abbreviations: TB: Tuberculosis. SINAN - Sistema de Informação de Agravos de Notificação. (Brazilian Notification Information System). AFB: acid fast bacilli. ART: Antiretroviral therapy.

FIGURE LEGENDS**Figure 1. Overview of tuberculosis epidemiology in Brazil (2015-2109).**

(A) Political map of Brazil shows the different 26 Brazilian states and one federal district colored according to the 5 macroregions (geopolitical subdivision). The states that include RePORT research centers are displayed in red. (B) Total number of tuberculosis cases reported in each Brazilian state, between 2015 and 2019. The states that include RePORT research centers are indicated by red bars. The upper right panel shows the total number of tuberculosis cases between 2015 and 2019 in the top 10 Brazilian cities with the highest number of tuberculosis cases reported within the study period. The cities where RePORT has research centers are highlighted by red bars. (C) Tuberculosis incidence (per 100,000 person-years) for the years 2015 -2019, in Brazil the states (left panel) or in the cities (right panel) that host RePORT sites are shown. Data represent incidence and 95% confidence interval. Data were obtained from the National Plan for the End of Tuberculosis as a Public Health Problem (*Plano Nacional pelo Fim da Tuberculose como Problema de Saúde Pública*) (Ministério da Saúde do Brasil, 2017) (D) Total number of TB cases reported by the Brazilian Notification Information System (SINAN; grey bars) and by RePORT sites (red dots and connecting lines), in the study period. Right panel shows a donut pie chart illustrating the percentage that the TB cases recruited in the RePORT Brazil protocol represents among the total number of cases reported in SINAN between 2015 and 2019.

Abbreviations: TB: Tuberculosis. RePORT: Regional Prospective Observational Research for Tuberculosis. SINAN: Sistema de Informação de Agravos de Notificação.




Figure 2. Characteristics of the TB cases in RePORT and SINAN (2015-2019).

(A) Comparison of the age distribution (median and interquartile range values) between sex, diabetes condition and HIV infection among RePORT participants and TB cases reported by SINAN within the study period. (B) Proportion of new and re-treatment cases of tuberculosis. (C) Comparison between RePORT and SINAN TB cases (prior to initiation of anti-TB treatment) in regard to frequency of individuals stratified by acid-fast bacilli (AFB) and *M. tuberculosis* culture results as well as of abnormal chest radiograph presentation. (D) Comparison between RePORT and SINAN TB cases with regard to smoking habit and alcohol consumption (smoking and alcohol: in the past or at the time of evaluation before anti-TB treatment). The proportion of TB-diabetes comorbidity is also shown. See the **Supplemental Figure 1** for interpretation of the $p\delta$ -value.

Abbreviations: Δ = delta-gap. $p\delta$ -value: second-generation p-value. RePORT: Regional Prospective Observational Research for Tuberculosis. SINAN: Sistema de Informação de Agravos de Notificação. TB: Tuberculosis.




Figure 3. Characteristics associated with unfavorable treatment outcomes in tuberculosis patients.

Two logistic regression models (LASSO regression and mixed effects) were performed to evaluate the independent associations between clinical characteristics of tuberculosis patients and antitubercular treatment unfavorable outcomes (failure, death and lost to follow-up) in either RePORT (left panel) or SINAN (right panel; where the CI are present, but narrow due to the large sample size). Optimal parameters were found by a cross-validation step, which was repeated 100 times to stabilize the results (**Supplementary Table 2** for RePORT and/or **Supplementary Table 3** for SINAN) and were included in the adjusted models.

Adjusted model 1: LASSO regression using a cross-validation step

Adjusted model 2: The mixed effects model, including the variables "Brazilian states" as a random effect.

In addition to the p-value, the second-generation p-value ($p\delta$ -value) and the *delta-gap* (Δ) (when applicable) are shown. See the **Supplemental Figure 2** for interpretation of the $p\delta$ -value.

Abbreviations: Δ = delta-gap. $p\delta$ -value: second-generation p-value. RePORT: Regional Prospective Observational Research for Tuberculosis. SINAN: Sistema de Informação de Agravos de Notificação. TB: Tuberculosis.

