

CASE REPORT

Open Access

# Severe phenotype in an apparent homozygosity caused by a large deletion in the *CFTR* gene: a case report

Raisa da Silva Martins<sup>1</sup>, Ana Carolina Proença Fonseca<sup>1</sup>, Franklyn Enrique Samudio Acosta<sup>2,3</sup>, Tania Wrobel Folescu<sup>4</sup>, Laurinda Yoko Shinzato Higa<sup>4</sup>, Izabela Rocha Sad<sup>4</sup>, Célia Regina Moutinho de Miranda Chaves<sup>5</sup>, Pedro Hernan Cabello<sup>1,6</sup> and Giselda Maria Kalil Cabello<sup>1\*</sup>

## Abstract

**Background:** Over 1900 mutations have been identified in the cystic fibrosis conductance transmembrane regulator gene, including single nucleotide substitutions, insertions, and deletions. Unidentified mutations may still lie in introns or in regulatory regions, which are not routinely investigated, or in large genomic deletions, which are not revealed by conventional molecular analysis. The apparent homozygosity for a rare, cystic fibrosis conductance transmembrane regulator mutation screened by standard molecular analysis should be further investigated to confirm if the mutation is in fact homozygous. We describe a patient presenting with an apparent homozygous *S4X* mutation.

**Case presentation:** A 13-year-old female patient of African descent with clinical symptoms of classic cystic fibrosis and a positive sweat test (97 mEq/L, diagnosed at age 3 years) presented with pancreatic insufficiency and severe pulmonary symptoms (initial lung colonization with *Pseudomonas aeruginosa* at age 4 years; forced vital capacity: 69%; forced expiratory volume: 51%; 2011). Furthermore, she developed severe acute lung disease and recurrent episodes of dehydration requiring hospitalization. The girl carried the *CFTR* mutation *S4X* in apparent homozygosity. However, further analysis revealed a large deletion in the second allele that included the region of the mutation. The deletion that we describe includes nucleotides 120–142, which correspond to a loss of 23 nucleotides that abolishes the normal translation initiation codon.

**Conclusion:** This study reiterates the view that large, cystic fibrosis conductance transmembrane regulator deletions are an important cause of severe cystic fibrosis and emphasizes the importance of including large deletions/duplications in cystic fibrosis conductance transmembrane regulator diagnostic tests.

**Keywords:** Cystic fibrosis, *CFTR*, Apparent homozygosity, Large deletion, Severe phenotype, Brazilian patient

## Background

Cystic fibrosis (CF; Omin #219700), the most frequent, life-limiting autosomal recessive disorder among Caucasians, is caused by mutations in the cystic fibrosis conductance transmembrane regulator (*CFTR*) gene [1]. In Europe, the carrier frequency is 1:25, resulting in a disease incidence of 1 in 2500 live births [2]. However, this incidence is quite variable, with a range from 1/500 in Ohio Amish to 1/

90000 in Hawaiian Orientals [3]. The *CFTR* gene is characterized by an extremely large number of mutations (more than 1900) [4], the most common being the *F508del*. Other disease-causing mutations are distributed throughout all regions of the world, often with very low frequencies, making the molecular diagnosis difficult, especially in countries outside the European axis where the *F508del* frequency is relatively low, as observed in countries where ethnic composition is not predominantly Caucasian. Brazil is a typical example where the molecular diagnosis of CF is difficult owing to its multi-ethnic characteristics with a significant African component.

\* Correspondence: gkalil.fiocruz@gmail.com

<sup>1</sup>Laboratório de Genética Humana, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Pavilhão Leônidas Deane sala 611. CEP: 21040-360 Avenida Brasil 4365, Rio de Janeiro, Brazil

Full list of author information is available at the end of the article

The design of diagnostic tests available on the market is usually based on the mutation panel of European or North American populations, which is unsuitable for Brazil. Little is known about the incidence of CF in Brazil, but it does vary among different regions of the country [5-8]. Among the European countries that contributed to the current ethnic composition of our population, the *F508del* allele frequency is 71.8% in Germany, 51% in Italy, 53% in Spain, 57% in Poland, 58% in Slovenia, and 45% in Portugal; the latter is the main Brazilian Caucasian component [9,10]. The migratory waves from different regions of Europe, the intense inter-ethnic gene flow and different proportions of mixing genes of Caucasian, African, and Amerindian origins may explain the variation in the prevalence of the *F508del* mutation among different Brazilian regions [11].

Over the past 10 years, the *F508del* mutation contributed 30% of CF alleles in the State of Rio de Janeiro and disease incidence was estimated at 1/6902 live births [12]. Given the current annual number of births of 220,603 (Records of the Ministry of Health of Brazil, 2011; <http://www2.datasus.gov.br/DATASUS/index.php>), approximately 32 patients are born with CF each year. Until now, mutation screening allowed us to identify 29 different mutant alleles in CF patients from Rio de Janeiro. However, the detection rate is only 60%, which means that 40% of alleles remain unknown.

Over 1900 mutations have been identified throughout the *CFTR* gene, including single nucleotide substitutions, insertions, and deletions. Unidentified mutations may still lie in introns or in regulatory regions, which are not routinely investigated, or in large genomic deletions, which are not revealed by conventional molecular analysis. The apparent homozygosity for a rare *CFTR* mutation screened by standard molecular analysis should be further investigated to confirm if the mutation is in fact homozygous. Apparent homozygosity can result from a mutation of one allele and the presence of a large deletion encompassing the location of the first mutation on the second allele [13].

We describe a patient carrying the *S4X* mutation (CM930095) in apparent homozygosity; further analysis revealed that it was heterozygous with a large deletion that includes nucleotides 120–142, which corresponds to a loss of 23 nucleotides coding the first exon of the *CFTR* gene. This deletion removes a region from position -12 to position 10, including codons 1, 2, 3, and 4 of exon 1.

### Case presentation

A 13-year-old female patient of African descent with clinical symptoms of classic CF, a positive sweat test (97 mEq/L; diagnosed at age 3 years), pancreatic insufficiency (PI), and severe pulmonary symptoms (initial lung

colonization with *Pseudomonas aeruginosa* at the age of 4 years) had a forced vital capacity of 69% and a forced expiratory volume of 51%. Furthermore, the girl developed severe acute lung disease and recurrent episodes of dehydration requiring hospitalization.

### Methods

Informed consent in *CFTR* studies was obtained from the proband and her parents at the time of referral. Genomic DNA was extracted from peripheral blood by a Purelink Genomic DNA Mini Kit (Invitrogen, Carlsbad, CA, USA). The DNA was tested for a proper panel of *CFTR* mutations from Rio de Janeiro [6] and the *S4X* mutation in apparent homozygosity was identified by allele-specific polymerase chain reaction (PCR) amplification. The apparent homozygosity for a rare mutation and the absence of the *S4X* mutant allele in the maternal DNA raised the suspicion that it was a false homozygosity. To evaluate the presence of a large deletion encompassing the same sequence region, amplified products were cloned using the pGEM-T<sup>®</sup> Easy Vector System (Fitchburg WI, USA) following manufacturer recommendations and all plasmids were purified by QIAprep<sup>®</sup> Spin Miniprep Kit (Qiagen, GmbH, Hilden) . Purified plasmids were sequenced by Sanger protocol with the BigDye<sup>®</sup> Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City CA, USA) according to the manufacturer's instructions. Single nucleotide polymorphism from intragenic markers (*TUB9-T854T-M470V*) were genotyped using TaqMan<sup>®</sup> allelic discrimination assays (Assay ID: C\_59055676\_10; C\_\_3021356\_10; C\_\_3021372\_10, respectively, Applied Biosystems). The TaqMan<sup>®</sup> genotyping reaction was amplified on an Applied Biosystems<sup>®</sup> 7500 Fast Real-Time PCR System.

### Results

The suspected presence of a large deletion in exon 1 was confirmed by cloning and sequencing analysis, performed on the DNA samples from both parents and the patient (Figure 1). This deletion removes 23 nucleotides from position 120–142, which affects the 5' untranslated region upstream of exon 1, including the translation initiation codon (codons 133–135). This result confirms our initial suspicion that the patient was compound heterozygous for one rearrangement, instead of being homozygous for a rare allele. Analysis of intragenic markers showed that she was homozygous for the 2-1-2 (*TUB9-T854T-M470V*) haplotype.

### Discussion

When a rare mutation is detected in homozygosity in a molecular analysis the result should be confirmed. The distinction between real and apparent homozygosity has important implications for genetic counseling, prenatal



## Ethics approval

The Ethics Committee of the Institute Oswaldo Cruz/Fiocruz approved this study.

## Competing interests

The authors declare that they have no competing interest.

## Authors' contributions

GMKC conceptualized and designed the study, reviewed the literature, participated in the sequence analysis, and drafted the manuscript. RSM carried out the molecular genetic studies and helped draft the manuscript. ACPF carried out the molecular genetic studies and helped draft the manuscript. FESA contributed to sequence alignment and participated in the draft of the manuscript. TWF carried out data collection and wrote the clinical course. LYSH provided all clinical findings. IRS participated in clinical follow-up. CRMMC assessed the child clinically and participated in clinical follow-up. PHC participated in designing the study, contributed to data analysis, and revised the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

The authors thank Dr. Mário Campos Junior, Laboratório de Genética Humana, Instituto Oswaldo Cruz/Fiocruz, for assistance in the assembly of the Figure. Additionally, they thank the Plataforma de Análises de Fragmentos de Eletroforese Capilar PDTIS/FIOCRUZ for DNA sequencing. Most importantly, we are grateful to the patient and her parents who kindly agreed to participate in this study. This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq, and the Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro – FAPERJ.

## Electronic database

DATASUS: <http://www2.datasus.gov.br/DATASUS/index.php>.

## Author details

<sup>1</sup>Laboratório de Genética Humana, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Pavilhão Leônidas Deane sala 611. CEP: 21040-360 Avenida Brasil 4365, Rio de Janeiro, Brazil. <sup>2</sup>Laboratório Interdisciplinar de Pesquisas Médicas, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil. <sup>3</sup>Laboratório de Parasitologia, Instituto Conmemorativo Gorgas de Estudios de la Salud, Panama City, Panama. <sup>4</sup>Departamento de Pneumologia, Instituto Nacional de Saúde da Mulher, da Criança e do Adolescente Fernandes Figueira, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil. <sup>5</sup>Departamento de Nutrição, Instituto Nacional de Saúde da Mulher, da Criança e do Adolescente Fernandes Figueira, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil. <sup>6</sup>Laboratório de Genética Humana, Universidade do Grande Rio, Unigranrio, Rio de Janeiro, Brazil.

Received: 8 March 2014 Accepted: 26 August 2014

Published: 30 August 2014

## References

1. Welsh MJ, Tsui L-C, Boat TF, Beaudet AL: **Cystic fibrosis**. In *The Metabolic and Molecular Basis of Inherited Disease*. Edited by Scriver CR, Beaudet AL, Sly WS, Valle D. New York: McGraw-Hill; 1995:3799–3876.
2. Hamosh A, Corey M: **Correlation between genotype and phenotype in patients with cystic fibrosis (Cystic Fibrosis Genotype-Phenotype Consortium)**. *N Engl J Med* 1993, **329**:1308–1313.
3. Cutting GR: **Cystic Fibrosis**. In *Principles and Practice of Medical Genetics*. 4th edition. Edited by Rimoin DL, Connor JM, Pyeritz RE, Korf BR. London: Harcourt publishers; 2002:1561–1606.
4. **Cystic fibrosis mutation database**. <http://www.genet.sickkids.on.ca/cftr/>.
5. Bernardino ALF, Ferri A, Passos-Bueno MR, Kim CEA, Nakaie CMA, Gomes CE, Damaceno N, Zatz M: **Molecular analysis in Brazilian cystic fibrosis patients reveals five novel mutations**. *Genet Test* 2000, **4**(1):69–74.
6. Cabello GMK, Cabello PH, Otsuki K, Gombarovits ME, Llerena J Jr, Fernandes O: **Molecular analysis of 23 exons of the CFTR gene in Brazilian patients leads to the finding of rare cystic fibrosis mutations**. *Hum Biol* 2005, **77**:125–135.
7. Raskin S, Phillips JA III, Krishnamani MRS, Vnencak-Jones C, Parker RA, Rosov T, Gombarovits ME, Llerena JC Jr, Fernandes O: **DNA analysis of**

**cystic fibrosis in Brazil by direct PCR amplification from Guthrie Cards**. *Am J Med Genet* 1993, **46**:665–669.

8. Streit C, Burlamaque-Neto AC, Abreu e Silva F, Giugliani R, Pereira MLS: **CFTR gene: molecular analysis in patients from South Brazil**. *Mol Genet and Metab* 2003, **78**:259–264.
9. Cystic Fibrosis Genetic Analysis Consortium (CFGAC) 1994: **Population variation of common cystic fibrosis mutations**. *Hum Mutat* 1996, **4**:167–177.
10. Estivill X, Bancells C, Ramos C, Biomed CF Mutation Analysis Consortium: **Geographic distribution and regional origin of 272 cystic fibrosis mutations in European populations**. *Hum Mutat* 1997, **10**:135–154.
11. Cabello GMK, Cabello PH, Llerena JC, Fernandes O: **Polymorphic markers suggest a gene flow of CFTR gene from Sub-Saharan/Arabian and Mediterranean to Brazilian population**. *J Hered* 2006, **97**:313–317.
12. Cabello GMK, Moreira AF, Horovitz D, Correia P, Santa Rosa A, Llerena J Jr, Greg J, Grody WW, Degraive WM, Fernandes O, Cabello PH: **Cystic fibrosis: low frequency of DF508 mutation in 2 population samples from Rio de Janeiro**. *Brazil Hum Biol* 1999, **71**(2):189–196.
13. Hantash FM, Rebuyon A, Peng M, Redman JB, Sun W, Strom CM: **Apparent homozygosity of a novel frame shift mutation in the CFTR gene because of a large deletion**. *J Mol Diag* 2009, **11**(3):253–256.
14. Ramalho AS, Lewandowska MA, Farinha CM, Mendes F, Gonçalves J, Barreto C, Harris A, Amaral MD: **Deletion of CFTR translation start site reveals functional isoforms of the protein in CF patients**. *Cell Physiol Biochem* 2009, **24**:335–346.
15. Cabello GMK, Cabello PH, Lopez-Camelo JS, Llerena JC, Fernandes O: **Haplotype distribution and linkage disequilibrium between four polymorphic markers with CFTR locus in Brazilian cystic fibrosis patients**. *Hum Biol* 2005, **77**:853–865.
16. **Provean (protein variation effect analyzer)**. <http://provean.jcvi.org/>. 2013.

doi:10.1186/1756-0500-7-583

**Cite this article as:** da Silva Martins et al: Severe phenotype in an apparent homozygosity caused by a large deletion in the CFTR gene: a case report. *BMC Research Notes* 2014 **7**:583.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

