

References

- Patel G, Bonomo RA. 'Stormy waters ahead': global emergence of carbapenemases. *Front Microbiol* 2013; **4**: 48.
- Nordmann P, Poirel L, Walsh TR et al. The emerging NDM carbapenemases. *Trends Microbiol* 2011; **19**: 588–95.
- Fischer J, Rodríguez I, Schmoger S et al. *Escherichia coli* producing VIM-1 carbapenemase isolated on a pig farm. *J Antimicrob Chemother* 2012; **67**: 1793–95.
- Fischer J, Rodríguez I, Schmoger S et al. *Salmonella enterica* subsp. *enterica* producing VIM-1 carbapenemase isolated from livestock farms. *J Antimicrob Chemother* 2013; **68**: 478–80.
- Wang Y, Wu C, Zhang Q et al. Identification of New Delhi metallo- β -lactamase 1 in *Acinetobacter lwoffii* of food animal origin. *PLoS One* 2012; **7**: e37152.
- Zhang WJ, Lu Z, Schwarz S et al. Complete sequence of the *bla*_{NDM-1}-carrying plasmid pNDM-AB from *Acinetobacter baumannii* of food animal origin. *J Antimicrob Chemother* 2013; **68**: 1681–2.
- Cabanes F, Lemant J, Picot S et al. Emergence of *Klebsiella pneumoniae* and *Salmonella* metallo- β -lactamase (NDM-1) producers on Reunion Island. *J Clin Microbiol* 2012; **50**: 3812.
- Carattoli A. Plasmids and the spread of resistance. *Int J Med Microbiol* 2013; doi:10.1016/j.ijmm.2013.02.001.
- Hendriksen RS, Vieira AR, Karlsmose S et al. Global monitoring of *Salmonella* serovar distribution from the World Health Organization Global Foodborne Infections Network Country Data Bank: results of quality assured laboratories from 2001 to 2007. *Foodborne Pathog Dis* 2011; **8**: 887–900.
- Guenther S, Ewers C, Wieler LH. Extended-spectrum β -lactamases producing *E. coli* in wildlife, yet another form of environmental pollution? *Front Microbiol* 2011; **2**: 246.

J Antimicrob Chemother 2013

doi:10.1093/jac/dkt298

Advance Access publication 18 July 2013

Isolation of NDM-producing *Providencia rettgeri* in Brazil

Ana Paula D'Alincourt Carvalho-Assef^{1*}, Polyana Silva Pereira¹, Rodolpho Mattos Albano², Gabriela Casemiro Berião¹, Thiago Pavoni Gomes Chagas¹, Loeci Natalina Timm³, Renato Cassol Ferreira Da Silva⁴, Diego Rodrigues Falci⁴ and Marise Dutra Asensi¹

¹Laboratório de Pesquisa em Infecção Hospitalar (LAPIH), Instituto Oswaldo Cruz-FIOCRUZ, Rio de Janeiro, RJ, Brazil; ²Departamento de Bioquímica, Instituto de Biologia Roberto Alcântara Gomes, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brazil; ³Fundação Estadual de Produção e Pesquisa em Saúde (FEPPS IPB-LACEN-RS), Porto Alegre, RS, Brazil; ⁴Hospital Nossa Senhora da Conceição, Porto Alegre, RS, Brazil

*Corresponding author. Tel: +55-2125621636; Fax: +55-2125621634; E-mail: anapdca@ioc.fiocruz.br

Keywords: NDM-1, Tn125, Enterobacteriaceae

Sir,

New Delhi metallo- β -lactamase (NDM) was first identified in 2009 in a *Klebsiella pneumoniae* isolated from a patient in a Swedish hospital who had been previously hospitalized in India in 2008.¹ Since then, different bacterial species carrying the *bla*_{NDM} gene have been isolated, mostly in the Indian subcontinent, but also in several other countries around the world. Here, we report the first description (to our knowledge) of NDM-1 in Brazil.

In early 2013, a diabetic patient with peripheral vascular disease was admitted to a public hospital in southern Brazil (Rio Grande do Sul state) with a diabetic foot infection. Three weeks later, he was discharged from the hospital for ambulatory treatment with 500 mg of ciprofloxacin orally every 12 h. However, a month later he returned to the hospital with worsening of clinical signs and underwent toe amputation. Amoxicillin/clavulanate (1000/200 mg) intravenously every 8 h was initiated prior to surgery and maintained until patient discharge. A fragment of soft tissue from the toe sent for culture yielded growth of a carbapenem-resistant *Providencia rettgeri*, in which *bla*_{NDM-1} was detected. The patient recovered well and was discharged. Previous travel history to other countries was not established.

The identification of the isolate was confirmed by the API 20E system (bioMérieux, Marcy l'Étoile, France) and 16S rRNA gene sequencing. It was resistant to imipenem (MIC 4 mg/L) and susceptible to ertapenem (MIC 0.5 mg/L) and meropenem (MIC 0.75 mg/L) using Etest (AB bioMérieux, Solna, Sweden) according to CLSI 2013 breakpoints.² This resistance profile is very unusual in carbapenemase-producing bacteria. However, ertapenem susceptibility was also described in an NDM-producing *P. rettgeri* isolated from Israel.³ Screening for carbapenemase/metallo- β -lactamase production yielded positive results with imipenem/EDTA,⁴ but yielded negative results when using the modified Hodge test. PCR screening for β -lactamase genes (*bla*_{NDM}, *bla*_{KPC}, *bla*_{OXA-48}, *bla*_{VIM}, *bla*_{IMP}, *bla*_{CTX-M}, *bla*_{TEM}, *bla*_{SHV} and *bla*_{GES}) followed by DNA sequencing identified only the presence of *bla*_{NDM-1}.

Antimicrobial susceptibility profiling analysis performed with Etest indicated that the isolate displayed a multidrug resistance profile. According to CLSI 2013,² it was resistant to ceftazidime, cefotaxime and ciprofloxacin (MICs >256, 16 and >32 mg/L, respectively) and showed intermediate susceptibility to gentamicin (MIC 8 mg/L) and susceptibility to amikacin (MIC 3 mg/L). According to EUCAST breakpoints,⁵ it was resistant to tigecycline (MIC 2 mg/L).

Infections due to multidrug-resistant carbapenemase-producing strains are commonly treated with polymyxin B and/or tigecycline. However, this isolate was intrinsically resistant to polymyxin B and presented intermediate susceptibility to tigecycline, creating a serious problem for the choice of therapy.

The presence of visible plasmid DNA in the *P. rettgeri* isolate was not observed using alkaline lysis methodology.⁶ We also performed hybridization experiments to determine whether the *bla*_{NDM-1} gene was located in a low-copy plasmid (not detectable by ethidium bromide staining), but no hybridization was observed. These observations suggest that the *bla*_{NDM-1} gene in this strain is most likely chromosomally integrated.

To characterize the genetic structure surrounding the *bla*_{NDM-1} gene, a tagmentation library from genomic DNA was made with the Nextera XT DNA Sample Preparation Kit (Illumina) and pair-end sequenced on an Illumina Miseq system. Contigs obtained after *de novo* assembling the reads with Velvet algorithms⁷ were

used for Blast searches against GenBank. A contig that carried the NDM gene was then used as a reference to map the reads and check contig assembly with Geneious 6.1.5 (Biomatters, New Zealand). By this approach, we observed that *bla*_{NDM-1} was located inside a composite transposon, named Tn125 (10092 bp), the same as that described in NDM-1-producing *Acinetobacter baumannii* isolated from Germany.⁸ Tn125 is bracketed by two copies of the insertion sequence IS*Aba125*. In the *A. baumannii* isolate, Tn125 was found on the chromosome, corroborating our results.

*bla*_{NDM-1} has been associated with different mobile elements; however, a complete form or variations of IS*Aba125* have been found upstream of the *bla*_{NDM-1} gene, suggesting that IS*Aba125* is responsible for mobilization of the gene. This insertion sequence has primarily been found in *A. baumannii*.⁸

In South America, the first description of NDM was in Uruguay in 2012, detected in the same species identified in our study (*P. rettgeri*).⁹ The Rio Grande do Sul state (Brazil) and Uruguay border one another. However, to assert that Uruguay was the source of NDM, it would be necessary to perform molecular typing of the isolates to assess their clonality. The other reports of NDM in South America were of a *K. pneumoniae* outbreak in Colombia¹⁰ and of *A. baumannii* in Paraguay,⁹ other countries that border Brazil.

This first known detection of NDM in Brazil brings attention to the need for adopting effective measures to control the spread of this important carbapenemase in this country.

Acknowledgements

We thank the Genomic Platform for DNA Sequencing PDTIS (Instituto Oswaldo Cruz).

Funding

This work was funded by research grants from Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq, Fundação Carlos Chagas de Amparo a Pesquisa – FAPERJ and Instituto Oswaldo Cruz – IOC – FIOCRUZ.

Transparency declarations

None to declare.

References

- 1 Yong D, Toleman MA, Giske CG *et al.* Characterization of a new metallo- β -lactamase gene, *bla*_{NDM-1}, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* 2009; **53**: 5046–54.
- 2 Clinical Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing: Twenty-third Informational Supplement M100-S23*. CLSI, Wayne, PA, USA, 2013.
- 3 Gefen-Halevi S, Hindiyeh MY, Ben-David D *et al.* Isolation of genetically unrelated *bla*_{NDM-1}-positive *Providencia rettgeri* strains in Israel. *J Clin Microbiol* 2013; **51**: 1642–3.
- 4 Nordmann P, Poirel L, Carrère A *et al.* How to detect NDM-1 producers. *J Clin Microbiol* 2011; **49**: 718–21.
- 5 EUCAST. *Breakpoint Tables for Interpretation of MICs and Zone Diameters, Version 3.0, January 2013*. http://www.eucast.org/fileadmin/src/medial/PDFs/EUCAST_files/Disk_test_documents/EUCAST_Breakpoint_table_v_3.0.pdf (24 June 2013, date last accessed).

6 Kado CI, Liu ST. Rapid procedure for detection and isolation of large and small plasmids. *J Bacteriol* 1981; **145**: 1365–73.

7 Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 2008; **18**: 821–9.

8 Poirel L, Bonnin RA, Boulanger A *et al.* Tn125-related acquisition of *bla*_{NDM}-like genes in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2012; **56**: 1087–9.

9 PAHO/WHO. *Epidemiological Alert: Nosocomial Transmission of NDM-Type Multiresistant Bacteria*, 19 December 2012. http://new.paho.org/hq/index.php?option=com_docman&task=doc_view&gid=19623&Itemid=11 (11 June 2013, date last accessed).

10 Escobar Pérez JA, Olarte Escobar NM, Castro-Cardozo B *et al.* Outbreak of NDM-1-producing *Klebsiella pneumoniae* in a neonatal unit in Colombia. *Antimicrob Agents Chemother* 2013; **57**: 1957–60.

J Antimicrob Chemother 2013

doi:10.1093/jac/dkt266

Advance Access publication 3 July 2013

Rectal colonization with New Delhi metallo- β -lactamase-1-producing *Escherichia coli* prior to transrectal ultrasound (TRUS)-guided prostate biopsy

Deborah A. Williamson^{1,2*}, Joshua T. Freeman¹, Sally A. Roberts¹, Helen Heffernan², Kristin Dyet², David L. Paterson³, Benjamin A. Rogers³, Hanna E. Sidjabat³ and Jonathan Masters⁴

¹Department of Clinical Microbiology, Auckland District Health Board, Auckland, New Zealand; ²Institute of Environmental Science and Research, Wellington, New Zealand; ³University of Queensland Centre for Clinical Research, Brisbane, Australia; ⁴Department of Urology, Auckland District Health Board, Auckland, New Zealand

*Corresponding author. Department of Clinical Microbiology, Auckland District Health Board, Auckland, New Zealand. Tel: +0064 (9) 307 4949, ext. 23687; Fax: +0064 (9) 307 6922; E-mail: dwilliamson@adhb.govt.nz

Keywords: carbapenemases, antimicrobial resistance, fluoroquinolones

Sir,

The global spread of New Delhi metallo- β -lactamase (NDM)-producing Enterobacteriaceae is of significant public health concern.¹ To date, NDM-producing Enterobacteriaceae have been isolated from numerous geographical regions, including Europe, North America, Australia and New Zealand.¹ In addition to hydrolysing carbapenems, NDM-producing organisms display resistance to a broad range of antimicrobial classes, primarily due to the presence of additional acquired plasmid-associated resistance genes.² As a result, infections caused by these organisms pose a considerable therapeutic challenge.

Transrectal ultrasound (TRUS)-guided prostate biopsy is a commonly performed urological outpatient procedure, with