

## CORRESPONDENCE



## Neurodevelopment in Infants Exposed to Zika Virus In Utero

**TO THE EDITOR:** During the Zika virus (ZIKV) epidemic in Rio de Janeiro from September 2015 through June 2016, a prospective cohort study involving symptomatic pregnant women who had ZIKV infection confirmed by reverse-transcriptase–polymerase-chain-reaction assay was established.<sup>1</sup> The study was approved by the institutional review boards at Fundação Oswaldo Cruz in Rio de Janeiro and the University of California, Los Angeles, and all the women provided written informed consent for themselves and their children.

A total of 182 children who were exposed to ZIKV in utero were followed longitudinally with specialized testing (Fig. S1 in the Supplementary Appendix, available with the full text of this letter at NEJM.org). Of these children, 131 (72%) were brought by their parents for at least one of the following evaluations: brain imaging (in 115 children), complete eye examinations (in 112),<sup>2</sup> assessment with the Bayley Scales of Infant and Toddler Development, third edition (Bayley-III)<sup>3</sup> (in 104), and assessment of brain-stem auditory evoked response (in 49). Brain-imaging studies consisted of transfontanelle cerebral ultrasonography (in 98 children), computed tomography (in 25), magnetic resonance imaging (MRI) (in 47), or all of these tests; 45 children who underwent brain imaging (39%) underwent more than one type of study. Transfontanelle cerebral ultrasonography was usually performed first, and further imaging was performed according to clinical discretion.


Trained personnel assessed 104 children who were between 12 months and 18 months of age and who had similar socioeconomic backgrounds. Bayley-III, a developmental tool validated cross-culturally in Brazil,<sup>4</sup> was used to assess three domains (cognitive, language, and motor functions). A total of 94 children underwent both Bayley-III and neuroimaging assess-

ments. Abnormal findings on neuroimaging were identified in 39 of 115 children (34%) overall and in 35 of 94 children (37%) who also underwent neuropsychological testing. Among 94 children who underwent both neuroimaging and Bayley-III testing, neuroradiologists found that 10 (11%) had structural abnormalities, 5 (5%) had nonstructural abnormalities, and 20 (21%) had abnormal results that were limited to a non-specific T<sub>2</sub>-weighted hypersignal on MRI.

As shown in Figure 1, and in Table S1 in the Supplementary Appendix, of the 94 children who had undergone neuroimaging and Bayley-III testing, 59 (63%) had Bayley-III scores above 85 for all three domains (1 SD below the mean [ $\pm$ SD] score of  $100\pm 15$  [scores range from 55 to 145, with lower scores indicating a greater degree of developmental delay]); 24 (26%) had one or more Bayley-III scores between 85 and 71 (1 to 2 SD below the mean [ $\pm$ SD] score); and 11 (12%) had one or more scores below 70 (2 SD below the mean [ $\pm$ SD] score). No microcephaly was detected and findings on brain imaging were normal in 44 children (47%) with Bayley scores higher than 85, in 13 (14%) with scores between 85 and 71, and in 2 (2%) with scores of 70 or

### THIS WEEK'S LETTERS

- 2377** Neurodevelopment in Infants Exposed to Zika Virus In Utero
- 2379** Mutation Clearance after Transplantation for Myelodysplastic Syndrome
- 2380** Heat-Stable Carbetocin to Prevent Postpartum Hemorrhage
- 2381** A Man with Epistaxis, Pain and Erythema of the Forearm, and Pancytopenia



less in any domain. Conversely, results of brain imaging were abnormal in 15 children (16%) with Bayley-III scores higher than 85, in 11 (12%) with scores between 85 and 71, and in 9 (10%) with scores of 70 or less in any domain.

Children with normal brain imaging were 20% less likely to have a Bayley-III score that was 2 SD below the mean ( $\pm$ SD) score than those with abnormal brain imaging (odds ratio, 0.80, 95% confidence interval, 0.70 to 0.91). Among children with abnormal findings on brain imaging, 7 of 112 (6%) had an abnormal eye examination and 6 of 49 (12%) had an abnormal hearing assessment. Among 131 children who were exposed to ZIKV in utero and who underwent imaging, neurodevelopmental assessment, sensory organ assessment, or all of these tests, 19 (14%) were found to have severe neurodevelopmental delay (2 SD below the mean [ $\pm$ SD] score), sensory organ dysfunction, or both; this rate is higher than that reported in previous studies.<sup>5</sup> Although a significant association was noted between normal results on brain imaging and higher Bayley-III scores, neuroimaging did not predict developmental delay in 2% of children and normal development in 16% of children.

M. Elisabeth Lopes Moreira, M.D.

Fundação Oswaldo Cruz  
Rio de Janeiro, Brazil

Karin Nielsen-Saines, M.D.

David Geffen School of Medicine at the University of California, Los Angeles (UCLA)  
Los Angeles, CA  
knielsen@mednet.ucla.edu

Patricia Brasil, M.D.

Fundação Oswaldo Cruz  
Rio de Janeiro, Brazil

Tara Kerin, Ph.D.

David Geffen School of Medicine at UCLA  
Los Angeles, CA

Luana Damasceno, M.S.

Marcos Pone, M.D.

Liege M.A. Carvalho, M.D.

Sheila M. Pone, M.D.

Zilton Vasconcelos, Ph.D.

Ieda P. Ribeiro, Ph.D.

Andrea A. Zin, M.D.

Fundação Oswaldo Cruz  
Rio de Janeiro, Brazil

Irena Tsui, M.D.

Kristina Adachi, M.D.

David Geffen School of Medicine at UCLA  
Los Angeles, CA

Stephanie L. Gaw, M.D.

University of California, San Francisco, School of Medicine  
San Francisco, CA

Umme-Aiman Halai, M.D.

David Geffen School of Medicine at UCLA  
Los Angeles, CA

Tania S. Salles, M.D.

Denise C. da Cunha, M.D.

Myrna C. Bonaldo, Ph.D.

Fundação Oswaldo Cruz  
Rio de Janeiro, Brazil

Claudia Raja Gabaglia, M.D.

Biomedical Research Institute of Southern California  
Oceanside, CA

Leticia Guida, Ph.D.

Jociele Malacarne, Ph.D.

Roozemerie P. Costa, B.S.

S. Clair Gomes, Jr., Ph.D.

A. Beatriz Reis, M.S.

Fernanda V.M. Soares, Ph.D.

Fundação Oswaldo Cruz  
Rio de Janeiro, Brazil

Renata H. Hasue, Ph.D.

Carolina Y.P. Aizawa, M.S.

Fernanda F. Genovesi, M.S.

University of São Paulo  
São Paulo, Brazil

Mitsue Aibe, M.D.

Fundação Oswaldo Cruz  
Rio de Janeiro, Brazil

Christa Einspieler, Ph.D.

Peter B. Marschik, Ph.D.

Medical University of Graz  
Graz, Austria

J. Paulo Pereira, Jr., M.D.

Elyzabeth A. Portari, M.D.

Fundação Oswaldo Cruz  
Rio de Janeiro, Brazil

Carla Janzen, M.D.

James D. Cherry, M.D.

David Geffen School of Medicine at UCLA  
Los Angeles, CA

Supported by grants from the Departamento de Ciência e Tecnologia (DECIT) do Ministério da Saúde do Brasil (25000.072811/2016-17); Coordenacao de Aperfeiçoamento de Pessoal de Nivel Superior (CAPES) (88887.116627/2016-01); Brazilian National Council for Scientific and Technological Development (CNPq) (441098/2016-9); Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) (E\_18/2015TXB); ZIKAlliance; the Thrasher Research Fund (20164370); the National Institute of Allergy and Infectious Diseases of the National Institutes of Health (NIH) (AI28697 and AI1259534-01); the National Eye Institute of the NIH (AI129847-01); the Wellcome Trust and the United Kingdom Department for International Development (205377-Z16-Z); and the European Union Horizon 2020 research and innovation program (ZikaPLAN grant agreement no. 734584).

Disclosure forms provided by the authors are available with the full text of this letter at NEJM.org.

1. Brasil P, Pereira JP Jr, Moreira ME, et al. Zika virus infection in pregnant women in Rio de Janeiro. *N Engl J Med* 2016;375:2321-34.
2. Zin AA, Tsui I, Rossetto J, et al. Screening criteria for ophthalmic manifestations of congenital Zika virus infection. *JAMA Pediatr* 2017;171:847-54.
3. Bayley N. Bayley scales of infant and toddler development: administration manual. 3rd ed. San Antonio, TX: Harcourt Assessment, 2006.
4. Madaschi V, Macedo EC, Mecca TP, Pauls CS. Bayley-III scales of infant and toddler development: transcultural adaptation and psychometric properties. *Paidéia (Ribeirão Preto)* 2016; 26:189-97.
5. Reynolds MR, Jones AM, Petersen EE, et al. Vital signs: update on Zika virus-associated birth defects and evaluation of all U.S. infants with congenital Zika virus exposure — U.S. Zika Pregnancy Registry, 2016. *MMWR Morb Mortal Wkly Rep* 2017; 66:366-73.

DOI: 10.1056/NEJMc1800098

## Mutation Clearance after Transplantation for Myelodysplastic Syndrome

**TO THE EDITOR:** Duncavage and colleagues (Sept. 13 issue)<sup>1</sup> address the molecular predictors of disease progression and survival among patients with myelodysplastic syndrome (MDS) who underwent allogeneic hematopoietic stem-cell transplantation. With the use of enhanced exome sequencing, they found that detection of at least one mutation with a maximum variant allele frequency of at least 0.5% at day 30 after transplantation was associated with a higher risk of progression than the absence of such a mutation.

The article offers a new perspective on the understanding of the biologic properties of MDS that may be further explored in the context of the unique immunologic milieu observed after transplantation. In fact, it is well known that most of the therapeutic efficacy of allogeneic hematopoietic stem-cell transplantation relies on the graft-versus-leukemia effect,<sup>2</sup> which is also observed in patients with MDS<sup>3</sup> and which often occurs in parallel with graft-versus-host disease (GVHD). In this context, it would be crucial to