

A General Description of Apocynaceae Iridoids Chromatography

Ana Cláudia F. Amaral, Aline de S. Ramos,
 José Luiz P. Ferreira, Arith R. dos Santos,
 Deborah Q. Falcão, Bianca O. da Silva,
 Debora T. Ohana and Jefferson Rocha de A. Silva

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/55784>

1. Introduction

1.1. Iridoids

Iridoids are considered atypical monoterpene compounds, based on a methylcyclopentan-[C]-pyran skeleton, often fused to a six-membered oxygen ring consisting of ten, nine or in rare cases, eight carbon atoms (Figure 1a) [1, 2]. More than 2500 iridoid compounds have been described in nature, with structural differences related mainly to the degree and type of substitution in the cyclopentane ring skeleton [3]. Iridoids can be found in nature as secoiridoids (Figure 1b), a large group characterized by cleavage of the 7,8-bond on the cyclopentane ring, glycosides, mainly 1-*O*-glucosides, and nor-iridoids, originating from oxidative decarboxylation on C₁₀ or C₁₁ (Figure 1) [3, 4].




Figure 1. Basic skeleton a) iridoid; b) seco-iridoid (R=H or glucose)

Iridoids are derived from isoprene units, which are considered the universal building blocks of all terpenoids, formed through intermediates of the mevalonic acid (MVA) pathway in the cytosol, and the novel 2-methyl-D-erythritol 4-phosphate (MEP) pathway in the plastids of plant cells [2, 5, 6]. The participation of two pathways in iridoid biosynthesis has not yet been clarified, but recent analyses have described the major role of the MEP in the yield of the source for the iridoid isoprene units when compared with the MVA pathway [7, 8, 9]. Iridoid biosynthesis shows two pathways, called route I and II, in which secoiridoids and carboxylated or decarboxylated iridoids are formed, respectively. Route I, considered the main pathway, is responsible for yielding the precursor of the carboxylic iridoids, from iridodial which is oxidized a iridotrial and subsequently converted to a series the iridoids, as occurs in loganin, secologanin, derived secoiridoids, and complex indole alkaloids. In route II, the presence of 8-epi-iridodial, 8-epi-iridotrial and 8-epi-deoxyloganic acid have been reported, forming a source of decarboxylated carbocyclic iridoids such as aucubin and catalpol [10, 3, 11, 12].

Iridoids have shown a broad range of biological activities, such as an antibacterial, antifungal, anti-inflammatory, antitumoral, hepatoprotective, cardioprotective, antioxidative, anti-protozoal and anti-insect properties [13, 14, 15, 16, 17, 18, 19]. *In vitro* activities inhibiting the hepatitis C virus, the differentiation of the adipocyte, and PPAR α activation activities have been also described [20, 21].

The distribution of iridoids in the Eudicotyledoneae has potential usefulness in the taxonomy of the families, related to their presence in a restricted number of families. Iridoid are considered good chemotaxonomic markers of different taxonomic groups, and can be used, in combination with order, tribe and family, to establish the phylogenetic relationship [22, 23, 10, 3, 24, 25].

According to an update of the Angiosperm Phylogeny Group (APGIII) [28,29], the presence of iridoids has been reported in approximately fifty plant families, and can be considered as one of the synapomorphies of the Asterid clade. It is divided into Lamiids, which presents iridoids of the Gentiales, Garryales and Lamiales orders, and Campanulids, which presents secoiridoids of the Asterales and Dipsacales orders. The Gentiales order comprises five families: Apocynaceae, Gelsemiaceae, Gentianaceae, Loganiaceae and Rubiaceae (APG III, 2009), highlighted for the diversity of their iridoids [11, 29, 30].

1.2. Apocynaceae family

Apocynaceae is the most important family within this order, with ca 5000 species distributed worldwide. Seventy percent of the genus, and half of the species distributed in the Neotropical region, are found in the native Brazilian flora [31, 33]. Today, five subfamilies are described for the Apocynaceae family: Rauvolfioideae, Apocynoideae, Secamonoideae and Asclepiadoideae. The latter is the major subfamily of the Apocynaceae, and comprises approximately 3000 species divided into 172 genera, distributed mainly in Neotropic areas of South America, such as Brazil, where the highest diversity of the species has been found [32, 33, 34].

The iridoids class has significant distribution within Apocynaceae family, but is concentrated in just a few genera. The most representative of these are: *Plumeria*, *Himatanthus*, *Allamanda*

and *Cerbera*. According to the traditional classification, the family comprises approximately 87 iridoids, the main ones being plumeride, plumericin and isoplumericin. A review of literature comprising works published on the identification of this class of constituents within Apocynaceae showed, as major natural sources, the following species (numbers in brackets indicate species as shown in Table 1) [36]:

- (1). *Plumeria rubra* L. [*P. acuminata* W. T. Aiton; *P. acutifolia* Poir.; *P. bicolor* Ruiz & Pav.]; (2). *P. lancifolia* Müll. Arg.; (3). *P. acutifolia* Poir. [*P. rubra* L.]; (4). *P. alba* L. [*P. alba* var. *jacquiniana* A. DC.; *P. hypoleuca* Gasp.; *P. hypoleuca* var. *angustifolia* Gasp.]; (5). *P. bracteata* A. DC.; (6). *P. obtusifolia* Steud.; (7). *P. obtusa* L. [*P. multiflora* Standl.]; (8). *P. rubra* var. *alba*; (9). *P. multiflora* Standl.; (10). *Allamanda neriifolia* Hook. [*A. cathartica* var. *Schottii* (Pohl) L.H. Bailey & Raffill]; (11). *Allamanda cathartica* L. [*A. grandiflora* (Aubl.) Lam.; *A. schottii* Hook.]; (12). *Alstonia boonei* De Wild. [*A. congensis* Engl.]; (13). *Cerbera manghas* L.; (14). *Alyxia reinwardtii* Blume; (15). *Alstonia scholaris* (L.) R.Br.; (16). *Himatanthus sucuuba* (Spruce ex Müll. Arg.) Woodson [*H. tarapotensis* (K. Schum. Ex Markgr.) Plumel; *Plumeria floribunda* Müll. Arg.; *P. tarapotensis* K. Schum. ex Markgr.]; (17). *Vinca* sp. L.; (18). *Thevetia peruviana* (Pers.) K. Schum. [*T. neriifolia* Juss. ex Steud.]; (19). *Himatanthus phagedaenicus* (Mart.) Woodson [*Plumeria floribunda* var. *crassipes* Müll. Arg.; *P. lancifolia* var. *major* Müll. Arg.; *P. phagedaenica* Mart.]; (20). *Plumeria bicolor* Seem.; (21). *Plumeria acuminata* W.T. Aiton [*P. rubra* L.]; (22). *Cerbera odollam* Gaertn.; (23). *Allamanda schottii* Pohl [*A. brasiliensis* Schott ex Pohl; *A. cathartica* Schrad; *A. neriifolia* Hook.]; (24). *Himatanthus articulatus* (Vahl) Woodson [*H. rigidus* Wild. Ex Roem. & Schult.; *Plumeria articulata* Vahl; *P. drastica* Mart.; *P. microcalyx* Standl.]; (25). *Rauwolfia grandiflora* Mart. Ex A. DC.; (26). *Plumeria inodora* Jacq. [*P. alba* L.; *P. alba* var. *fragrans* Kunth; *P. alba* var. *fragrantissima* G. Don; *P. alba* var. *inodora* (Jacq.) G. Don]; (27). *Himatanthus bracteatus* (A. DC.) Woodson; (28). *Himatanthus stenophyllus* Plumel; (29). *Himatanthus fallax* (Müll. Arg.) Plumel [*Plumeria fallax* Müll. Arg.]; (30). *Himatanthus obovatus* (Müll. Arg.) Woodson; (31). *Allamanda doniana* Müll. Arg.; (32). *Catharanthus roseus* (L.) G. Don; (33). *Nerium indicum* Mill. [*N. oleander* L.]; (34). *Alstonia macrophylla* Wall. ex G. Don; (35). *Winchia calophylla* A. DC.

1.3. Iridoids of Apocynaceae family

IRIDIDS	SPECIES	PLANT MATERIAL	REF.
6''O-acetylplumeride <i>p</i> -E-coumarate	7: <i>Plumeria obtusa</i>	7:leaves	7: [37]; 7: [38]
6''O-acetylplumeride <i>p</i> -Z-coumarate	7: <i>Plumeria obtusa</i>	7:leaves	7: [37]; 7: [38]
13-O-acetylplumeride	10: <i>Allamanda neriifolia</i>	10:stem; 10:leaves	10: [39]; 10: [40]
allamancin	10: <i>Allamanda neriifolia</i>	10:stem	10: [39]; 10: [40]
allamandicin	10: <i>Allamanda neriifolia</i>	10:stem	10: [39]; 10: [41]; 10: [42]
allamandin	1: <i>Plumeria rubra</i> ; 10: <i>A. neriifolia</i> ; 11: <i>A. cathartica</i> ;	1:stem bark; 19,23:stem; 10,11:leaves; 11,16:root	1: [43]; 1: [44]; 1: [48]; 1: [49]; 10: [39]; 11: [41];

IRIDOIDS	SPECIES	PLANT MATERIAL	REF.
	23: <i>A. schottii</i> ; 16: <i>Himatanthus sucuuba</i> ; 19: <i>H.</i> <i>phagedaenicus</i> ;		11: [42]; 16: [50]; 16: [51]; 19: [45]; 19: [46]; 23: [47]
allamandoid	10: <i>Allamanda neriifolia</i>	10: aerial parts	10: [52]; 10: [53]
allamcidin A	1: <i>Plumeria rubra</i> ; 10: <i>Allamanda neriifolia</i>	1: bark; 10: leaves	1: [43]; 1: [44]; 10: [39]; 10: [40]
allamcidin B	1: <i>Plumeria rubra</i> ; 10: <i>Allamanda neriifolia</i>	1: bark; 10: leaves	1: [43]; 1: [44]; 10: [39]; 10: [40]
allamcidin B β -D-glucose	10: <i>Allamanda neriifolia</i>	10: stem	10: [39]; 10: [40]
allamcin	1: <i>Plumeria rubra</i> ; 10: <i>Allamanda neriifolia</i> ; 23: <i>A. schottii</i>	1: stem bark; 10: leaves; 23: stem	1: [43]; 1: [44]; 1: [48]; 10: [39]; 10: [40]; 23: [47]
allamdin	11: <i>Allamanda cathartica</i>	11: root	11: [41]; 11: [42]
allaneroside	10: <i>Allamanda neriifolia</i>	10: leaves	10: [40]; 10: [54]
3,10-bis-O-allosylcerberidol	13: <i>Cerbera manghas</i> ; 22: <i>C.</i> <i>odollam</i>	13,22: leaves	13: [40]; 13: [56]; 13,22: [55]
3-O-allosylcerberidol	13: <i>Cerbera manghas</i> ; 22: <i>C.</i> <i>odollam</i>	13,22: leaves	13: [40]; 13: [56]; 13,22: [55]
3-O-allosylcyclocerberidol	13: <i>Cerbera manghas</i> ; 22: <i>C.</i> <i>odollam</i>	13,22: leaves	13: [40]; 13: [56]; 13,22: [55]
3-O-allosylepoxycerberidol	13: <i>Cerbera manghas</i> ; 22: <i>C.</i> <i>odollam</i>	13,22: leaves	13: [40]; 13: [56]; 13,22: [55]
alstonoside	15: <i>Alstonia scholaris</i>	15: stem	15: [57]; 15: [58]
alyxialactone	14: <i>Alyxia reinwardtii</i> ; 15: <i>Alstonia scholaris</i>	14: leaves; 15: bark	14: [59]; 15: [60]
10-O-benzoyltheveside	13: <i>Cerbera manghas</i>	13: leaves	13: [56]; 13: [61]
10-O-benzoyltheviridoside	13: <i>Cerbera manghas</i> ; 18: <i>Thevetia peruviana</i>	13,18: leaves	13: [56]; 13,18: [62]
boonein	12: <i>Alstonia boonei</i> ; 25: <i>Rauwolfia grandiflora</i>	12: root; 25: bark	12: [40]; 12: [63]; 12: [64]; 25: [1]
13-O-caffeoylplumieride	1: <i>Plumeria rubra</i> ; 3: <i>P.</i> <i>acutifolia</i>	1,3: root	1: [65]; 3: [38]; 3: [40]; 3: [43]; 3: [66]
10-carboxyloganin	13: <i>Cerbera manghas</i>	13: leaves	13: [56]; 13: [62]
cerberic acid	13: <i>Cerbera manghas</i>	13: bark	13: [40]; 13: [56]; 13: [67]
cerberidol	13: <i>Cerbera manghas</i> ; 22: <i>C.</i> <i>odollam</i>	13,22: leaves	13: [40]; 13: [56]; 13,22 [55]
cerberinic acid	13: <i>Cerbera manghas</i>	13: bark	13: [40]; 13: [56]; 13: [67]

IRIDOIDS	SPECIES	PLANT MATERIAL	REF.
cerbinal	13: <i>Cerbera manghas</i>	13:bark; 13:cortex	13: [56]; 13: [67]; 13: [68]
champalinin ([(E)-p-methoxycinnamoyloxy] plumieride)	7: <i>Plumeria obtusa</i>	7:leaves; 7:stem bark	7: [43]; 7: [69]
cyclocerberidol	13: <i>Cerbera manghas</i> ; 22: <i>C. odollam</i>	13,22:leaves	13: [40]; 13: [56]; 13,22: [55]
cyclocerberidol-3-O-β-D-glucoside	13: <i>Cerbera manghas</i>	13:leaves	13: [56]; 13: [62]
deglycosylplumieride	10: <i>Allamanda neriifolia</i>	10:stem	10: [39]
dehydrogardenoside-10	10: <i>Allamanda neriifolia</i>	10: leaves	10: [39]
10-dehydrogeniposide	13: <i>Cerbera manghas</i>	13:leaves	13: [56]; 13: [61]
15-demethylisoplumieride	1: <i>Plumeria rubra</i> ; 16: <i>Himatanthus succuba</i>	1,16:bark; 16:latex	1: [43]; 1,16: [53]; 1,16: [70]; 16: [51]
15-demethylplumieride	1: <i>Plumeria rubra</i> ; 3: <i>P. acutifolia</i> ; 4: <i>P. alba</i> ; 7: <i>P. obtusa</i> ; 16: <i>Himatanthus succuba</i>	1,16:bark; 3,4:leaves; 4:root; 7:aerial parts; 16:latex	1: [44]; 1,4: [43]; 3: [71]; 7: [72]; 16: [51]; 16: [73]
13-deoxyplumieride	1: <i>Plumeria rubra</i> ; 3: <i>P. acutifolia</i>	1,3:root	1: [65]; 3: [38]; 3: [40]; 3: [43]; 3: [66]
β-dihydroplumericin acid	1: <i>Plumeria rubra</i>	1:root	1: [38]; 1: [42]; 1: [43]; 1: [74]
β-dihydroplumericin	1: <i>Plumeria rubra</i> ; 16: <i>Himatanthus succuba</i>	1,16:root; 1:stem bark	1: [38]; 1: [42]; 1: [43]; 1: [48]; 1: [74]; 16: [51]
epiplumeridoid C	1: <i>Plumeria rubra</i>	1: stem bark	1: [48]
epoxycerberidol	13: <i>Cerbera manghas</i> ; 22: <i>C. odollam</i>	13,22:leaves	13: [40]; 13: [56]; 13,22: [55]
epoxycerberidol-3-O-β-D-glucoside	13: <i>Cerbera manghas</i>	13:leaves	13: [56]; 13: [62]
fulvoplumierin	1: <i>Plumeria rubra</i> L., 3: <i>P. acutifolia</i> , 5: <i>P. bracteata</i> , 8: <i>P. rubra</i> var. <i>alba</i> ; 21: <i>P. acuminata</i> ; 16: <i>Himatanthus succuba</i>	1,3,8,21:stem bark; 1,16:bark; 1:root, 1:descorticated stem, 3:rind, 5:root cortex	1: [43]; 1: [48]; 1: [65]; 1: [74]; 1: [75]; 1: [76]; 1: [77]; 1,3,5,8: [38]; 1,3,21: [44]; 3: [78]; 3: [79]; 3: [80]; 3,8: [42]; 3,8: [83]; 5: [81]; 8: [82]; 16: [51]
gardenoside	10: <i>Allamanda neriifolia</i>	10:stem;10:leaves	10: [39]
isoallamandicin	10: <i>Allamanda neriifolia</i>	10:stem	10: [39]; 10: [40]

IRIDOIDS	SPECIES	PLANT MATERIAL	REF.
isoboonein	15: <i>Alstonia scholaris</i> ; 25: <i>Rauwolfia grandiflora</i>	15,25:bark	15: [60]; 25: [1]
isoplumericin	1: <i>Plumeria rubra</i> , 3: <i>P. acutifolia</i> ; 4: <i>P. alba</i> , 7: <i>P. obtusa</i> ; 20: <i>P. bicolor</i> ; 10: <i>Allamanda neriifolia</i> ; 11: <i>A. cathartica</i> ; 23: <i>A. schottii</i> ; 31: <i>A. doniana</i> ; 16: <i>Himatanthus sucuuba</i> ; 19: <i>H. phagedaenicus</i> ; 24: <i>H. articulatus</i> ; 29: <i>H. fallax</i> ; 30: <i>H. obovatus</i> ;	1,16,20: stem bark; 1,4,7,16,30,31:root; 1,3: wood; 10,19,23,29:stem; 11,16:leaves; 16,24:bark; 16:latex	1: [42]; 1: [48]; 1: [58]; 1: [74]; 1: [75]; 1: [84]; 1,3: [85]; 1,4,7: [38]; 1,4,7: [43]; 4,7: [84]; 10: [39]; 11: [49]; 11,29,30,31: [89]; 16: [51]; 16: [88]; 16: [90]; 16: [91]; 16: [92]; 19: [45]; 19: [46]; 20: [86]; 23: [47]; 24: [87]; 30,31: [93]; 30,31: [94]
isoplumieride	1: <i>Plumeria rubra</i> ; 3: <i>P. acutifolia</i> ; 16: <i>H. sucuuba</i> ; 27: <i>H. bracteatus</i> ; 28: <i>H. stenophyllus</i>	1,3:root; 16,27,28:leaves; 16,27,28:bark; 16,27,28:latex	1: [65]; 3: [38]; 3: [40]; 3: [43]; 3: [66]; 16: [51]; 16: [73]; 16,27,28: [95]
loganic acid	25: <i>Rauwolfia grandiflora</i> ; 32: <i>Catharanthus roseus</i>	25: bark; 32:seed	25: [1]; 32: [4]
loganin	12: <i>Alstonia boonei</i> ; 15: <i>A. scholaris</i> ; 13: <i>Cerbera manghas</i> ; 17: <i>Vinca sp.</i> ; 25: <i>Rauwolfia grandiflora</i> ; 32: <i>Catharanthus roseus</i> ; 35: <i>Winchia calophylla</i>	12,13,17:leaves; 15,25:bark; 32:seed; 35:stem bark	12: [64]; 13: [46]; 13: [56]; 13: [61]; 13: [62]; 15: [60]; 17: [42]; 17: [96]; 25: [1]; 32: [4]; 35: [129]
3-O-methylallamancin	10: <i>Allamanda neriifolia</i>	10:leaves	10: [39]; 10: [40]
3-O-methylallamcim	10: <i>Allamanda neriifolia</i>	10:leaves	10: [39]; 10: [40]
naresuanoside	34: <i>Alstonia macrophylla</i>	34:stem	34: [128]
obtusadoid A	7: <i>Plumeria obtusa</i>	7:aerial parts	7: [72]
obtusadoid B	7: <i>Plumeria obtusa</i>	7:aerial parts	7: [72]
plumenoside (β -dihydroplumericinic acid glucosylester)	1: <i>Plumeria rubra</i> ; 3: <i>P. acutifolia</i>	1,3:root	1: [65]; 3: [38]; 3: [40]; 3: [43]; 3: [66]
plumeric acid 1- β -O- β -D-glucopyranosyl	24: <i>Himatanthus articulata</i>	24:bark	24: [87]
plumericidine	1: <i>Plumeria rubra</i>	1: flowers	1: [97]
plumericin	1: <i>Plumeria. rubra</i> , 3: <i>P. acutifolia</i> ; 4: <i>P. alba</i> , 5: <i>P. bracteata</i> , 7: <i>P. obtusa</i> ,	1,16,20:stem bark; 1,4,7,8,9,16,33:root; 1,3:wood; 5,8,16:root cortex; 1,19,23,29:stem;	1: [42]; 1: [48]; 1: [58]; 1: [74]; 1: [75]; 1: [82]; 1: [98]; 1,3: [85]; 1, 4, 7: [43]; 1,4,5,7,8,9: [38];

IRIDOIDS	SPECIES	PLANT MATERIAL	REF.
	8: <i>P. rubra</i> var. <i>alba</i> ; 9: <i>P. multiflora</i> ; 20: <i>P. bicolor</i> ; 21: <i>P. acuminata</i> ; 10: <i>Allamanda neriifolia</i> ; 11: <i>A. cathartica</i> ; 23: <i>A. schottii</i> ; 31: <i>A. doniana</i> ; 16: <i>Himatanthus</i>	1,10,11,16,23:leaves; 1,16,20,24:bark; 1:fruits;	1,21: [44]; 4,7: [84]; 5: [81]; 8: [82]; 9: [99]; 10: [39]; 11: [49]; 11,29,30,31: [89]; 16: [50]; 16: [51]; 16: [88]; 16: [90]; 16: [91]; 16: [92]; 16: [100]; 19: [45]; 19: [46]; 20: [86]; 20: [101]; 23: [47]; 23: [102]; 24: [87]; 33: [84]; 33: [130]
plumeridoid A	1: <i>Plumeria rubra</i>	1: stem bark	1: [48]
plumeridoid B	1: <i>Plumeria rubra</i>	1: stem bark	1: [48]
plumeridoid C	1: <i>Plumeria rubra</i> ; 16: <i>Himatanthus succuba</i>	1,16: stem bark	1: [48]; 16: [103]
plumiepoixide	10: <i>Allamanda neriifolia</i>	10:stem; 10:leaves	10: [39]; 10: [40]
plumieride	1: <i>Plumeria rubra</i> L., 2: <i>P. lancifolia</i> , 3: <i>P. acutifolia</i> , 4: <i>P. alba</i> , 5: <i>P. bracteata</i> , 6: <i>P. obtusifolia</i> , 7: <i>P. obtusa</i> , 8: <i>P. rubra</i> var. <i>alba</i> ; 20: <i>P. bicolor</i> ; 26: <i>P. inodora</i> ; 10: <i>Allamanda neriifolia</i> ; 11: <i>A. cathartica</i> ; 23: <i>A. schottii</i> ; 16: <i>Himatanthus succuba</i> ; 24: <i>H. articulatus</i> ; 27: <i>H. bracteatus</i> ; 28: <i>H. stenophyllus</i> ; 29: <i>H. fallax</i>	1,2,7,8:stem bark, 1,3,4,8:root, 1,2,3,4,5,7,8,10,11,16,27 ,28:leaves, 1,8:stem wood, 1,2,3,5:wood, 1,4,7:seed, 1,2,3,4,5,7:flowers, 1:descorticated stem, 2,3,5:root cortex, 2,3,5:pith, 2,3,4,5:cortex, 3:rind; 10,20,23,26,29: stem; 16,20,24,27,28: bark; 16,23,27,28: latex	1: [42]; 1: [48]; 1: [65]; 1: [75]; 1: [76]; 1: [77]; 1: [82]; 1: [84]; 1: [105]; 1: [111]; 1,2,3,4,5,6,7,8,; [38]; 1,4,7,20: [86]; 1,3: [85]; 1,8: [104]; 2: [81]; 3: [66]; 3: [80]; 3: [81]; 3: [106]; 3: [107]; 4: [84]; 4: [108]; 4: [109]; 5: [81]; 6: [110]; 7: [84]; 8: [105]; 10: [39]; 11: [49]; 11: [112]; 16: [51]; 16: [73]; 16,27,28: [95]; 20: [114]; 23: [102]; 24: [87]; 24: [115]; 26: [113]; 29: [89]
plumieride coumarate (13-O-coumaroylplumieride; plumieride <i>p-E</i> -coumarate)	1: <i>Plumeria rubra</i> ; 3: <i>P. acutifolia</i> ; 4: <i>P. alba</i> ; 7: <i>P. obtusa</i> ; 10: <i>Allamanda neriifolia</i> ; 11: <i>A. cathartica</i>	1:bark; 1,3,4,7,11:root; 1,4,7:stem; 1,4,7,10: leaves; 1,7: flowers; 1,3: wood	1: [44]; 1,3: [85]; 1,3,4,7: [38]; 1,3,4,7: [43]; 1,4,7: [84]; 3: [66]; 7: [37]; 10: [39]; 11: [89]; 11: [116]
plumieride coumarate glucoside (protoplumericin A; 13-O- <i>p</i> -O-glucosylcoumaroyl plumieride)	1: <i>Plumeria rubra</i> ; 3: <i>P. acutifolia</i> ; 4: <i>P. alba</i> , 7: <i>P. obtusa</i> ; 10: <i>Allamanda neriifolia</i> ; 11: <i>A. cathartica</i>	1,3,4,11:root; 1,3:wood, 4,7,10:stem; 4,7,10:leaves; 4,7:flowers; 1:bark	1,3: [85]; 1,3,4,7: [38]; 1,4,7: [43]; 1,4,7: [84]; 1,7,11: [118]; 3: [66]; 10: [39]; 10: [40]; 10: [116]; 10: [117]; 11: [89]
plumieride <i>p-Z</i> -coumarate	7: <i>Plumeria obtusa</i>	7:leaves	7: [37]; 7: [38]

IRIDOIDS	SPECIES	PLANT MATERIAL	REF.
plumieride-1 α	1: <i>Plumeria rubra</i> ; 3: <i>P. acutifolia</i> ; 7: <i>P. obtusa</i>	1,3:root; 7:leaves; 7:aerial parts	1: [65]; 3: [38]; 3: [40]; 3: [66]; 7: [72]; 7: [119]
plumieride-1 β -O- β -D-glucopyranosyl	24: <i>Himatanthus articulatus</i>	24:bark	24: [87]
plumieridin A	1: <i>Plumeria rubra</i> ; 7: <i>Plumeria obtusa</i>	1:flowers; 7:aerial parts	1: [43]; 1: [118]; 1: [120]; 7: [72]
plumieridin B	1: <i>Plumeria rubra</i>	1: flowers	1: [43]; 1: [120]
protoplumericin A-1 α	1: <i>Plumeria rubra</i> ; 3: <i>P. acutifolia</i> ; 10: <i>Allamanda neriifolia</i>	1, 3:root; 10:stem; 10:leaves	1: [65]; 3: [38]; 3: [40]; 3: [66]; 10: [39]
protoplumericin B (13-O- <i>p</i> -O-glucosylcaffeoyl-plumieride)	10: <i>Allamanda neriifolia</i>	10:stem; 10:leaves	10: [39]; 10: [40]
pulosarioside	14: <i>Alyxia reinwardtii</i>	14:bark	14: [40]; 14: [121]
scholarein A	15: <i>Alstonia scholaris</i>	15:bark	15: [60]
scholarein B	15: <i>Alstonia scholaris</i>	15:bark	15: [60]
scholarein C	15: <i>Alstonia scholaris</i>	15:bark	15: [60]
scholarein D	15: <i>Alstonia scholaris</i>	15:bark	15: [60]
secologanin	32: <i>Catharanthus roseus</i>	32:leaves	32: [4]
sweroside	34: <i>Alstonia macrophylla</i>	34:stem	34: [128]
theside	13: <i>Cerbera manghas</i> ; 18: <i>Thevetia peruviana</i>	13:fruit; 13,18:leaves; 18:root	13: [46]; 13: [56]; 13: [61]; 13: [124]; 13,18: [42]; 13,18: [62]; 18: [122]; 18: [123]; 18: [125]
theviridoside	13: <i>Cerbera manghas</i> ; 18: <i>Thevetia peruviana</i>	13:fruit; 13:cortex; 13,18:leaves; 18:root	13: [46]; 13: [56]; 13: [61]; 13: [68]; 13: [124]; 13,18: [42]; 13,18: [62]; 18: [125]; 18: [126]
theviridoside-10-O- β -D-fructofuranosyl	18: <i>Thevetia peruviana</i>	18:leaves; 18:root	18: [125]; 18: [127]
theviridoside-10-O- β -D-glucopyranosyl	18: <i>Thevetia peruviana</i>	18:root	18: [125]
theviridoside-3'-O- β -D-glucopyranosyl	18: <i>Thevetia peruviana</i>	18:root	18: [125]
theviridoside-6'-O- β -D-glucopyranosyl	18: <i>Thevetia peruviana</i>	18:leaves; 18:root	18: [125]; 18: [127]

Table 1. Iridoids found in Apocynaceae family.

2. Extracts preparation and iridoids isolated

Table 2 illustrates the extraction methods and the iridoids isolated of some genera of Apocynaceae.

SPECIES / PLANT MATERIAL	SOLVENT/ EXTRACTION METHOD	ISOLATED IRIDOIDS	REF.
<i>Allamanda neriifolia</i> Hook./ stem and leaves	MeOH/ Percolation	Isoallamandicin, allamcin, 3-O-methylallamcin, allamancin, 3-O-methylallamancin, allamcidin, plumericin, isoplumericin, allamandin, allamandicin, plumieride 13-O-acetate, deglucosylplumieride, 13-O-coumaroylplumieride, allamcidin B β -D-glucoside, plumiepoixide, protoplumericin B, plumieride, protoplumericin A, gardenoside, 10-dehydrogardenoside	[39]
<i>A. neriifolia</i> Hook./ aerial parts	EtOH/ Maceration	Allamanoid, plumieride, protoplumericin, nicotiflorin	[52]
<i>A. neriifolia</i> Hook./ stem	MeOH/ Percolation	Plumericin, plumieride, protoplumericin	[117]
<i>A. schottii</i> Pohl/ stem	EtOH/ Maceration	Allamandin, allamcin, plumericin, isoplumericin, scoparone, scopoletin, pinosresinol	[47]
<i>A. cathartica</i> L. / root bark	EtOH/ Hot extraction	Allamandin, allamandicin, allamandin, plumericin, isoplumericin	[41]
<i>A. cathartica</i> L. /root	MeOH/ Hot extraction	Isoplumericin, plumericin, plumieride, plumieride coumarate, plumieride coumarate glucoside	[116]
<i>A. cathartica</i> L. / root bark	MeOH and CHCl ₃ / Not described	Plumericin, isoplumericin, plumieride, plumieride coumarate, plumieride coumarate glucoside	[84]
<i>A. cathartica</i> L./ root (inner part)	MeOH and CHCl ₃ / Soxhlet apparatus	Plumieride, plumieride coumarate, plumieride coumarate glucoside	[84]
<i>A. cathartica</i> L. / leaves	EtOAc/ Not described	Isoplumericin, plumericin	[89]
<i>Alstonia macrophylla</i> Wall. ex G. Don / stem	EtOH/ Maceration	Sweroside, naresuanoside	[128]
<i>A. scholaris</i> (L.) R. Br./ bark	EtOH/ Maceration	Scholarein A, scholarcin B, scholarcin C, scholarcin D, isoboonein, alyxialactone, loganin	[60]


SPECIES / PLANT MATERIAL	SOLVENT/ EXTRACTION METHOD	ISOLATED IRIDOIDS	REF.
<i>Alyxia reinwardtii</i> Blume/ leaves	MeOH/ Maceration	Alyxialactone, 4-epi-alyxialactone	[59]
<i>Cerbera manghas</i> L./ leaves	MeOH/ Not described	Theviridoside, theveside	[124]
<i>C. manghas</i> L./ stem bark and root bark	MeOH/ Percolation	Cerbinal, cerberic acid, cerberinic acid	[67]
<i>C. manghas</i> L./ leaves	MeOH/ Percolation	Cerberidol, epoxycerberidol, cyclocerberidol, cerberidol-3-O- β -D-allopyranoside, cerberidol-3,10-bis-O- β -D- allopyranoside, epoxycerberidol-3-O- β -D-allopyranoside, cyclocerberidol-3-O- β -D-allopyranoside	[55]
<i>C. manghas</i> L./ leaves	MeOH/ Percolation	Theviridoside, 10-carboxyloganin, loganin, cyclocerberidol-3-O- β -D-glucoside, epoxycerberidol-3-O- β - D-glucoside	[62]
<i>C. manghas</i> L./ leaves	MeOH/ Percolation	10-O-benzoyltheveside, 10-dehydrogeniposide, loganin, theviridoside, theveside	[61]
<i>C. odollam</i> Gaertn./ leaves	MeOH/ Percolation	Cerberidol, cyclocerberidol, cerberidol-3-O- β -D- allopyranoside, cyclocerberidol-3-O- β -D-allopyranoside	[55]
<i>Himatanthus articulatus</i> (Vahl) Woodson/ bark	MeOH/ Maceration	1 β -O- β -D-glucopyranosylplumeric acid, plumeride-1 β -O- β - -D-glucopyranosyl, plumericin, isoplumericin	[87]
<i>H. fallax</i> (Müll. Arg.) Plumel/ stem	EtOAc/ Maceration	Isoplumericin, plumericin, plumeride	[89]
<i>H. sucuuba</i> (Spruce ex Müll. Arg.) Woodson / latex	H ₂ O/ Not described	15-demethylisoplumeride, 15- demethylplumeride, plumeride, isoplumeride	[70]
<i>H. sucuuba</i> (Spruce ex Müll. Arg.) Woodson/ bark	EtOH/ Maceration	Isoplumericin, plumericin	[88]
<i>H. sucuuba</i> (Spruce ex Müll. Arg.) Woodson / latex	H ₂ O/ Not described	Plumericin, isoplumericin	[92]
<i>H. sucuuba</i> (Spruce ex Müll. Arg.) Woodson /	BuOH/ Partition	Plumeride, isoplumeride, 15-demethylplumeride	[73]

SPECIES / PLANT MATERIAL	SOLVENT/ EXTRACTION METHOD	ISOLATED IRIDOIDS	REF.
latex			
<i>H. succuba</i> , (Spruce ex Müll. Arg.) Woodson / bark	EtOH/ Maceration	Plumericin	[100]
<i>Nerium indicum</i> Mill. / root	Petrol/ Soxhlet apparatus	Plumericin	[130]
<i>Plumeria acutifolia</i> Poir. / root	MeOH/ Percolation	Plumericin, 13-O-coumaroylplumieride, 13-O-caffeoylplumieride, 13-deoxyplumieride, plumenoside, 1 α -plumieride, 1 α -protoplumericin A, plumieride, protoplumericin A, 8-isoplumieride	[66]
<i>P. acutifolia</i> Poir. / leaves	MeOH/ Exhaustive Maceration	15-demethylplumieride	[71]
<i>P. bicolor</i> Seem. / bark	MeOH/ Exhaustive Maceration	Plumericin, isoplumericin	[86]
<i>P. rubra</i> L. var. <i>acutifolia</i> (Poir.) Woodson / bark	MeOH/ Maceration	15-demethylisoplumieride, 15-demetilplumieride, plumieride, isoplumieride	[70]
<i>P. rubra</i> L. / steam bark	CH ₂ Cl ₂ :MeOH (1:1) and MeOH/ Maceration	Plumericin, isoplumericin, plumieride, fulvoplumerin	[75]
<i>P. rubra</i> L. / heartwood	EtOAc/ Percolation	Plumericin, isoplumericin, plumieride, 13-O-coumaroylplumieride, protoplumericine A	[85]
<i>P. rubra</i> L. / stem bark	MeOH/ Maceration	Fulvoplumerin, allamandin, allamcin, plumericin, 15-demethylplumieride, plumieride, α -allamcinidin, β -allamcinidin, 13-O- <i>trans</i> -p-coumaroylplumieride	[44]
<i>P. rubra</i> L. / stem bark	MeOH/ Maceration	Fulvoplumerin, dihydroplumericin, plumieride, plumeridoids A, B and C, isoplumericin, plumericin, allamcin, allamandin, mixture of plumeridoid C and epiplumeridoid C	[48]
<i>P. rubra</i> L. var. <i>acutifolia</i> (Poir.) Woodson /	EtOH/ Maceration	Plumieridin A, plumieridin B, plumericin, plumieride	[120]


SPECIES / PLANT MATERIAL	SOLVENT/ EXTRACTION METHOD	ISOLATED IRIDOIDS	REF.
flowers			
<i>P. rubra</i> L. var. <i>acutifolia</i> (Poir.) Woodson/ flowers	EtOH/ Maceration	Plumericidine	[97]
<i>P. obtusa</i> L. / flower	EtOH/ Maceration	Plumieride coumarate glucoside	[118]
<i>P. obtusa</i> L. / aerial parts	MeOH/ Maceration	Obtusadoids A and B, plumieridin A, plumieridine, 1 α -plumieride, 15-demethylplumieride	[72]
<i>P. obtusa</i> L. / leaves	MeOH/ Maceration	6''-O-acetylplumieride <i>p</i> -E-coumarate, 6''-O-acetylplumieride, <i>p</i> -Z-coumarate, plumieride, plumieride <i>p</i> -Z-coumarate, plumieride <i>p</i> -E-coumarate	[37]
<i>P. obtusa</i> L. / leaves	MeOH/ Maceration	1 α -plumieride	[119]
<i>P. inodora</i> Jacq. / stems	H ₂ O/ Maceration	Plumieride	[113]
<i>P. bicolor</i> Seem. / bark	MeOH/ Maceration	Plumieride	[114]
<i>Rauwolfia grandiflora</i> Mart. Ex A. DC. / bark	EtOH/ Maceration	Loganic acid, loganin, boonein, isoboonein	[1]
<i>Thevetia peruviana</i> (Pers.) K. Schum. / Leaves	MeOH/ Percolation	10-O- β -D-fructofuranosyltheviridoside, 6'O- β -D-glucopyranosyltheviridoside	[125]
<i>T. peruviana</i> (Pers.) K. Schum. / root	MeOH/ Percolation	Theveside, theviridoside, 10-O- β -D-fructofuranosyltheviridoside, 6'O- β -D-glucopyranosyltheviridoside, 10-O- β -D-glucopyranosyltheviridoside, 3'O- β -D-glucopyranosyltheviridoside	[127]
<i>Winchia calophylla</i> A. DC. / stem bark	EtOH/ Hot extraction	Loganin	[129]

Table 2. Main extraction methods to obtain iridoids of Apocynaceae family.


3. Chemical structures




1 *Plumericin* R = H; R¹ = CH₃



3 *Allamcin*




4 *Allamandin*




5 *Fulvoplumierin*


2 *Isoplumericin* R = CH₃; R¹ = H



6 *Plumieride* R = H; R¹ = CH₃




8 *Isoplumieride*




9 *Loganin*

7 *15-demethylplumieride* R=R¹= H



10 *Plumieride coumarate*



11 *Plumieride coumarate glucoside*

Figure 2. The most isolated iridoids of Apocynaceae family

4. Chromatographic separation

4.1. Thin layer chromatography

Preparative chromatography was performed with thin layer chromatography (TLC) aluminum sheets and 8:2 chloroform/ methanol as mobile phase of a fraction from the 95% EtOH extract of stems of *Alstonia macrophylla*. This procedure led to the isolation of the compounds sweroside (2 mg) and naresuanoside (3 mg) [128]

Another interesting application of preparative chromatography is described in [47] with ethanol extract of ground stems of *A. schottii*. This extract was fractionated by bioassay monitoring and after successive liquid-liquid partition and flash column chromatography; the authors obtained the iridoids allamandin; allamcin and a mixture of plumericin and isoplumericin. This separation was performed with a Chromatotron rotor (silica gel (Si gel), 2-mm) with 1 % methanol in chloroform as solvent system.

The bioassay-guided fractionation of the extracts of *Plumeria rubra* barks also proved to be a successful strategy, leading to the isolation of eleven substances, of which eight were iridoids. In this context, the aqueous extract of *Plumeria rubra* bark has iridoids such as the epimers, α -allamcidin and β -allamcidin, which were resolved by preparative TLC on Si-gel G plates (20 × 20 cm, 250 μ m, Merck®), using chloroform/ethyl acetate/methanol (3:3:1) as solvent system [44].

Ferreira and coworkers [95] describe the HPTLC analysis of the bark, latex and leaf extracts and substances of *Himatanthus sucuuba*. The solutions of the extracts in methanol (10 mg mL⁻¹) and the isolated iridoids (3 mg mL⁻¹ - plumieride; 1 mg mL⁻¹ - isoplumieride) were applied (5 μ L) using Linomat IV, a Camag semi-automatic spotter. The analysis was carried out on a precoated silica gel 60 F254 (Merck) HPTLC plate (0.2 mm of layer thickness and 10 × 10 cm size) using chloroform/ methanol (8:2) as developing system. The resulting chromatogram was dried and the spots were visualized by spraying with vanillin – sulfuric acid solution, followed by heating at 100 °C. Table 3 shows other studies using TLC analysis of Apocynaceae iridoids.

SPECIES	PLANT MATERIAL	IRIDOIDS	STATIONARY PHASE	MOBILE PHASE	REF.
<i>Plumeria obtusa</i>	Flowers	Plumieride coumarate glucoside	Si gel 60G F ₂₅₄	CHCl ₃ -MeOH (3:1)	[118]
<i>Plumeria obtusa</i>	Leaves	6 ⁿ -O-acetylplumieride <i>p</i> -E-coumarate	1- Si gel GF ₂₅₄	1- CHCl ₃ -MeOH (8.5:1.5)	[37]
		6 ⁿ -O-acetylplumieride <i>p</i> -Z-coumarate	2- Si gel	2- Gradients of Petrol-EtOAc	
<i>Plumeria rubra</i>	Stem bark	Plumeridoid A	1- Si gel GF ₂₅₄	1- Hexane-EtOAc (20%)	[48]
		Plumeridoid B	2- Si gel GF ₂₅₄	2- Hexane-EtOAc (4%)	
<i>Himatanthus fallax</i>	Stems	Plumericin	Si gel	10% Me ₂ CO in CHCl ₃	[89]
		Isoplumericin			
		Plumieride		CHCl ₃ -MeOH (85:15)	

SPECIES	PLANT MATERIAL	IRIDOIDS	STATIONARY PHASE	MOBILE PHASE	REF.
<i>Himatanthus succuba</i>	Latex	Plumericin	Si gel GF ₂₅₄	Hexane-EtOAc (6:4)	[92]
		Isoplumericin			
<i>Alyxia reinwardti</i>	Leaves	Alyxialactone	Si gel GF ₂₅₄	CHCl ₃ -MeOH (95:5)	[59]
		4- <i>epi</i> -alyxialactone			
<i>Allamanda cathartica</i>	Roots	Plumericin	1- Si gel	1- CHCl ₃	[41]
		Isoplumericin			
		Allamandicin	2- Si gel	2- CHCl ₃ -MeOH (95:5)	
		Allamandin	3- Si gel	3- ether-hexane (1:1)	

Table 3. TLC analyses of Apocynaceae iridoids.

4.2. Open column chromatography

The methanol extract of the leaves of *Cerbera manghas* and its fruit contain the iridoids theveside and theviridoside, as described in [124]. The methanol extract of the leaves, after the addition of water, was sequentially partitioned with chloroform, acetic acid and butanol. This extract and the aqueous phase were submitted to column chromatography with charcoal and water/methanol as eluent. Theveside was isolated from the aqueous phase. Fractions of the butanol extract, which turn blue after heating with mineral acid, were chromatographed over a silica gel column with a gradient of increasing polarity of chloroform/methanol to afford theviridoside.

Cerberidol, epoxycerberidol, cyclocerberidol, cerberidol-3-*O*- β -D-allopyranoside, cerberidol-3,10-bis-*O*- β -D-allopyranoside, epoxycerberidol-3-*O*- β -D-allopyranoside and cyclocerberidol-3-*O*- β -D-allopyranoside are present in the leaves of *C. manghas* [55]. The methanol extract from 1.45 kg of dried leaves was concentrated, re-suspended in water, and sequentially partitioned with benzene and chloroform. The aqueous phase was chromatographed using MCI-gel column (elution with gradients of water/methanol) and the fraction eluted with 20% methanol was chromatographed in two steps: using RQ-1 (Fuji-gel phase) column (elution with water/acetonitrile) and using a silica gel column (elution with gradient of chloroform/methanol/water 7:3:1 to 7:3:1.2). Fractionation afforded cerberidol (75 mg), epoxycerberidol (10 mg), cyclocerberidol (160 mg), cerberidol-3-*O*- β -D-allopyranoside (135 mg), cerberidol-3,10-bis-*O*- β -D-allopyranoside (20 mg), epoxycerberidol-3-*O*- β -D-allopyranoside (27 mg) and cyclocerberidol-3-*O*- β -D-allopyranoside (250 mg). Some of these iridoids are also found in dried leaves of *Cerbera odollam* Gaertn [55]. The same methodology was used to isolate cerberidol (15 mg), cyclocerberidol (48 mg), cerberidol-3-*O*- β -D-allopyranoside (28 mg) and cyclocerberidol-3-*O*- β -D-allopyranoside (14 mg).

The iridoids cerbinal, cerberic acid and cerberinic acid are found in the methanol extract of the bark of *C. manghas* [67]. The crude extract obtained by percolation from 4 kg of stem bark and 1.9 kg of root bark were diluted with water to 50% water:methanol. The mixture was washed with hexane and partitioned with benzene. Benzene fractions were re-suspended in methanol and cerbinal (120 mg from stem bark extract and 300 mg from root bark extract) precipitated.

The supernatant was chromatographed on a silica gel column and cerbinal, cerberic acid and cerberinic acid were eluted with benzene/acetone.

The iridoids isoplumericin, plumericin, plumieride, plumieride coumarate and plumieride coumarate glucoside can be detected and quantified in several species of *Plumeria* and *Allamanda* by TLC using silica gel 60 and the following mobile phases: benzene/ethyl acetate 4:1, chloroform/methanol 4:1, chloroform/methanol 7:3, propanol/ethyl acetate/water 7:2:1 [84]. Visualization of the chromatograms is achieved by spraying with 50% sulfuric acid/ethanol solution and heating. For the analyses, it is necessary to use iridoids as standards that can be isolated from the roots of *Allamanda cathartica*. For the isolation of iridoids during the development of the method, chloroform and methanol extracts, sequentially obtained in a Soxhlet apparatus from the root bark and inner roots of *Allamanda cathartica* (15.0 g for each extract) were successively fractionated on column chromatography. In the first chromatography step, the chloroform extract from the bark (1.7 g) was applied to a silica gel column (80 g) and eluted with gradient of petrol, ethyl ether, chloroform and methanol. This procedure yielded a mixture containing isoplumericin and plumericin (160 mg). Fractions eluted with chloroform/methanol (3:2) were mixed with the methanol extract of the bark (total mass: 2.8 g) and chromatographed on a silica gel column (150 g deactivated with water) with gradient of chloroform and methanol as mobile phase to furnish plumieride coumarate (150 mg), plumieride coumarate glucoside (480 mg) and two mixtures: one containing plumieride and plumieride coumarate, and the other containing plumieride and plumieride coumarate glucoside. The first mixture was further resolved on partition between water and ethyl acetate and afforded 180 mg of plumieride and 150 mg of plumieride coumarate. The other mixture was rechromatographed to afford plumieride (300 mg) and plumieride coumarate glucoside (410 mg). Neither the chloroform (810 mg) nor the methanol (920 mg) extracts from the inner part of the roots contained isoplumericin and plumericin. These extracts were purified on silica gel column (75 g) deactivated with water and eluted with chloroform and methanol gradient, to give plumieride (70 mg), plumieride coumarate (80 mg) and plumieride coumarate glucoside (200 mg).

The fresh leaves of *Cerbera manghas* contain the iridoids 10-*O*-benzoyltheveside, 10-dehydrogeniposide, loganin, theviridoside and theveside [124]. For the isolation, the methanol extract obtained by percolation using 2.6 kg of fresh leaves was extracted with butanol, and this extract was partitioned with benzene. After partition, the remaining butanol fraction (22.8 g) was subjected to column chromatography using MCI-gel (CHP-20) as stationary phase and a gradient of methanol/water as eluent. The fraction eluted with 20% methanol was subjected to C-18 column (elution with acetonitrile/water) to afford 20 mg of 10-dehydrogeniposide and 20 mg of loganin. The fraction eluted with pure water (2.6g) in the first chromatographic step was also subjected to C-18 column (elution with acetonitrile/ water) to furnish 23 mg of 10-*O*-benzoyltheveside and 270 mg of theveside.

According to [44] the iridoids fulvoplumerin, allamandin, α - and β -allamcidin, plumieride, 15-demethylplumieride, 13-*O*-*trans-p*-coumaroylplumieride and plumericin are present in extracts of *Plumeria rubra*. For isolation of these iridoids, the stem bark (2.5 kg) was successively extracted with petroleum ether and methanol followed by bioguided fractionation to investi-

gate the cytotoxic activity against various cancer cells, particularly murine lymphocytic leukemia cells (P-388). Petroleum ether extract (30 g) was submitted to flash column chromatography over silica gel (750 g, 230-400 mesh) with chloroform/petroleum ether (1:1) as eluent. Fraction 002 (200 mg) was rechromatographed using silica gel (120 g) and chloroform/methanol (99:1) to isolate fulvoplumierin (25 mg after recrystallization from petroleum ether/chloroform 1:1). The methanol extract (295 g) was partitioned between chloroform and water. The chloroform extract (60 g) was successively chromatographed in a silica gel column (1.5 kg) with gradients of increasing polarity, with chloroform and methanol, and then in a silica gel (600 g) with petroleum ether/chloroform/ethyl acetate (1:3:1) as eluent to furnish allamandin (12 mg after recrystallization from chloroform). Another fraction from the first column chromatography of the methanol extract (1.2 g) was purified in a silica gel column (400 g) and ethyl acetate/methanol (97:3) and recrystallized from chloroform to furnish 9 mg of allamcin. The aqueous extract (200 g) was also subjected to column chromatography using silica gel column (2 kg) and gradient of chloroform and methanol. Plumieride (55 g) was obtained directly from fraction F 022 after recrystallization from methanol. Fraction F022 (29 g) also furnished 15-demethylplumieride and 13-*O-trans-p*-coumaroylplumieride (3 g) by column chromatography over silica gel and gradient of chloroform and methanol. Fraction F018 (800 mg) was purified over silica gel (400 g) with gradient of ethyl acetate and methanol to afford plumericin (18 mg after recrystallization with ethyl acetate). Another fraction (F019) was purified over silica gel (250 g) with ethyl acetate/chloroform/methanol (6:6:1) solvent system to furnish an unstable iridoid aldehyde and a mixture of α - and β -allamcin (12 mg and 16 mg, respectively) further isolated by TLC on silica gel plates (20 cm x 20 cm, 250 μ m) with ethyl acetate/chloroform/methanol (3:3:1).

Plumericin, isoplumericin, plumieride, 13-*O*-coumaroylplumieride and protoplumericine A were isolated from the ethyl acetate extract of *P. rubra* heartwood [85]. The extract (44 g) obtained by percolation was submitted to column chromatography on silica gel with light petroleum ether/ethyl acetate (1:1) and methanol as eluent to furnish four fractions (A-D). Fraction C directly afforded plumericin (1.22 g) after recrystallization from toluene/ethyl acetate. Fraction B was rechromatographed on a silica gel with toluene/ethyl acetate (9:1), and isoplumericin (140 mg) was isolated from fraction 2. Fraction D was also rechromatographed on a silica gel column, but with chloroform/methanol/water (90:10:0.5 to 70:30:10) as solvent system. Fraction 3' was 13-*O*-coumaroylplumieride (2.0 g) and fraction 7' was protoplumericine (1.6 g). Fraction 5' was subjected to Sephadex LH20 column eluted with methanol, followed by silica gel column eluted with chloroform/methanol/water (85:15:0.7) to furnish plumieride (120 mg).

The iridoids, plumericin, isoplumericin, plumieride and fulvoplumierin, were present in the extracts of *Plumeria rubra* bark. After maceration of the powdered bark (3.5 kg) with dichloromethane/methanol (1:1) and pure methanol, the combined extracts were partitioned between water and ethyl acetate. To isolate the four iridoids, the organic layer was chromatographed twice in a column using silica gel and gradient of increasing polarity with hexane and ethyl acetate, ethyl acetate and methanol, and then pure methanol. The amounts of the compounds isolated were not reported [75].

The flowers of *Plumeria rubra* L. cv. *acutifolia* can provide plumericidine, as described by [97]. The ethanol (95%) extract, obtained from 2.9 kg of flowers, was successively partitioned with petroleum ether, ethyl acetate and butanol. The ethyl acetate fraction was sequentially submitted twice to column chromatography using silica gel and gradient of chloroform/methanol as mobile phase. Chromatography on a Sephadex LH20 column yielded 20 mg of plumericidine.

According to [72], several iridoids can be isolated from the aerial parts of *Plumeria obtusa*: obtusadoid A, obtusadoid B, plumieridin A, 1 α -plumieride, 15-demethylplumieride and plumieridine. The methanol extract (400 g) was obtained from 10 kg of the plant material and sequentially partitioned with hexane and ethyl acetate. The ethyl acetate extract was chromatographed using a silica gel column and gradients of hexane, ethyl acetate and methanol. The less polar fractions were rechromatographed in the same stationary phase, and eluted with hexane/dichloromethane (1:1) to afford obtusadoid A (6 mg), obtusadoid B (11.5 mg), plumieridin A (8 mg) and plumieridine (12 mg). The more polar fraction obtained in the first chromatography was filtered on a Sephadex LH20 column using methanol, and further submitted to RP-8 flash column chromatography. Elution with 50% methanol afforded 1 α -plumieride (22 mg) and 15-demethylplumieride (13 mg).

Plumieride also can be isolated from the bark of *Plumeria bicolor* [114]. Powdered bark (4 kg) was extracted in methanol, and the crude extract was washed with acetonitrile. The material was re-extracted with chloroform, and this extract was fractionated in column chromatography using silica gel (900 g) and different solvents of increasing polarity. Plumieride was eluted with chloroform/ethyl acetate (1:1) and recrystallized from methanol.

The bark of *Plumeria bicolor* also contains plumericin and isoplumericin, as described in [86]. The methanol extract (100 g), after washing with acetonitrile, was extracted with chloroform and chromatographed on a column containing 800 g of silica gel G (60-120 mesh). Elution was carried out using gradients of increasing polarities with benzene, chloroform and methanol. Plumericin and isoplumericin was recrystallized from methanol.

Isoplumericin and plumericin are present in the bark of *Himatanthus sucuuba* [88]. For the isolation of these iridoids, 95% ethanol extract (2 g), obtained from 50 g of plant material was submitted to column chromatography using silica gel and gradients of increasing polarities with hexane, ethyl acetate and methanol. After recrystallization, isoplumericin (18 mg) was obtained from ethyl acetate and plumericin (70 mg) from methanol.

The stem bark of *Winchia calophylla* contains loganin (1.25 g) [129]. The 95% ethanol extract (600 g) from the dried stem bark (10.5 kg) was partitioned between petroleum ether and water. The petroleum ether extract was submitted to acid-base extraction and after adjustment to pH 9-10 with ammonium hydroxide; the aqueous layer was extracted with petroleum ether, chloroform and butanol. The chromatography of the butanol fraction using silica gel H column led to the isolation of loganin.

The iridoids, scholarein A, B, C and D, can be obtained by the fractionation of the ethanol extract from bark of *Alstonia scholaris* [60]. The crude extract, obtained from 15 kg of the plant material, was partitioned between ethyl acetate and water. The organic layer (190 g) was

sequentially chromatographed on a column. The first chromatography, over silica gel (2.1 kg) and using gradient of chloroform and ethyl ether, furnished five fractions (1-5). From fraction 2, 8 mg of scholarein B and 7 mg of scholarein D were isolated after silica gel chromatography and elution with petroleum ether/ethyl ether (3:1). From fraction 3, 25 mg of scholarein A and 60 mg of scholarein C were obtained after successive columns using silica gel and chloroform/ethyl ether as stationary and mobile phases, respectively.

4.3. Gas Chromatography (GC)

In the study on iridoids, the technique of gas chromatography is generally used for analytical purposes. Gas chromatography represents an advantage over thin layer chromatography, particularly for detecting substances in small amounts, and mass spectrometry can be used to distinguish most iridoid and secoiridoid glucosides by fragmentation patterns [112].

Methods have been developed by [112] for the detection of 33 iridoids and secoiridoid glucosides in mixtures and plant extracts using gas chromatography sometimes coupled to mass spectrometry. For the gas chromatography analyses, a Shimadzu Model GC-1C gas chromatograph with hydrogen FID-1B flame ionization detector was used. Columns were packed with 1.5% OV-1, 1.5% OV-17, 2% OV-210 and 2% OV-225 on 80-100-mesh Shimalite W AW/DMCS. In all, 33 iridoids and secoiridoids glucosides are analyzed as TMS-derivatives. Using the 1.5% OV-17 column with 1.8 m in length and 4 mm in I.D. at 270 °C, the elution order was: aucubin, 7-deoxyloganic acid and catalpol (retention time = 1.37 min), 7-deoxyloganin, monotropein, gardenoside, secologanin, loganin, scandoside, theviridoside, geniposide, scandoside methyl ester, 7-dehydrologanin, morroniside, hastatoside and forsythide (retention time = 2.55 min), forsythide 10-methyl ester, verbenalin, sweroside, gentiopicroside and swertiamarin (retention time = 3.08 min), bankakosin, kingside, amaroswerin, amarogentin and asperuloside. The 1.5% OV-17 column, 0.5 m in length and 3 mm in I.D., at 230°C, furnished the same elution order as above, but forsythide 10-methyl ester, verbenalin and sweroside eluted together (retention time = 3.20 min), while the separation of hastatoside and forsythide, gentiopicroside and swertiamarin were better. When the non-polar 1.5% OV-1 column with 1.8 m in length and 4 mm in I.D. was used at 270 °C, better separation between loganin and secologanin occurred. Better results were achieved for the 7-deoxyloganic, 7-deoxyloganic acid and catalpol. However, verbenalin and sweroside eluted together, and amaroswerin and amarogentin were not detected. The OV-17 column was slightly polar and, in general, it influenced the larger range of the retention times. The more polar columns with 2% OV-210 at 215 °C, and 2% OV-225 at 230 °C, both with 0.5 m in length, showed important differences from keto compounds, such as 7-dehydrologanin and verbenalin, and lactonic compounds, such as sweroside, gentiopicroside and gentiopicroside, reflected in their longer retention times. Amaroswerin and amarogentin were not detected. Sweroside and gentiopicroside were well-separated on OV-210 column, which was not observed using other columns. Paederoside, ligustroside, catalposide, oleuropein, 10-acetoxyligustroside and 10-acetoxyleuropein were only detected and well-separated in OV-17 and OV-1 columns with 0.5m in length at 270 °C.

For the gas chromatography-mass spectrometry studies, a Hitachi K-53 gas chromatograph and a Hitachi RMU-6 E mass spectrometer were used. The glass columns, 0.5 m x 3 mm in I.D.,

were packed with 1.5% OV-17 on 80-100-mesh Shimalite W AW/DMCS and 1.5 % OV-1 on 80-100-mesh Shimalite W AW/DMCS, and were used to the oleuropein-type glucosides detection. The authors considered the GC-MS identification of some iridoids was not satisfactory: any important peak different from the sugar moiety was detected in asperuloside and paederoside TMS-derivatives of amaroswerin, amarogaentin asperuloside and paederoside exhibited different retention times, but the same fragmentation pattern; separation using this technique was unsuccessful [112].

Aqueous extracts of different plant species with known presence of iridoid and secoridoid glucosides were analyzed by GC-FID and GC-MS [112]. One of these species was *Allamanda cathartica* var. *schottii* (Pohl) Rafill (Apocynaceae) cultivated in a greenhouse. Aqueous extracts obtained from 3-5 g of fresh plant material and hot water were treated in a column of charcoal (active carbon for column chromatography) for removal of sugars by elution with water. The sample was eluted with methanol and concentrated under reduced pressure. TMS-derivatives were prepared. For the analyses, a 1.5% OV-17 column with 1.8 m in length at 280 °C was used. GC-FID and GC-MS (70 eV) showed the presence of one iridoid glycoside. The fragmentation pattern indicated that the original glucoside was plumieride, and that relative retention time was the same as that of asperuloside.

Isoplumericin, plumericin, plumieride, plumieride coumarate and plumieride coumarate glucoside can be detected by GC-FID [116]. For the development of the method, it was necessary to isolate these iridoids for use as standard. The methanol extract of *Allamanda cathartica* L. roots, obtained with 500 g of the dried plant material and boiling methanol, was submitted to silica gel 60 column (2.5 kg) with the eluents: petrol, petrol/ethyl ether, ethyl ether, ethyl ether/chloroform, chloroform/ methanol and pure methanol [116]. Fractionation was monitored by TLC [84]. The fractions were eluted with petrol/ethyl ether until chloroform/methanol contained isoplumericin and plumericin (4.2 g). This mixture (1.0 g) was suspended in ethyl ether and rechromatographed on silica gel (40 g). Elution with petrol and a gradient of increasing polarity with petrol/ethyl ether furnished isoplumericin (250 mg), plumericin (140 mg) and a mixture of both (290 mg). A second fraction eluted with chloroform/methanol in the first chromatographic step (8.0 g) was resubmitted to column chromatography on silica gel (300 g). Elution with a gradient of chloroform and methanol led to the isolation of plumieride coumarate (5.1 g). The third fraction of the first chromatograph step (10.0 g), eluted with chloroform/methanol, was partitioned between water and ethyl acetate to give plumieride coumarate in the organic phase (3.8 g) and plumieride in the aqueous phase (1.8 g). Finally, 30.0 g of the forth fraction of the first chromatograph step, eluted with chloroform/methanol to methanol, was submitted to column chromatography on silica gel (1.0 kg) deactivated with water. The elution was carried out with a gradient of chloroform and methanol and furnished plumieride coumarate glucoside (6.2 g). GC-FID analyses were used to evaluate the pure grade of fractions and isolated substances. For these analyses, trimethylsilylation of iridoids using HMDS-TMCS and pyridine, and acetylations with acetic anidride and pyridine, were necessary. Plumieride coumarate (isomer mixture) and plumieride coumarate glucoside were acetylated and their products were purified on chromatography with silica gel column and ethyl ether as eluents, to afford pure penta-acetylplumieride coumarate and octa-acetylplu-

mieride coumarate glucoside, respectively. Furthermore, plumieride, plumieride coumarate and plumieride glucoside were hydrolyzed under heating and acid conditions (sulfuric acid, 1 N for 2-3 h), extracted with ethyl acetate and both organic phase and aqueous phase (after neutralization with Amberlite) were analyzed by TLC and GC. Analyses were performed on a 1.5% OV-17 glass column with 0.4 m length and 4 mm I.D. Other analytical conditions were: nitrogen as carrier gas at 50 mL/min; detector temperature, 320 °C; column temperatures: 190 °C for isoplumericin and plumericin, 240 °C for TMS-plumieride, 300 °C TMS-plumieride coumarate. Glucose was detected in aqueous phases from acid hydrolyses of plumieride, plumieride coumarate and plumieride coumarate glucoside, while *p*-coumaric acid was detected in the organic phases of plumieride coumarate and plumieride coumarate glucoside. GC analyses also showed that plumieride coumarate was isolated as an isomer mixture (20% cis and 80% trans isomers) [116].

4.4. High, Medium and Low Performance Liquid Chromatography (HPLC, MPLC and LPLC)

Studies on HPLC with iridoids of Apocynaceae focus mainly on the separation of components from extracts or fraction. The chromatography profile, the identification and quantification of these terpenes in the extracts are described. Table 4 shows the principal references on iridoids isolated from Apocynaceae by HPLC, MPLC and LPLC.

TECHNIQUE	MOBILE PHASE	COLUMN	SPECIES/ PLANT MATERIAL	SAMPLE	IRIDOID	REF
HPLC	50% methanol in water	μBondapack C-18	<i>Rauwolfia grandiflora</i> / bark	ethanol extract	boonein and isoboonein	[1]
HPLC	10% acetonitrile	Octadecylsilane (ODS)	<i>Thevetia peruviana</i> / root	methanol extract	theviridoside, 10-O-β-D-fructofuranosyltheviridoside, 6'-O-β-D-glucopyranosyltheviridoside, 10-O-β-D-glucopyranosyltheviridoside and 3'-O-β-D-glucopyranosyltheviridoside	[125]
HPLC	10% acetonitrile	Octadecylsilane (ODS)	<i>Thevetia peruviana</i> / leaves	methanol extract	10-O-β-D-fructofuranosyltheviridoside and 6'-O-β-D-glucopyranosyltheviridoside	[127]
HPLC	20 mM KH ₂ PO ₄ and acetonitrile Flow: 0.8 mL/min	Cosmosil 5 C ₁₈ -AR (5 μm, 25 cm x 4.6 mm I.D.)	<i>Rauwolfia</i> / barks	ethanol extract	loganin, loganic acid and gardenoside	[131]

TECHNIQUE	MOBILE PHASE	COLUMN	SPECIES/ PLANT MATERIAL	SAMPLE	IRIDOID	REF
HPLC	water/ acetonitrile	Octadecyl-silane (ODS) (250×10mm)	<i>Plumeria acutifolia</i> / leaves	methanol extract	15-demethylplumieride	[71]
HPLC	not informed	Zorbax SB-C18	<i>Plumeria rubra</i> L. cv. <i>acutifolia</i> / flowers	ethyl acetate fraction	plumieridin A and plumieridin B	[120]
HPLC	not informed	not informed	<i>Allamanda neriifolia</i> / aerial parts	ethanol extract	allamanoid, plumieride and protoplumericin	[52]
LPLC	5% ethyl acetate in chloroform	Michel-Miller column (200 g Silica gel)	<i>Allamanda schottii</i> / stem	ethanol extract	allamandin and allamcin	[47]
MPLC	water to 50% methanol in water	RP-18 (25–40 mm particle size, 460 × 36 mm I.D., 460 × 15 mm I.D.)	<i>Himatanthus sucuuba</i> / latex	methanol/ water (1:1)	plumieride	[73]
TECHNIQUE	MOBILE PHASE	COLUMN	SPECIES/ PLANT MATERIAL	SAMPLE	IRIDOID	REF.
HPLC	Acetonitrile and water containing 0.05% trifluoro- acetic acid Flow: 1 mL/min	Lichrospher C18 (5mm, 250 mm × 4.6 mm I.D.)	<i>Himatanthus sucuuba</i> / bark and latex	methanol/ water (1:1)	plumieride, isoplumieride and demethylplumieride	[73]
MPLC and HPLC	water to 50% methanol; 30% acetonitrile in water to 50% acetonitrile MPLC Flow: 6 mL/min	LiChroprep C-18 (45 × 3.5 cm) and Shimpack C-18 (10 μm, 45 cm × 250 mm I.D.)	<i>Himatanthus sucuuba</i> / latex	aqueous fraction and fractions from this procedure	15-demethylisoplumieride, 15- demethylplumieride, plumieride and isoplumieride	[70]

Table 4. Separation by Pressure Liquid Chromatography

4.5. Counterflow

Protoplumericin and plumieride can be extracted from the methanol extract of *Allamanda neriifolia* stems [117] by droplet counter-current chromatography. Crude extract, obtained from 1.6 kg of plant material, was successively partitioned with benzene, chloro-

form and butanol. Plumericin (420 mg) was directly obtained from the benzene fraction (0.02% yield). The butanol fraction was subjected to sequential chromatographic steps, using XAD-2 column and gradient of water/methanol (mobile phase), silica gel column and solvent system containing chloroform/methanol/water or chloroform/methanol, followed by droplet current chromatography with chloroform/methanol/ water. Plumieride (1.3 g) and protoplumericin (13.2 g) were obtained with yields of 0.08% and 0.83%, respectively.

The iridoids allamcidin B β -D-glucoside, plumiepoixide and protoplumericin B were isolated from *Allamanda nerifolia* extract obtained by percolation with methanol, also by droplet counter-current chromatography [39]. The crude methanol extracts from 2.6 kg of stem and 6.7 kg of leaves were sequentially fractionated with benzene, chloroform and butanol. Previous chromatographic treatment with MCI gel (elution with water/methanol), silica gel column (mobile phases: chloroform/methanol/water; benzene/acetone; ethyl acetate/methanol/ water and ethyl acetate/hexane) and Sephadex LH20 column (mobile phase: chloroform/methanol) led to the isolation of isoallamandicin (10 mg from the stem), allamcin (230 mg from the leaves), 3-O-methylallamcin (30 mg from the leaves), allamancin (102 mg from the stem), 3-O-methylallamancin (41 mg from the leaves), allamcidin (125 mg from the leaves), plumieride 13-O-acetate (760 mg from the stem and ca. 2 g from the leaves). Fractions of butanol extracts from the stem and leaves were subjected to droplet counter-current chromatography using chloroform/methanol/water (5:6:4, ascending mode) to obtain allamcidin B β -D-glucoside (17 mg from the stem), plumiepoixide (7 mg from the stem and 374 mg from the leaves) and protoplumericin B (70 mg from the leaves).

Iridoids can be isolated from *Plumeria acutifolia* roots following a similar methodology [66]. Successive liquid-liquid partitions of the crude methanol extract (obtained from 6 kg of plant material) with benzene, chloroform and butanol, followed by several chromatographic steps for fractionation of the butanol fraction (using polystyrene, silica gel and octadecyl silica columns) led to the isolation of 13-O-coumaroylplumieride (43 g), plumieride (7.5 g), 13-O-caffeoylplumieride (60 mg), 1 α -plumieride (20 mg) and protoplumericin A (9 g). A further purification step involving droplet counter-current chromatography and a mixture of chloroform/methanol/water (4:6:5, ascending mode) led to the isolation of 13-deoxyplumieride (200mg), plumenoside (50 mg) and 8-isoplumieride (700 mg).

4.6. Capillary electrophoresis

For analytical purposes, iridoids can be analyzed by capillary electrophoresis. A method to separate nine iridoids described in [131] uses a Hewlett-Packard (HP^{3D} CE) capillary electrophoresis system coupled to a photodiode array detector (210 nm and 230 nm) and equipped with a fused-silica capillary tube (90 cm \times 75 μ m I.D.). The distance to the detector was 81.5 cm. Other conditions: sample injection at 50 mbar for 3 s and further deionized water injection at 50 mbar for 3 s; constant voltage, 16 kV (positive to negative); cartridge temperature, 30 $^{\circ}$ C; electrolyte (buffer), 50 mM sodium borate and 30 mg/mL 2,6-di-O-methyl- β -cyclodextrin (DM- β -CD); run time, 32 min. Before the analyses, the capillary column was sequentially purged with 0.5 M NaOH, 0.1 M NaOH, deionized water and buffer solution. The iridoids studied eluted in the following order: geniposide, loganin, shanzhiside, aucubin, catalpol, harpago-

side, gardenoside, geniposidic acid and loganic acid. All were commercially purchased and only loganin, loganic acid and gardenoside were described for the Apocynaceae family. Several conditions of analyses were studied, including different pH, surfactants, concentrations of sodium borate and the addition of cyclodextrins (CD) to the buffer, and it was concluded that the less polar DM- β -CD added to 50 mM borate solution was the most suitable running buffer. In this condition, only aucubin and catalpol could not be separated, even with the addition of organic solvents and/or valine, urea and barium ion. The greatest advantage of capillary electrophoresis compared to HPLC analyses (the most commonly used technique) is its speed.

According to [132], capillary electrophoresis can be used to analyze a mixture of eleven iridoid glycosides: unedoside, harpagide, methyl catalpol, morroniside, asperuloside, griselinoside, catalpol, ketologanin, verbenalin, loganin and 10-cinnamoyl catalpol. Only loganin was found in the Apocynaceae family. For the analyses, iridoids were diluted in purified water. A Hewlett-Packard ^{3D}CE system coupled to a diode array detector and equipped with an air-cooling device was used. The fused-silica capillary tube measured 80 cm in length, 50 μ m in I.D. and 375 μ m in O.D. Distance to the detector was 71.5 cm only for UV detection (197 nm, 235 nm, 239 nm and 283 nm). When coupled to a mass spectrometer system (Bruker ESQUIRE) with an electrospray ionization source, the drying gas was nitrogen at 200 °C and flow-rate 100 L/h. In this case, the distance between injector and UV detector was 20 cm. Other conditions: sample injection at 50 mbar for 5 s (only UV detection) or 25 s (with MS system); voltage, +20 kV; cartridge temperature, 25 °C; electrolyte solution, 20 mM ammonium acetate with 100 mM sodium dodecyl sulfate (SDS), pH 9.5; sheath liquid, 1 mM lithium acetate mixture to water/methanol (1:1 v/v) at a flow rate of 200 μ L/h. When the MS system was used: scan range, 100-550 *m/z*; cut-off, 80 *m/z*; glass capillary exit, 95 V; skimmer, 32 V; electrospray voltage for the capillary, -4.0 kV; for the cylinder, -1,8 kV; for the end plate, -3.5 kV. In the comparison among the counterions sodium dodecyl sulfate (SDS), ammonium dodecyl sulfate and lithium dodecyl sulfate, diluted in water and running buffer, the best resolution for separating iridoid glucosides, lower noise in the MS system, and better repeatability and sensitivity were found with SDS in the running buffer. The volatility of ammonium acetate in buffer enables MS analyses, and concentrations higher than 20 mM did not represent better resolution. Quite the contrary, higher SDS concentrations furnished better results. In the study of the influence of pH, the best one was 9.5, although its influence in the range of 8.7-10.0 was lower than the SDS effect. Good linearity was observed for all the iridoids glucosides analyzed, but in different ranges.

The literature describes chromatographic techniques related to the characterization, isolation and purification of iridoids. Most reports show the open column technique as the principal technique used to isolate this class. Also, there have been few studies on counterflow and capillary electrophoresis chromatographies. In general, there has been little scientific investment in the area of obtaining iridoids of the Apocynaceae family, despite the great pharmacological importance of this class of constituents.

Author details

Ana Cláudia F. Amaral¹, Aline de S. Ramos¹, José Luiz P. Ferreira^{1,2}, Arith R. dos Santos¹, Deborah Q. Falcão², Bianca O. da Silva⁴, Debora T. Ohana^{1,5} and Jefferson Rocha de A. Silva³

*Address all correspondence to: acamaral@fiocruz.br

1 Laboratório de Plantas Medicinais e Derivados, Depto de Produtos Naturais, Farmanguinhos – FIOCRUZ, Mangunhos, Brazil

2 Faculdade de Farmácia – UFF, Niterói, Brazil

3 Laboratório de Cromatografia – Depto. de Química – UFAM, Japiim, Manaus, Brazil

4 Instituto de Pesquisas Biomédicas, Hospital Naval Marcílio Dias, Brazil

5 Fac. de Ciências Farmacêuticas – UFAM, Manaus, Brazil

References

- [1] Bianco A, De Luca A, Mazzei RA, Nicoleit M, Passacantilli P, Limas RA. Iridoids of *Rauwolfia grandiflora*. *Phytochemistry* 1994; 35(6) 1485-1487.
- [2] Dewick PM. *Medical Natural Products: A biosynthetic Approach*. 2^o Edition. John Wiley & Sons, LTD; 2002.
- [3] Sampaio-Santos MI, Kaplan MAC. Biosynthesis Significance of Iridoids in Chemo-systematics. *Journal of the Brazilian Chemical Society* 2001; 12(2) 144-153.
- [4] Oudin A, Courtois M, Rideau M, Clastre M. The Iridoid Pathway in *Catharanthus roseus* Alkaloid Biosynthesis. *Phytochemical Reviews* 2007; 6 259–276.
- [5] Eisenreich W, Bacher A, Arigoni D, Rohdich F. Biosynthesis of Isoprenoids Via the Non-Mevalonate Pathway. *Cellular and Molecular Life Sciences*. 2004; 61 1401–1426.
- [6] Hunter WN. The Non-Mevalonate Pathway of Isoprenoid Precursor Biosynthesis. *Journal of Biological Chemistry* 2007; 282 21573–21577.
- [7] Contin A, Van der Heijden R, Lefeber AWM, Verpoorte R. The Iridoid Glucoside Secologanin is Derived from the Novel Triose Phosphate/Pyruvate Pathway in a *Catharanthus roseus* Cell Culture. *FEBS Letters* 1998; 434 413–416.
- [8] Eichinger D, Bacher A, Zenk MH, Eisenreich, W. Analysis of metabolic pathways via quantitative prediction of isotope labeling patterns: a retrobiosynthetic ¹³C NMR study on the monoterpene loganin. *Phytochemistry* 1999; 51 223–236.

- [9] Li H, Yang S, Wang H, Tian J, Gao W. Biosynthesis of the Iridoid Glucoside, Lamalbid, in *Lamium barbatum*. *Phytochemistry* 2010; 7 1690–1694.
- [10] Inouye H, Uesato S. Biosynthesis of Iridoids and Secoiridoids. In: *Progress in the Chemistry of Organic Natural Products*, Vol. 50 (Herz W., Grisebach H., Kirby G. W., and Tamm Ch., eds.). Springer, New York, 1986 169–236.
- [11] Jensen SR, Franzyk H, Wallander E. Review: Chemotaxonomy of the Oleaceae: Iridoids as Taxonomic Markers. *Phytochemistry* 2002; 60 213–231.
- [12] Nagatoshi M, Terasaka K, Nagatsu A, Mizukami H. An Iridoid-Specific Glucosyltransferase from *Gardenia jasminoides*. *JBC* July, 2011: 1–19. Available from <http://www.jbc.org/cgi/doi/10.1074/jbc.M111.242586> (accessed 20 September 2012).
- [13] Galvez M, Martin-Cordero C, Ayuso MJ. Pharmacological Activities of Iridoids Biosynthesized by Route II. *Studies in Natural Products Chemistry* 2005; 32 365–394.
- [14] Cimanga K, Kambu K, Tona L, Hermans N, Apers S, Totte J, Pieters L, Vlietinck AJ. Antiamoebic Activity of Iridoids from *Morinda morindoides* Leaves. *Planta Medica*. 2006; 72 751–753.
- [15] Yang XP, Li EW, Zhang Q, Yuan CS, Jia ZJ. Five New Iridoids from *Patrinia rupestris* *Chemistry & Biodiversity*. 2006; 3 762–770.
- [16] Tzakou O, Mylonas P, Vagias C, Petrakis PV. Iridoid Glucosides with Insecticidal Activity from *Galium melanantherum*. *Zeitschrift für Naturforschung, C: Journal of Biosciences* 2007; 62 597–602.
- [17] da Silva VC, Giannini MJSM, Carbone V, Piacente S, Pizza C, Bolzani VS, Lopes MN. New Antifungal Terpenoid Glycosides from *Alibertia edulis* (Rubiaceae). *Helvetica Chimica Acta* 2008; 91 1355–1362.
- [18] Tundis R, Loizzo MR, Menichini F, Statti GA, Menichini F. Biological and Pharmacological Activities of Iridoids: Recent Developments. *Mini-Reviews in Medicinal Chemistry* 2008; 8 399–420.
- [19] Castillo L, Rossini C. Bignoniaceae Metabolites as Semiochemicals. *Molecules* 2010; 15 7090–7105.
- [20] Zhang HJ, Rothwangl K, Mesecar AD, Sabahi A, Rong LJ, Fong HHS. Lamiridosins, Hepatitis C Virus Entry Inhibitors from *Lamium album*. *Journal of Natural Products* 2009; 72 2158–2162.
- [21] Bai NS., HK, Ibarra A, Bily A, Roller M, Chen XZ, Rühl R. Iridoids from *Fraxinus excelsior* with Adipocyte Differentiation-Inhibitory and PPAR α Activation Activity. *Journal of Natural Products* 2010 (73) 2–6.
- [22] Dahlgren G. An updated angiosperm classification. *Botanical Journal of the Linnean Society*. 1989; 100: 197–203. doi: 10.1111/j.1095-8339.1989.tb01717.x (accessed 18 september 2012).

- [23] Hegnauer, R. *Chemotaxonomie der Pflanzen*. Birkhäuser Verlag, Basel. 1986.
- [24] Taskova R, Peev D, Handjieva N. Iridoid Glucosides of the Genus *Veronica* s.l. and their Systematic Significance. *Plant Systematics and Evolution* 2002; 231 1-17.
- [25] Rivière C, Goossens L, Guerardel Y, Maes E, Garénaux E, Pommery J, Pommery N, Désiré O, Lemoine A, Telliez A, Delelis A, Hénichart J.P. Chemotaxonomic interest of iridoids isolated from a Malagasy species: *Perichlaena richardii*. *Biochemical Systematics and Ecology* 2011; 39: 797–825.
- [26] Wink M, Waterman P. Chemotaxonomy in relation to molecular phylogeny of plants. In: Wink, M. (ed.), *Biochemistry of plant secondary metabolism*, Annual Plant Reviews. Boca Raton: Sheffield Academic Press and CRC Press. 1999: 300–341.
- [27] Wink, M. Evolution of Secondary Metabolites from an Ecological and Molecular Phylogenetic Perspective. *Phytochemistry* 2003; 64 3–19.
- [28] Dahlgren RMT. A revised System of Classification of Angiosperms. *Botanical Journal of the Linnean Society*. 1980; 80 91-124.
- [29] Angiosperm Phylogeny Group III. An Update of the Angiosperm Phylogeny Group Classification for the Orders and Families of Flowering Plants: APG III. *Botanical Journal of the Linnean Society* 2009; 161 105–121.
- [30] Taskova, RM, Gotfredsen CH, Jensen SR. Chemotaxonomy of Veroniceae and its Allies Plantaginaceae. *Phytochemistry* 2006; 67 286–301.
- [31] Koch, I, Rapini A. Apocynaceae. In: *Lista de espécies da flora do Brasil*. Jardim Botânico do Rio de Janeiro, Rio de Janeiro. 2011. <http://floradobrasil.jbrj.gov.br/2011/FB00004>. (accessed 10 September 2012).
- [32] Endress ME, Liede-Schumann S, Meve U. Advances in Apocynaceae: the Enlightenment, an Introduction. *Annals of the Missouri Botanical Garden* 2007; 94 259–267.
- [33] Rapini A. Taxonomy “Under Construction”: Advances in the Systematic of Apocynaceae, with Emphasis on the Brazilian Asclepiadoideae. *Rodriguésia* 2012; 63(1): 75-88.
- [34] Meve U. Species Numbers and Progress in Asclepiad Taxonomy. *Kew Bulletin* 2002; 57 459–464.
- [35] The International Plant Names Index. <http://www.inpi.org/> (accessed 04 October 2012).
- [36] Tropicos. org. Missouri Botanical Garden. <http://www.tropicos.org> (accessed 04 October 2012).
- [37] Siddiqui BS, Naeed A, Begum S, Siddiqui S. Minor Iridoids from the Leaves of *Plumeria obtusa*. *Phytochemistry* 1994; 37(3) 769-771.
- [38] Begun S, Naeed A, Siddiqui BS, Siddiqui S. Chemical Constituents of the Genus *Plumeria*. *Journal of the Chemical Society of Pakistan* 1994; 16(4) 280-299.

- [39] Abe F, Mōri T, Yamauchi T. Iridoids of Apocinaceae. III. Minor Iridoids from *Allamanda neriifolia*. Chemical Pharmaceutical Bulletin 1984; 32(8) 2947-2956.
- [40] Boros CA, Stermitz FR. Iridoids. An Update Review, Part II. Journal of Natural Products 1991; 54(5) 1173-1246.
- [41] Kupchan SM, Dessertine AL, Blaylock BT, Bryan RF. Isolation and Structural Elucidation of Allamandin, an Antileukemic Iridoid Lactone from *Allamanda cathartica*. Journal of Organic Chemistry 1974; 39(17) 2477-2482.
- [42] El-Naggar LJ, Beal JL. Iridoids. A Review. Journal of Natural Products 1980; 43(6) 649-707.
- [43] Sharma G, Chahar MK, Dobhal S, Sharma N, Sharma TC, Sharma MC, Joshi YC, Dobhal MP. Phytochemical Constituents, Traditional Uses and Pharmacological Properties of the Genus *Plumeria*. Chemical & Biodiversity 2011; 8(8) 1357-1369.
- [44] Kardono LBS, Tsauri S, Padmawinata K, Pezzuto JM, Kinghorn AD. Cytotoxic Constituents of the Bark of *Plumeria rubra* Collected in Indonesia. Journal of Natural Products 1990; 53(6) 1447-1455.
- [45] Vanderlei MF, Silva MS, Gottlieb HF, Braz-Filho R. Iridoids and Triterpenes from *Himatanthus phagedaenica*: the Complete Assignment of the Proton and Carbon-13 NMR Spectra of Two Iridoid Glycosides. Journal of the Brazilian Chemical Society 1991; 2(2) 51-55.
- [46] Al-Hazimi HMG, Alkathlan H. Naturally Occurring Iridoids During the Period 1990-1993. Journal of the Chemical Society of Pakistan 1996; 18(4) 336-357.
- [47] Anderson JE, Chang CJ, McLaughlin JL. Bioactive Components of *Allamanda schottii*. Journal of Natural Products 1988; 51(2) 307-308.
- [48] Kuigoua GM, Kouam SF, Ngadjui BT, Schulz B, Green IR, Choudhary MI, Krohn K. Minor Secondary Metabolic Products from the Stem Bark of *Plumeria rubra* Linn. Displaying Antimicrobial Activities. Planta Medica 2010; 76(6) 620-625.
- [49] Lopes RK, Ritter MR, Rates SMK. Revisão das Atividades Biológicas e Toxicidade das Plantas Ornamentais mais Utilizadas no Rio Grande do Sul, Brasil. Brazilian Journal of Biosciences 2009; 7(3) 305-315.
- [50] Morel AF, Graebner IB, Porto C, Dalcol II. Study on the Antimicrobial Activity of *Himatanthus sucuuba*. Fitoterapia 2006; 77(1) 50-53.
- [51] Amaral ACF, Ferreira JLP, Pinheiro MLB, Silva JR de A. Monograph of *Himatanthus sucuuba*, a Plant of Amazonian Folk Medicine. Pharmacognosy Reviews 2007; 1(2) 305-313.
- [52] Yu YL, Li X, Ke CA, Tang CP, Yang XZ, Li XQ, Ye Y. Iridoid Glucosides from *Allamanda neriifolia*. Chinese Chemical Letters 2010; 21(6) 709-711.

- [53] Dinda B, Debnath S, Banik R. Naturally Occurring Iridoids and Secoiridoids. An Update review, Part 4. *Chemical Pharmaceutical Bulletin* 2011; 59(7) 803-833.
- [54] Shen YC, Chem CH. Allaneroside and Bioactive Iridoid Glycosides from *Allamanda neriiifolia*. *Taiwan Yaoxue Zazhi* 1986; 38(4) 203-213.
- [55] Abe F, Yamauchi T, Wan ASC. Studies on *Cerbera*: VIII. Normonoterpenoids and their Allopyranosides from the Leaves of *Cerbera* Species. *Chemical & Pharmaceutical Bulletin* 1989; 37(10) 2639-2642.
- [56] Shen LR, Jin SM, Yin BW, Du XF, Wang YL, Huo CH. Chemical Constituents of Plants from the Genus *Cerbera*. *Chemistry & Biodiversity* 2007; 4(7) 1438-1449.
- [57] Thomas PS, Kanaujia A, Ghosh D, Duggar R, Katiyar CK. Alstonoside, a Secoiridoid Glucoside from *Alstonia scholaris*. *Indian Journal of Chemistry, section B: Organic Chemistry Including Medicinal Chemistry* 2008; 47B(8), 1298-1302.
- [58] Dinda B, Chowdhury DR, Mohanta BC. Naturally Occurring Iridoids, Secoiridoids and Their Bioactivity. An Update Review, Part 3. *Chemical Pharmaceutical Bulletin* 2009; 57(8) 765-796.
- [59] Topcu G, Che CT, Cordell GA, Ruangrunsi N. Iridolactones from *Alyxia reinwardtii*. *Phytochemistry* 1990; 29(10) 3197-3199.
- [60] Feng T, Cai XH, Du ZZ, Luo XD. Iridoids from the Bark of *Alstonia scholaris*. *Helvetica Chimica Acta* 2008; 91(12) 2247-2251.
- [61] Yamauchi T, Abe F, Wan ASC. 10-O-Benzoyltheveside and 10-Dehydrogeniposide from the Leaves of *Cerbera manghas*. *Phytochemistry* 1990; 29(7) 2327-2328.
- [62] Abe F, Yamauchi T. 10-Carboxyloganin, Normonoterpenoid Glucosides and Dinormonoterpenoid Glucosides from the Leaves of *Cerbera manghas* (Studies on *Cerbera*. X). *Chemical & Pharmaceutical Bulletin* 1996; 44(10) 1797-1800.
- [63] Marini-Bettolo GB, Nicoletti M, Messana I, Patamia M, Galeffi C, Oguakwa JU, Portalone G, Vaciago A. Research on African Medicinal Plants - IV. Boonein, a New C-9 Terpenoid Lactone from *Alstonia boonei*: A Possible Precursor in the Indole Alkaloid Biogenesis. *Tetrahedron* 1983; 39(2) 323-329.
- [64] Adotey JPK, Adukpo YDB, Armah FA. A Review of the Ethnobotany and Pharmacological Importance of *Alstonia boonei* De Wild (Apocynaceae). *ISRN Pharmacology* 2012; Article ID 587160, 9 pages. <http://www.hindawi.com/isrn/pharmacology/2012/587160/> (accessed 15 september 2012). doi: 10.5402/2012/587160.
- [65] Patil CD, Patil SV, Borase HP, Salunke BK, Salunkhe RB. Larvicidal Activity of Silver Nanoparticles synthesized Using *Plumeria rubra* Plant Latex Against *Aedes aegypti* and *Anopheles stephensi*. *Parasitological Research* 2012; 110(5) 1815-1822.

- [66] Abe F, Chen RF, Yamauchi T. Studies on the Constituents of *Plumeria*. Part I. Minor Iridoids from the Roots of *Plumeria acutifolia*. Chemical & Pharmaceutical Bulletin 1988; 36(8) 2784-2789.
- [67] Abe F, Okabe H, Yamauchi T. Studies on *Cerbera*: II. Cerbinal and its Derivatives, Yellow Pigments in the Bark of *Cerbera manghas* L.. Chemical & Pharmaceutical Bulletin 1977; 25(12) 3422-3424.
- [68] Zhang X, Zhang J, Liu M, Pei Y. Chemical Components in Cortex of *Cerbera manghas* L. Zhongcaoyao 2008; 39(8) 1138-1140.
- [69] Ali FI, Hashmi IL, Siddiqui BS. A novel iridoid from *Plumeria obtusa*. Natural Product Communications 2008; 3(2) 125-128.
- [70] Barreto A de S, Amaral ACF, Silva JR de A, Schripsema J, Rezende CM, Pinto AC. Ácido 15-desmetilisoplumierideo, um Novo Iridóide Isolado das Cascas de *Plumeria rubra* e do Látex de *Himatanthus sucuuba*. Química Nova 2007; 30(5) 1133-1135.
- [71] Hassan EM, Shahat AA, Ibrahim NA, Vlietinck A, Apers S, Pieters L. A New Monoterpene Alkaloid and other Constituents of *Plumeria acutifolia*. Planta Medica 2008; 74(14) 1749-1750.
- [72] Saleem M, Akhtar N, Riaz N, Ali MS, Jabbar A. Isolation and Characterization of Secondary Metabolites from *Plumeria obtusa*. Journal of Asian Natural Products Research 2011; 13(12) 1122-1127.
- [73] Silva JR de A, Amaral ACF, Silveira CV, Rezende CM, Pinto AC. Quantitative Determination by HPLC of Iridoids in the Bark and Latex of *Himatanthus sucuuba*. Acta Amazonica 2007; 37(1) 119-122.
- [74] Albers-Schoenberg G, Schmid H. Über Die Struktur von Plumericin, Isoplumericin, β -Dihydroplumericin Acid. Helvetica Chimica Acta 1961; 44(181) 1447-1473.
- [75] Elsässer B, Krohn K, Akhtar MN, Florke U, Kouam SF, Kuigoua MG, Ngadjui BT, Abegaz BM, Antus S, Kurtán T. Revision of the Absolute Configuration of Plumericin and Isoplumericin from *Plumeria rubra*. Chemistry & Biodiversity 2005; 2(6) 799-808.
- [76] Vankata Rao E, Anjaneyulu TSR. Chemical Components of the Bark of *Plumeria rubra*. Indian Journal of Pharmacy 1967; 29(9) 273-274.
- [77] Tandon SP, Tiwarik P, Rathore YKS. Chemical Constituents of the Decorticated Stem of *Plumeria rubra* Linn. Proceedings of the National Academy of Sciences, India Section A 1976; 46(2) 109-110.
- [78] Grumbach A, Schmid H, Bencze W. An Antibiotic from *Plumeria acutifolia*. Experientia 1952; 8 224-225.
- [79] Schmid H, Bencze W. Über die Konstitution des Fulvoplumierins I. Helvetica Chimica Acta 1953; 36(27) 205-214.

- [80] Rangaswami S, Venkata Rao E, Suryanarayana M. Chemical Examination of *Plumeria acutifolia*. Indian Journal of Pharmacy 1961; 23 122-124.
- [81] Wanner H, Zorn-Ahrens V. Distribution of Plumieride in *Plumeria acutifolia* and *Plumeria bracteata*. Berichte de Schweizerischen Botanischen Gesellschaft 1972; 81 27-39.
- [82] Mahran GH, Abdel-Wahab, SM, Ahmed MS Detection and Isolation of the Antibiotic Principles of *Plumeria rubra* L. and *Plumeria rubra* L. var. *alba*. Growing in Egypt. Bulletin of the Faculty of Pharmacy (Cairo University) 1975; 12(2) 151-60.
- [83] Albers-Schönberg, G, Philipsborn WV, Jackman LM, Schmid H. Die Struktur de Fulvoplumierins. Helvetica Chimica Acta 1962; XLV(164) 1406-1408.
- [84] Coppen JJW, Cobb AL. The Occurrence of Iridoids in *Plumeria* and *Allamanda*. Phytochemistry 1983; 22(1) 125-128.
- [85] Hamburger MO, Cordell GA, Ruangrunsi N. Traditional Medicinal Plants of Thailand XVII. Biologically Active Constituents of *Plumeria rubra*. Journal of Ethnopharmacology 1991; 33(3) 289-292.
- [86] Sharma U, Singh D, Kumar P, Dobhal MP, Singh S. Antiparasitic Activity of Plumericin & Isoplumericin Isolated from *Plumeria bicolor* against *Leishmania donovani*. Indian Journal of Medical Research 2011; 134(5) 709-716.
- [87] Barreto A de S, Carvalho MG, Neri I, Gonzaga L, Kaplan MAC. Chemical Constituents from *Himatanthus articulata*. Journal of the Brazilian Chemical Society 1998; 9(5) 430-434.
- [88] Castillo D, Arevalo J, Herrera F, Ruiz C, Rojas R, Rengifo E, Vaisberg A, Lock D, Lemesre JL, Gornitzka H, Sauvain M. Spirolactone Iridoids might be responsible for the Antileishmanial Activity of a Peruvian Traditional Remedy Made with *Himatanthus sucuuba* (Apocynaceae). Journal of Ethnopharmacology 2007; 112(2) 410-414.
- [89] Abdel-Kader MS, Wisse J, Evans R, van der Werff H, Kingston DGI. Bioactive Iridoids and a New Lignan from *Allamanda cathartica* and *Himatanthus fallax* from the Suriname Rainforest. Journal of Natural Products 1997; 60(12) 1294-1297.
- [90] Silva JR de A, Rezende CM, Pinto AC, Pinheiro MLB, Cordeiro MC, Tamborini E, Young CM, Bolzani V da S. Ésteres Triterpênicos de *Himatanthus sucuuba* (Spruce) Woodson. Química Nova 1998; 21(6) 702-704.
- [91] Miranda ALP, Silva JR de A, Rezende CM, Neves JS, Parrini SC, Pinheiro MLB, Cordeiro MC, Tamborini E, Pinto AC. Anti-inflammatory and Analgesic Activities of the Latex Containing Triterpenes from *Himatanthus sucuuba*. Planta Medica 2000; 66(3) 284-286.
- [92] Silva JR de A, Rezende CM, Pinto AC, Amaral ACF. Cytotoxicity and Antibacterial Studies of Iridoids and Phenolic Compounds Isolated from the Latex of *Himatanthus sucuuba*. African Journal of Biotechnology 2010; 9(43) 7357-7360.

- [93] Vilegas JHY, Hachich EM; Garcia M, Brasileiro A, Carneiro MAG, Campos VLB. Antifungal Compounds from Apocynaceae Species. *Revista Latinoamericana de Química* 1995; 23(2) 73-75.
- [94] Vilegas JHY, Hachich EM, Garcia M, Brasileiro A, Carneiro MAG, Campos VLB. Antifungal compounds from Apocynaceae species. *Revista Latinoamericana de Química* 1992; 23(1) 44-45.
- [95] Ferreira JLP, Amaral ACF, Araújo RB, Carvalho JR, Proença CEB, Fraga SAPM, Silva JR de A. Pharmacognostical Comparison of three Species of *Himatanthus*. *International Journal of Botany* 2009; 5(2) 171-175.
- [96] Bobbitt J.M., Segebarth K.P. The Iridoid Glycosides and Similar Substances. In: Battersby A.R. and Taylor W.I. (eds.) *Cyclopentanoid Terpene Derivatives*. New York: Marcel Dekker; 1969. p25-31.
- [97] Ye G, Li ZX, Xia GX, Peng H, Sun ZL, Huang CG. A New Iridoid Alkaloid from the Flowers of *Plumeria rubra* L. cv. *acutifolia*. *Helvetica Chimica Acta* 2009; 92(12) 2790-2794.
- [98] Stephens PJ, Pan JJ, Krohn K. Determination of the Absolute Configurations of Pharmacological Natural Products via Density Functional Theory Calculations of Vibrational Circular Dichroism: The New Cytotoxic Iridoid Prismatomerin. *Journal of Organic Chemistry* 2007; 72(20) 7641-7649.
- [99] Little JE, Johnstone DB. Plumericin: An Antimicrobial Agent from *Plumeria multiflora*. *Archives of Biochemistry* 1951; 30 445-452.
- [100] Wood CA, Lee K, Vaisberg AJ, Kingston DGI, Neto CC, Hammond GB. A Bioactive Spirolactone Iridoid and Triterpenoids from *Himatanthus sucuuba*. *Chemical & Pharmaceutical Bulletin* 2001; 49(11) 1477-1478.
- [101] Dobhal MP, Li G, Gryshuk A, Bhatanagar AK, Khaia SD, Joshi YC, Sharma MC, Oseroff A, Pandey RK. Structural Modifications of Plumieride Isolated from *Plumeria bicolor* and the Effect of these Modifications on In Vitro Anticancer Activity. *Journal of Organic Chemistry* 2004; 69(19) 6165-6172.
- [102] Sousa LMA, Neto, RLM, Schmidt FN, Oliveira MR. Anti-mitotic activity Towards Sea Urchin Eggs of Dichloromethane Fraction Obtained from *Allamanda schottii* Pohl (Apocynaceae). *Brazilian Journal of Pharmacognosy* 2009; 19(2A) 349-352.
- [103] Waltenberger B, Rollinger JM, Griesser UJ, Stuppner H, Gelbrich T. Plumeridoid C from the Amazonian Traditional Medicinal Plant *Himatanthus sucuuba*. *Acta Crystallographica Section C: Crystal Structure Communications* 2011; 67(10) 409-412.
- [104] Mahran GH, Abdel Wahab SM, Salah AM. Isolation and Quantitative Estimation of Plumieride from the different Organs *Plumeria rubra* and *Plumeria rubra* var. *alba*. *Planta Medica* 1974; 25(3) 226-230.

- [105] Mahran GH, Abdel-Wahab SM, Ahmed MS. Isolation and Estimation of Plumieride from Different Organs of *Plumeria rubra* L. and *Plumeria rubra* L. var. *alba*. Bulletin of the Faculty of Pharmacy (Cairo University) 1975a; 12(2)133-139.
- [106] Schmid IH, Bickel H, Meijer ThM. Zur Kenntnis des Plumierids. 1. Mitteilung. Helvetica Chimica Acta 1952; 35(1)415-427.
- [107] Yeowell DA, Schmid H. Zur Biosynthese des Plumierids. Experientia 1964; 20(5) 250-252.
- [108] Rangaswami S, Venkata Rao E. Chemical Components of *Plumeria alba*. Proceedings – Indian Academy of Sciences Section A 1960; 52A 173-181.
- [109] Afifi MS, Salama OM, Gohar AA, Marzouk NA. Iridoids with Antimicrobial Activity from *Plumeria alba*. Bulletin of Pharmaceutical Sciences, Assiut University 2006; 29(1) 215-223.
- [110] Adam G, Khoi NH, Bergner C, Lien NT. Natural Products from Vietnamese Plants. Part. 3. Plant Growth Inhibiting Properties of Plumieride from *Plumeria obtusifolia*. Phytochemistry 1979; 18(8) 1399-1400.
- [111] Halpern O, Schmid H. Zurkenntnis de Plumierids. 2. Mitteilung. Helvetica Chimica Acta 1958; 41(4) 1109-1154.
- [112] Inouye H, Uobe K, Hirai M, Masada Y, Hashimoto K. Studies on Monoterpene Glucosides and Related Natural Products. XXX. Gas Chromatography and Gas Chromatography-Mass Spectrometry of Iridoid and Secoiridoid Glucosides. Journal of Chromatography 1976; 118(2) 201-216.
- [113] Grignon-Dubois M, Rezzonico B, Usubillaga A, Vojas LB. Isolation of Plumieride from *Plumeria inodora*. Chemistry of Natural Compounds 2005; 41(6) 730-731.
- [114] Gupta RS, Bhatnager AK, Joshi YC, Sharma R, Sharma A. Effects of plumieride, an iridoid on spermatogenesis in male albino rats. Phytomedicine 2004; 11(2-3) 169-174.
- [115] Rebouças S de O, Grivicich I, dos Santos MS, Rodriguez P, Gomes MD, de Oliveira SQ, da Silva J, Ferraz ABF. Antiproliferative Effect of a Traditional Remedy, *Himatantthus articulatus* Bark, on Human Cancer Cell Lines. Journal of Ethnopharmacology 2011; 137(1) 926-929.
- [116] Coppen JJW. Iridoids with Algicidal Properties from *Allamanda cathartica*. Phytochemistry 1983; 22(1) 179-182.
- [117] Yamauchi T, Abe F, Taki M. Protoplumericin, an Iridoid Bis-Glucoside in *Allamanda neriifolia*. Chemical and Pharmaceutical Bulletin 1981; 29(10) 3051-3055.
- [118] Boonclarm D, Sornwatana T, Arthan D, Kongsaree P, Svasti J. β -Glucosidase Catalyzing Specific Hydrolysis of an Iridoid β -Glucoside from *Plumeria obtuse*. Acta Biochimica et Biophysica Sinica 2006; 38(8) 563-570.

- [119] Siddiqui BS, Ilyas F, Rasheed M, Begum S. Chemical Constituents of Leaves and Stem Bark of *Plumeria obtusa*. *Phytochemistry* 2004; 65(14) 2077-2084.
- [120] Ye G, Yang YL, Xia GX, Fan MS, Huang CG. Complete NMR Spectral Assignments of Two New Iridoid Diastereoisomers from the Flowers of *Plumeria rubra* L.CV. *acutifolia*. *Magnetic Resonance in Chemistry* 2008; 46(12) 1195-1197.
- [121] Kitagawa I, Shibuya H; Baek IN, Yokokawa Y, Nitta A, Wiriadinata H, Yoshikawa M. Pulosarioside, a new bitter trimeric-iridoid diglucoside, from an Indonesian jamu, the bark of *Alyxia reinwardtii* BL. (Apocynaceae). *Chemical & Pharmaceutical Bulletin* 1988; 36(10) 4232-4235.
- [122] Sticher O. Theveside, a New Iridoid Glucoside from *Thevetia peruviana* (Apocynaceae). *Tetrahedron Letters* 1970; 11(36) 3195-3196.
- [123] Sticher O. Theveside, a New Iridoid-Glucoside from *Thevetia peruviana* (*Thevetia neriiifolia*) (Apocynaceae). *Pharmaceutica Acta Helveticae* 1971; 46(3) 156-166.
- [124] Inouye H, Nishimura T. Iridoids Glucosides of *Cerbera manghas*. *Phytochemistry* 1972; 11(5) 1852.
- [125] Abe F., Chen R.F., Yamauchi T, Ohashi H. Iridoids from the Roots of *Thevetia peruviana*. *Chemical & Pharmaceutical Bulletin* 1995b; 43(3) 499-500.
- [126] Sticher O, Schmid H. Theviridosid, ein iridoid-glucosid aus *Thevetia peruviana* (Pers.) K. Schum. (*Thevetia neriiifolia* Juss.) Vorläufige Mitteilung. *Helvetica Chimica Acta* 1969; 52(2) 478-481.
- [127] Abe F, Yamauchi T, Xahara S, Nohara T. Minor Iridoids from *Thevetia peruviana*. *Phytochemistry* 1995; 38(3)793-794.
- [128] Changwicht K, Khorana N, Suwanborirux K, Waranuch N, Limpeanchob N, Wisuitiprot W, Suphrom N, Ingkaninan K. Bisindole alkaloids and secoiridoids from *Alstonia macrophylla* Wall. ex G. Don. *Fitoterapia* 2011; 82(6) 798-804.
- [129] Zhu W-M, He H-P, Fan L-M, Shen Y-M, Zhou J, Hao X-J. Components of Stem barks of *Winchia calophylla* A. DC. and their bronchodilator activities. *Acta Botanica Sinica* 2005; 47(7) 892-896.
- [130] Basu D, Chaterjee A. Occurrence of Plumericin in *Nerium indicum*. *Indian Journal of Chemistry* 1973; 11 297.
- [131] Wu H-K, Chuang W-C, Sheu S-J. Separation of nine iridoids by capillary electrophoresis and high-performance liquid chromatography. *Journal of Chromatography A* 1998; 803(1-2) 179-187.
- [132] Suomi J, Wiedmer SK, Jussila M, Riekkola M-L. Analysis of eleven iridoid glycosides by micellar electrokinetic capillary chromatography (MECC) and screening of plant samples by partial filling (MECC)-electrospray ionization mass spectrometry. *Journal of Chromatography A* 2002; 970(1-2) 287-296.