



## *Eratyrus mucronatus* Stål, 1859 and *Panstrongylus rufotuberculatus* (Champion, 1899) (Hemiptera, Reduviidae, Triatominae): first records in a riverside community of Rio Negro, Amazonas state, Brazil

Amanda Coutinho de Souza<sup>1\*</sup>, José Rodrigues Coura<sup>1</sup>, Catarina Macedo Lopes<sup>2</sup>,  
Angela Cristina Verissimo Junqueira<sup>1</sup>

**1** Laboratório de Doenças Parasitárias, Instituto Oswaldo Cruz, Rio de Janeiro, RJ, Brazil • ACS: amandacoutinhodesouza@gmail.com • JRC: coura@ioc.fiocruz.br • ACVJ: junqueir.rlk@terra.com.br

**2** Laboratório Interdisciplinar de Vigilância Entomológica em Diptera e Hemiptera, Instituto Oswaldo Cruz, Rio de Janeiro, RJ, Brazil • aniratac macedo@gmail.com

\* Corresponding author

### Abstract

We report the first records of *Eratyrus mucronatus* Stål, 1859 and *Panstrongylus rufotuberculatus* (Champion, 1899) (Hemiptera, Reduviidae, Triatominae) in a riverside community of the Rio Negro in the Brazilian Amazon. Health promotion activities were carried out with the human population and agents to combat endemic diseases. The entomological findings reinforce the importance of community participation in the surveillance and control of triatomines. Actions to disseminate knowledge on Chagas disease and vectors strengthened the entomological surveillance in the middle Rio Negro.

### Key words

Chagas disease, first record, Barcelos, health promotion, triatomines, entomological surveillance

---

Academic editor: Hélcio Gil-Santana | Received 20 November 2020 | Accepted 9 April 2021 | Published 11 June 2021

**Citation:** Souza AC, Coura JR, Lopes CM, Junqueira ACV (2021) *Eratyrus mucronatus* Stål, 1859 and *Panstrongylus rufotuberculatus* (Champion, 1899) (Hemiptera, Reduviidae, Triatominae): first records in a riverside community of Rio Negro, Amazonas state, Brazil. Check List 17 (3): 905–909. <https://doi.org/10.15560/17.3.905>

---

### Introduction

Currently, 153 living and three fossil species are known to belong to the subfamily Triatominae (Reduviidae) (Alevi et al. 2020; Galvão 2020; Zhao et al. 2021). In the Brazilian Amazon, some species of triatomines have been found to be naturally infected by *Trypanosoma cruzi* (Chagas, 1909), including *Eratyrus mucronatus* Stål, 1859, *Microtriatoma trinidadensis* (Lent, 1951),

*Triatoma sordida* (Stål, 1859), *Panstrongylus lignarius* (Walker, 1837), *Panstrongylus rufotuberculatus* (Champion, 1899), *Rhodnius brethesi* Matta, 1919, *Rhodnius neglectus* Lent, 1954, *Rhodnius paraensis* Sherlock, Guittón & Miles, 1977, *Rhodnius robustus* Larrousse, 1927, and *Rhodnius montenegrensis* Rosa et al. 2012 (Bilheiros et al. 2018; Coura et al. 2018; Ramos et al.

2018; Madeira et al. 2020).

Oral transmission in humans linked to the consumption of açaí juice contaminated with *T. cruzi* in the Rio Negro microregion (Amazonas state, Brazil) emphasizes the importance of strengthening the epidemiological surveillance service (Souza-Lima et al. 2013). An outbreak with 17 cases in Santa Isabel do Rio Negro municipality confirmed and seroepidemiological and clinical studies of *T. cruzi* infection using samples from the population, confirm the endemicity of Chagas disease in the region (Brum-Soares et al. 2010; Souza-Lima et al. 2013; Coura et al. 2018).

Health education is an important part of prevention of Chagas disease and improving the human population's general living conditions and overall health. Health knowledge educates the population in changes of behavior which can prevent disease and preserve and improve life (Oliveira and Gonçalves 2004).

The control of Chagas disease in the Brazilian Amazon is predominantly done through educating the general population and healthcare professionals. Technicians in laboratories for diagnosing malaria, need to be trained to diagnose the *T. cruzi* infection. The agents of endemic diseases such as *T. cruzi* should be searched for and triatomines identified. Doctors and nurses need the training to treat the disease (Coura and Junqueira 2015).

We report the first records of *E. mucronatus* and *P.*


*rufotuberculatus* in a riverside community in the Rio Negro microregion, Amazonas.

## Methods

Researchers of the Laboratório de Doenças Parasitárias developed health education to disseminate the knowledge about Chagas disease and vectors in the riverside community of Piloto, municipality of Barcelos on the middle Rio Negro, Amazonas state, Brazil (Fig. 1, 2). Health and endemic disease agents participated in all activities.

Information on Chagas disease and its transmission cycle, whether by labor or orally, were developed for extractors of piassava fiber and palm fruits and distributed to the general population. Triatomine diversity, *T. cruzi* transmission modes, and prevention measures were included in this information.

On 15 November 2017, a year after health education had begun in the Piloto, seven specimens of insects, supposedly triatomines, were delivered to the agents. The insects were manually collected near homes in the Piloto community. The specimens were sent to the Laboratório de Doenças Parasitárias at Oswaldo Cruz Institute, Oswaldo Cruz Foundation (Fiocruz), Rio de Janeiro. The specimens were morphologically identified in the Laboratório Interdisciplinar de Vigilância Entomológica



**Figure 1.** Maps showing the distribution of (A) *Eratyrus mucronatus* and (B) *Panstrongylus rufotuberculatus*. C. Location of the riverside community of Piloto, Barcelos, Amazonas, Brazil.



**Figure 2.** The riverside community of Piloto, Barcelos, Amazonas, Brazil.

em Diptera e Hemiptera, at Fiocruz using the dichotomous keys by Lent and Wygodzinsky (1979) and Galvão (2014). The triatomines were deposited in the Triatominae Collection at the Oswaldo Cruz Institute (CTIOC).

## Results

### *Eratyrus mucronatus* Stål, 1859

**New records.** BRAZIL – Amazonas • Barcelos, riverside community of Piloto; 00°53.60'S, 062°59.23'W; elev. 43 m; 15.XI.2017; collected by community residents

near houses, a place with several species palm trees; 1 ♂, CTIOC 12348.

**Identification.** In the genus *Eratyrus* Stål, 1859 the main characters used for identification were the head eyes longer than wide; antennae inserted away from the apex of the head and eyes; ocelli situated on sharp protrusions of the disc in the post-ocular region of the head and very long, oblique, spine-shaped scutellum. The species was determined by having the anterior pronotum lobe with a robust dorsal thorn, humeri of the pronotum with sharply spiny angles, and corium with a small, reddish, subapical spot (Fig. 3).

### *Panstrongylus rufotuberculatus* (Champion, 1899)

**New records.** BRAZIL – Amazonas • Barcelos, riverside community of Piloto; 00°53.60'S, 062°59.23'W; elev. 43 m; 15.XI.2017; collected by community residents near the houses, a place with several species palm trees; 1 ♂, CTIOC 12347.

**Identification.** The genus *Panstrongylus* Berg, 1879 was identified by the position of the antennae, which are inserted close to the eyes. The species was identified by the golden bristles on the dorsal surface of the body;



**Figure 3.** *Eratyrus mucronatus* Stål, 1859, male, dorsal view.



**Figure 4.** *Panstrongylus rufotuberculatus* (Champion, 1899), male, dorsal view.

anterior lobe of the pronotum with reddish tubers; short scutellum, which may be rounded, conical, or truncated at its tip; connexival segments with central dark spots; and light-green anterior wings (Fig. 4).

## Discussion

*Eratyrus mucronatus* occurs in the Brazilian states of Amazonas, Acre, Pará, Rondônia, Roraima, Maranhão, Goiás, and Mato Grosso, and extends to Colombia, Bolivia, Ecuador, French Guiana, Suriname, and Venezuela (Meneguetti et al. 2011; Obara et al. 2013; Monte et al. 2014; Gama Neto et al. 2020). It occurs in wild habitats and is generally associated with nests of Xenartha, Didelphidae, and Chiroptera in palms as well as in sympatry with colonies of *Triatoma maculata* (Erichson, 1848) and *Cavernicola pilosa* Barber, 1937 (Dias et al. 1942; Guerrero et al. 1981; Vivas et al. 2001). The natural infection of this species has already been recorded (Sherlock et al. 1997). Its first occurrence in Brazil was reported in the Ouro Preto do Oeste municipality, Rondônia state, in a wild environment of the Comissão Executiva do Plano da Lavoura Cacaueira reserve (Meneguetti et al. 2011). The triatomines captured in Acre were collected inside a home located in a secondary forest in a peri-urban area with palms and cattle (Obara et al. 2013).

*Panstrongylus rufotuberculatus* has been found in Mexico, Costa Rica, Panama, French Guiana, Colombia, Venezuela, Ecuador, Peru, Bolivia, and Argentina (Salomón et al. 1999; Galvão 2014). In Brazil, has been found in Acre, Amazonas, Pará, and Mato Grosso (Galvão 2014; Oliveira et al. 2019). The first specimen captured in Amazonas was at the mouth of the Rio Preto, a tributary of Rio Unini, part of the Rio Negro basin (Almeida and Santos 1973). However, infection of *P. rufotuberculatus* with *T. cruzi* has already been reported (Sousa et al. 1983; Salomón et al. 1999). Miles (1979) found specimens associated with opossums in the Brazilian Amazon; Cálderon et al. (1985) observed intra-domiciliary colonies, and Castro et al. (2010) reported the invasion of homes by adults that were attracted by the lights.

Health education builds popular participation in health services (Oliveira and Gonçalves 2004). Some studies point to an integrated action with the community, involving educational activities to guarantee the continuity of actions and strengthen the epidemiological and entomological surveillance (Falavigna-Guilherme et al. 2002; Mello et al. 1992).

We corroborate Abad-Franch et al. (2011), who emphasized the importance of encouraging public participation in insect collection. Using lectures, the human population was empowered with information on the vectors of *T. cruzi*, transmission modes, and prevention of the disease. Endemic disease agents had no knowledge of previous reports of *E. mucronatus* and *P. rufotuberculatus* in region. We suggest that dissemination of information continues in the Rio Negro microregion. Such

information reinforces entomological surveillance and supports a deeper knowledge of the region's triatomine fauna.

## Acknowledgements

We thank the people of Barcelos municipality for their contribution to the surveillance and control of Chagas disease in the middle Rio Negro. We also thank the endemic disease agent Maria José Andrade de Oliveira for the work in the municipality.

## Authors' Contributions

ACS and ACVJ developed the health promotion activities in the Piloto community. ACS, CML, and JRC wrote the manuscript; CML identified the triatomine specimens.

## References

- Abad-Franch F, Vega MC, Rolón MS, Santos WS, Rojas de Arias (2011) Community participation in Chagas disease vector surveillance: systematic review. PLoS Neglected Tropical Diseases 5 (6): e1207. <http://doi:10.1371/journal.pntd.0001207>
- Alevi KCC, Oliveira J, Garcia ACC, Cristal DC, Delgado LMG, Bittinelli IF, Reis YV, Ravazi A, Oliveira ABB, Galvão C, Azeredo-Oliveira MTV, Madeira FF (2020) *Triatoma rosai* sp. nov. (Hemiptera, Triatominae): a new species of Argentinian Chagas disease vector described based on integrative taxonomy. Insects 11: 830. <https://doi:10.3390/insects11120830>
- Almeida FB, Santos EI (1973) Triatomíneos da Amazônia II: encontro do *Panstrongylus rufotuberculatus* (Champion, 1899) no estado do Amazonas (Hemiptera, Reduviidae). Acta Amazonica 3 (3): 51–52. <https://doi.org/10.1590/1809-43921973033051>
- Bilheiros AB, Rosa JA, Oliveira J, Belintani T, Fontes G, Medeiros JF, Meneguetti DUO, Camargo LMA (2018) First report of natural infection with *Trypanosoma cruzi* in *Rhodnius montenegrensis* (Hemiptera, Reduviidae, Triatominae) in western Amazon, Brazil. Vector-Borne and Zoonotic Diseases 18 (11): 605–610. <https://doi.org/10.1089/vbz.2018.2266>
- Brum-Soares LM, Xavier SS, Sousa AS, Pereira JB, Ferreira JMBB, Costa IR, Junqueira ACV, Coura JR (2010) Morbidade da doença de Chagas em pacientes autóctones da Microrregião do Rio Negro, estado do Amazonas. Revista da Sociedade Brasileira de Medicina Tropical 43: 170–177. <https://doi.org/10.1590/S0037-86822010000200013>
- Castro MCM, Barrett TV, Santos WS, Abad-Franch F, Rafael JA (2010) Attraction of Chagas disease vectors (Triatominae) to artificial light sources in the canopy of primary Amazon rainforest. Memorias do Instituto Oswaldo Cruz 105 (8): 1061–1064. <https://doi.org/10.1590/S0074-02762010000800019>
- Coura JR, Junqueira ACV (2015) Surveillance, health promotion and control of Chagas disease in the Amazon Region medical attention in the Brazilian Amazon region: a proposal. Memorias do Instituto Oswaldo Cruz 110 (7): 825–830. <https://doi.org/10.1590/0074-02760150153>
- Coura JR, Junqueira ACV, Ferreira JMB (2018) Surveillance of seroepidemiology and morbidity of Chagas disease in the Negro River, Brazilian Amazon. Memorias do Instituto Oswaldo Cruz 113 (1): 17–23. <https://doi.org/10.1590/0074-02760170169>
- Dias E, Mello GB, Costa O, Damasceno RG, Azevedo M (1942) Investigações sobre esquistosípanose de morcegos no estado do Pará. Encontro do barbeiro “*Cavernicola pilosa*” como transmissor. Revista Brasileira de Biologia 2: 103–110.
- Jurberg J, Rodrigues JMS, Moreira FFF, Dale C, Cordeiro IRS, Lamas

- Jr VD, Galvão C, Rocha DS (2014) Coleção taxonômica. In: Cleber Galvão. (Ed.). Vetores da doença de Chagas no Brasil. Sociedade Brasileira de Zoologia, Curitiba, Brazil, 289 pp.
- Galvão C (2020) Taxonomia dos vetores da doença de Chagas: da forma à molécula, quase três séculos de história. In: Oliveira J, Alevi KCC, Camargo LMA, Meneguetti DUO (Eds.) Atualidades em medicina tropical no Brasil: vetores. Stricto Sensu, Rio Branco, Brazil, 9–37. <https://doi.org/10.35170/ss.ed.9786586283129.01>
- Gama Neto JL, Oliveira J, Rosa JA, Santos FM, Macedo VA, Silva WC (2020) Two new records of Triatominae (Hemiptera: Reduviidae) from Roraima state, Brazil. Revista Chilena de Entomología 46 (2): 321–327. <https://doi.org/10.35249/rche.46.2.20.23>
- Guerrero L, Scorza JV (1981) Las fuentes alimenticias de algunos Triatominae silvestres en los llanos centro occidentales de Venezuela. Boletín de Malaria y Salud Ambiental 21: 129–139.
- Lent H, Wygodzinsky P (1979) Revision of the triatominae (Hemiptera, Reduviidae), and their significance as vectors of Chagas disease. Bulletin American Museum of Natural History 163: 520–529.
- Madeira FP, Costa de Jesus A, Moraes MHS, Oliveira AS, Oliveira J, Melchior LAK, Camargo LMA, Meneguetti DUO, Bernarde PS (2020) Doença de Chagas: conceitos básicos de uma enfermidade negligenciada e seus vetores na amazônia occidental brasileira In: Oliveira J, Alevi KCC, Camargo LMA, Meneguetti DUO (Eds.) Atualidades em medicina tropical no Brasil: vetores. Stricto Sensu, Rio Branco, Brazil, 9–37. <https://doi.org/10.35170/ss.ed.9786586283129.01>
- Mascarenhas BM (1991) Triatomíneos da Amazônia: sobre o habitat e algumas considerações comportamentais de *Rhodnius brethesi* Matta, 1919 (Hemiptera: Reduviidae: Triatominae) na região do médio Rio Negro, Amazonas. Boletim do Museu Paraense Emílio Goeldi 7: 107–116.
- Matta A (1922) Sobre o gênero *Rhodnius* do Amazonas. Amazonas Medico 5: 161–162.
- Meneguetti DUO, Trevisan O, Rosa RM, Camargo LM (2011) First report *Eratyrus mucronatus*, Stål, 1859, (Hemiptera, Reduviidae, Triatominae), in the state of Rondonia, Brazil. Revista da Sociedade Brasileira de Medicina Tropical 44: 511–512. <https://doi.org/10.1590/S0037-86822011000400022>
- Miles MA (1979) Transmission cycles and heterogeneity of *Trypanosoma cruzi*, In: Lumsden WHR, Evans DA (Eds.) Biology of kinetoplastida. Academic Press, London, UK, 117–196.
- Monte GLS, Tadei WP, Farias TM (2014) Ecoepidemiology and biology of *Eratyrus mucronatus* Stål, 1859 (Hemiptera: Reduviidae: Triatominae), a sylvatic vector of Chagas disease in the Brazilian Amazon. Revista da Sociedade Brasileira de Medicina Tropical 47 (6): 723–727. <https://doi.org/10.1590/0037-8682-0263-2014>
- Obara MT, Cardoso AS, Pinto MCG, Souza CR, Albuquerque e Siva R, Gurgel-Gonçalves R (2013) *Eratyrus mucronatus* Stål, 1859 (Hemiptera: Reduviidae: Triatominae): first report in the state of Acre, Brazil, and updated geographic distribution in South America. Check List 9 (4): 851–854. <https://doi.org/10.15560/9.4.851>
- Oliveira AS, Ribeiro MAL, Castro GVS, Brilhante NA, Camargo LMA, Meneguetti DUO (2019) Confirmation of the occurrence of *Panstrongylus rufotuberculatus* (Champion, 1899) in the state of Acre, western Amazon. Revista da Sociedade Brasileira de Medicina Tropical 52: e20180388. <https://doi.org/10.1590/0037-8682-0388-2018>
- Oliveira HM, Gonçalves MJF (2004) Educação em Saúde: uma experiência transformadora. Revista Brasileira de Enfermagem 57 (6): 761–763. <https://doi.org/10.1590/S0034-71672004000600028>
- Ramos LJ, Souza JL, Souza CR, Oliveira J, Rosa JA, Camargo LMA, Cunha RM, Meneguetti DUO (2018) First report of *Triatoma sordida* Stål, 1859 (Hemiptera, Reduviidae, Triatominae) in the state of Acre and Brazilian western Amazon. Revista da Sociedade Brasileira de Medicina Tropical 51 (1): 77–79.
- Salomon O, Ripoll C, Rivetti E, Carcavallo R (1999) Presence of *Panstrongylus rufotuberculatus* (Champion, 1899) (Hemiptera: Triatominae) in Argentina. Memorias do Instituto Oswaldo Cruz 94: 285–288. <https://doi.org/10.1590/S0074-0276199900300002>
- Sherlock IA, Carcavallo RU, Galíndez-Girón I (1997) List of natural and experimental flagellates infections in several Triatominae species. In: Carcavallo RU, Galíndez Girón I (Eds.) Atlas of Chagas' disease vectors in the Americas. Editora Fundação Oswaldo Cruz; Rio de Janeiro, Brazil, 289–298.
- Sousa OE, Wolda H, Batista LF (1983) Triatominos encontrados en el ambiente silvestre en la isla de Barro Colorado. Revista de Medicina Panama 8: 50–55.
- Souza AC, Catalá S, Carbajal de La Fuente AL, Junqueira ACV (2017) Phenotype variability of the Amazon species *Rhodnius brethesi* (Hemiptera: Reduviidae). Journal Medical Entomology 54 (4): 909–916. <https://doi.org/10.1093/jme/tjx038>
- Souza-Lima RC, Barbosa MGV, Coura JR, Arcanjo ARL, Nascimento AS, Ferreira JMBB, Magalhães LK, Albuquerque BC, Araújo GAV, Guerra JAO (2013) Outbreak of acute Chagas disease associated with oral transmission in the Rio Negro region, Brazilian Amazon. Revista da Sociedade Brasileira de Medicina Tropical 46: 510–514.
- Vivas AS, Barazarte H, Fernández DM (2001) Primer registro de *Eratyrus mucronatus* Stål, 1959 (Hemiptera: Reduviidae) en el ambiente domiciliario en Venezuela. EntomoTropica 16 (3): 215–217.
- Zhao Y, Galvão C, Cai W (2021) *Rhodnius micki*, a new species of Triatominae (Hemiptera, Reduviidae) from Bolivia. ZooKeys 1012: 71–93. <https://doi.org/10.3897/zookeys.1012.54779>