



Distinct Roles of Hemocytes at Different Stages of Infection by Dengue and Zika Viruses in *Aedes aegypti* Mosquitoes

Thiago H. J. F. Leite^{1†}, Álvaro G. A. Ferreira^{1,2†}, Jean-Luc Imler³ and João T. Marques^{1,3*}

¹ Department of Biochemistry and Immunology, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil, ² Mosquitos Vetores: Endossimbiontes e Interação Patógeno-Vetor, Instituto René Rachou – Fiocruz, Belo Horizonte, Brazil, ³ Université de Strasbourg, CNRS UPR9022, Inserm U1257, Strasbourg, France

OPEN ACCESS

Edited by:

Luc Swevers,
National Centre of Scientific Research
Demokritos, Greece

Reviewed by:

Emilie Pondeville,
MRC-University of Glasgow Centre
For Virus Research (MRC),
United Kingdom
Ryan C. Smith,
Iowa State University, United States

*Correspondence:

João T. Marques
jtm@ufmg.br

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Comparative Immunology,
a section of the journal
Frontiers in Immunology

Received: 29 January 2021

Accepted: 30 April 2021

Published: 13 May 2021

Citation:

Leite THJF, Ferreira ÁGA, Imler J-L
and Marques JT (2021) Distinct Roles
of Hemocytes at Different Stages of
Infection by Dengue and Zika Viruses
in *Aedes aegypti* Mosquitoes.
Front. Immunol. 12:660873.
doi: 10.3389/fimmu.2021.660873

Aedes aegypti mosquitoes are vectors for arboviruses of medical importance such as dengue (DENV) and Zika (ZIKV) viruses. Different innate immune pathways contribute to the control of arboviruses in the mosquito vector including RNA interference, Toll and Jak-STAT pathways. However, the role of cellular responses mediated by circulating macrophage-like cells known as hemocytes remains unclear. Here we show that hemocytes are recruited to the midgut of *Ae. aegypti* mosquitoes in response to DENV or ZIKV. Blockade of the phagocytic function of hemocytes using latex beads induced increased accumulation of hemocytes in the midgut and a reduction in virus infection levels in this organ. In contrast, inhibition of phagocytosis by hemocytes led to increased systemic dissemination and replication of DENV and ZIKV. Hence, our work reveals a dual role for hemocytes in *Ae. aegypti* mosquitoes, whereby phagocytosis is not required to control viral infection in the midgut but is essential to restrict systemic dissemination. Further understanding of the mechanism behind this duality could help the design of vector-based strategies to prevent transmission of arboviruses.

Keywords: Zika virus, dengue virus, cellular immunity, macrophage-like cells, *Aedes aegypti*, vector mosquitoes, hemocytes

INTRODUCTION

Aedes aegypti mosquitoes are vectors for a wide variety of arthropod-borne viruses (arboviruses) (1). How these mosquitoes recognize and respond to viral infection is a central question that directly affects their vector competence. The understanding of antiviral responses in insects has greatly benefited from work in the fruit fly *Drosophila melanogaster* (2). Work in this model organism has identified many important antiviral defense mechanisms such as RNA interference (RNAi), Jak-STAT and STING (3–12). Later work in mosquitoes has shown that RNAi and Jak-STAT are important for the control of arbovirus infections (13–18). Interestingly, despite being widely conserved throughout evolution, STING has been lost in mosquitoes (19).

In addition to these well-known innate immunity pathways, the *Drosophila* model has also highlighted the role of circulating macrophage-like cells, referred to as hemocytes, in the control of

viral infection (20–22). Cellular immunity in insects includes phagocytosis of foreign bodies, nodulation, wound healing and the encapsulation of pathogens (23–27). Hemocytes can be freely circulating in the insect hemolymph or associated with tissues but these populations seem to be highly dynamic and interchangeable (28). Hemocytes are often recruited to infected tissues, which increases the chances of coming into contact with the pathogen to be cleared by phagocytosis (28, 29). A good example of hemocyte recruitment during an infection is in the case of *Plasmodium*, the malaria parasite. Invasion of the midgut of *Anopheles* mosquitoes by *Plasmodium* ookinetes promotes hemocyte recruitment and release of components of the mosquito complement system, promoting pathogen elimination (30–34). Despite the importance of hemocytes for the clearance of bacteria and *Plasmodium* in mosquitoes, little is known about their role during viral infections, particularly arboviruses such as dengue (DENV) and Zika (ZIKV) viruses. DENV and ZIKV belong to the *Flaviviridae* family and, together with the alphavirus chikungunya virus (CHIKV) are among the most important arboviruses transmitted by *Ae. aegypti* mosquitoes causing infections worldwide (1). Similar to the malaria parasite, arboviruses are acquired orally during blood feeding by mosquitoes, and the gut represents a physical barrier that hinders the passage of the viral particles to the mosquito hemocele (35). Reaching the hemocele is a necessary step for the virus to spread systemically and reach the salivary glands where it can be transmitted to a vertebrate host (21–23). During systemic infection, several tissues may host viral replication, including hemocytes themselves, but it is unclear how they contribute to amplification of the virus (36–38). Despite this increasing knowledge about the functions of hemocytes in mosquitoes, the role of cellular immunity in the antiviral defense remains largely unknown.

In this work, we investigated the involvement of hemocytes in the control of DENV and ZIKV in *Ae. aegypti* mosquitoes. Our results suggest a complex role for hemocytes. We show that hemocytes were recruited to the midgut in response to the presence of the virus but, once there, their phagocytic activity seems to facilitate viral replication although other functions may play a role in the antiviral defense. In contrast, during the systemic phase of the infection, inhibition of phagocytosis by hemocytes led to increased viral infection pointing to a more traditional role in antiviral immunity. Together our results indicate that hemocytes have dual roles in the control of arboviruses in *Ae. aegypti* mosquitoes depending on tissue affected and the stage of the infection in the vector.

MATERIALS AND METHODS

Indirect Immunofluorescence Assays

Mosquitoes were anaesthetized on ice and then were injected with 250 nanoliters of 20% paraformaldehyde for hemocyte fixation in midgut basal lamina. After 20 minutes, midguts were dissected in 4% paraformaldehyde diluted in phosphate-buffered saline (PBS) (13 mM NaCl, 0.7 mM Na₂HPO₄, 1 mM NaH₂PO₄ at pH 7.2) (PBS). The remaining midguts were fixed in the same solution

for 20 minutes, and then washed three times in PBS and then incubated with blocking solution PBSBT (1× PBS + 1% BSA + 0.1% Triton X-100) for 15 minutes at room temperature. Samples were then incubated overnight with 4G2 monoclonal antibody for Flavivirus E protein (ATCC: HB-112, used at 1:50 in PBSBT) at 4°C. Midguts were washed three times with PBSBT (5 min each) and incubated for 2 h with constant rocking at 25°C with goat anti-mouse IgG antibody (Invitrogen). Midguts were washed three times with PBSBT (5 min each) and incubated for 15 min with DAPI (Molecular Probes, 1:500), and phalloidin-rhodamine (Molecular Probes, 1:500). Then the midguts were washed in PBS and placed onto slides. Images were obtained with an LSM 880 microscope (Zeiss).

Mosquito Perfusion to Obtain Circulating Hemocytes

Circulating hemocytes were obtained by perfusion of adult mosquitoes as described (39) with modifications. Briefly, mosquitoes were injected with 1 µL of anticoagulant buffer solution (70% PBS 1x (pH 7.0) + 30% citrate buffer (98 mM NaOH, 186 mM NaCl, 1.7 mM EDTA and 41 mM citric acid, buffer pH 4.5) and were incubated on a petri dish on ice for 10 min to let hemocytes dissociate from tissues. The last two segments of abdomen were cut to create an opening, which was positioned onto a microscope slide. Each individual mosquito was positioned vertically and then injected with 3 µL of the same anticoagulant buffer solution in the lateral side of torax using a microinjector (Nanoject III). The injection pressure forced the diluted hemolymph to exit the opening made in the final portion of the abdomen and onto the microscope slide. The hemolymph was incubated at room temperature for 20 min in order to let the hemocytes adhere to the slides. Hemocytes were fixed in 4% paraformaldehyde for 20 min, washed three times in PBS and then incubated with blocking solution PBSBT (1× PBS + 1% BSA + 0.1% Triton X-100) for 15 minutes at room temperature. Slides were incubated for 15 min with DAPI (Molecular Probes, 1:500) and phalloidin-rhodamine (Molecular Probes, 1:40), followed by 3 washes in PBS. Cells were visualized in a fluorescence microscope for counting. To visualize infected hemocytes, the 4G2 monoclonal antibody against Flavivirus E protein was used.

Hemocyte Labeling *In Vivo*

For *in vivo* hemocyte staining we used Vybrant™ CM-DiI Cell-Labeling Solution (Invitrogen™) essentially as described (29). Briefly, female mosquitoes were placed on petri dish on ice and injected with 150 nanoliters containing 100 µM CM-DiI, freshly prepared in sterile water, after blood or sugar meal at specific time points. Injections were done using a nano-injector Nanoject III (Drummond Scientific Company). After injections, mosquitoes are placed on cages at 28°C until specific time points for midguts dissections.

Quantification of the Infection Area in the Midgut

Area measurements and hemocyte counting were performed using ImageJ v1.53c (<https://imagej.nih.gov/ij/>). All images

were acquired under identical conditions, digitized, converted to RGB image and stored in an uncompressed tagged image file format (.tiff). Infection area computing was performed using ImageJ. The following steps were performed for all images to quantify the area of infection in the midgut, as shown in **Supplementary Figure 1**: step 1, color-deconvolution was used to isolate red, green and blue spectra and select the image corresponding to virus infection staining; step 2, a projection final image was generated using all acquired series of z-stack confocal images using the tool “Image > Stacks > Z-project function”; step 3, the projection image was processed into 8 bits image type; step 4, the midgut outline was delimited; step 5, the area outside of midgut delimitation was erased by using the “clear outside” function; steps 6 and 7, optical density was assessed by setting a threshold using the “threshold tool”, and a maximum threshold was set; steps 8 and 9, the function “Measure” in the ‘Analyze’ tool menu was used to calculate the optical density and compute the midgut infection area.

Quantification of Hemocytes in the Midgut

Hemocyte numbers were quantified in the confocal microscopy images of midguts. The following steps were performed using ImageJ for all images: step 1, color-deconvolution was used to isolate red, green and blue spectra and select the images corresponding to hemocytes cell tracker and DNA staining; step 2, for each color, a projection final image was generated using all acquired series of z-stack confocal images using the tool “Image > Stacks > Z-project function”; step 3, the projection image was processed into 8 bits image type; step 4, the number of hemocytes was then counted using the ITCN (Image-based Tool for Counting Nuclei); step 5, for all hemocytes automatically identified in the hemocytes cell tracker color we additionally confirmed the presence of nuclei using the DNA staining image and reject the counts that do not presented a nucleus.

Inhibition of Phagocytosis by Injection of Latex Beads

To block the phagocytic activity of hemocytes in mosquitoes, we adapted protocols previously used for *Drosophila* (20). Adult mosquitoes were injected with 69 nanoliters of latex microspheres (CML Latex Beads, 4% w/v, 0.3 μm , ThermoFisher). Latex beads were washed and resuspended at a 2X concentration in PBS before injections. In order to quantify the inhibition of phagocytosis, we first injected regular latex beads followed by injection of red fluorescent beads (FluoSpheres™ Carboxylate-Modified Microspheres, 0.2 μm , dark red fluorescent (660/680), 2% solids, ThermoFisher) two days later. Perfusions were done 4 and 8 days after the first injection and the total number of hemocytes was counted as well as the percentage of cells with red beads.

RT-qPCR

Total RNA (200 ng) extracted from individual insects or individual tissues was reverse transcribed using Moloney murine leukaemia virus reverse transcriptase. cDNA was subjected to quantitative PCR (qPCR) using the kit Power SYBR Green Master Mix (Applied Biosystems), following the

manufacturer’s instructions. Primers used for quantitative PCR (qPCR) were as follows: RPL32 (forward, 5’-ACTTCTTCGTC CGCTTCTTG-3’; reverse, 5’-AGCCGCGTGTGTACTCTG-3’), DENV1 (forward, 5’-TCGGAAGCTTGCTTAACGTAG-3’; reverse, 5’ TCCGTTGGTTGTTTCATCAGA-3’), ZIKV (forward, 5’-TCAAACGAATGGCAGTCAGTG-3’; reverse, 5’-GCTTGTTGAAGTGGTGGGAG-3’) as previously described (14).

Mosquito Rearing and Infections

All experiments were carried out using *Ae. aegypti* Bangkok strain. Mosquitoes were maintained in an incubator at 28°C and 70–80% relative humidity, in a 12:12 h light:dark photoperiod, and with 10% sucrose solution ad libitum. For mosquito infections, we used previously described models for flavivirus infections using mice or artificial membrane feeding. Isolates of DENV4 (H241 strain), DENV1 (MV09) and ZIKV (PE243/2015) were previously described (14). As a mouse model, we utilized DENV1 and ZIKV infection of interferon alpha/beta and gamma receptor-deficient (AG129) animals (14). Mice were injected intraperitoneally with 10^6 pfu/mL of virus. Infected mice were anaesthetized at 3 days post injection (peak of viraemia) using ketamine/xylazine (80/8 mg kg⁻¹) and placed on top of the netting-covered containers with 5- to 6-day-old adult mosquito females. For infections by artificial membrane feeding, 5-6 day old adult females were starved for 24h and fed with a mixture of blood and virus supernatant containing 10^7 pfu/mL of DENV4 or 10^6 pfu/mL of ZIKV utilizing a glass artificial feeding system covered with pig intestine membrane, essentially as described (14). Mosquitoes were allowed to feed for 1 h. After blood feeding, fully engorged females were selected and harvested individually for midgut dissection at different time points. For direct systemic infections by intrathoracic injections, mosquitoes were anaesthetized with CO₂ and kept on ice during the whole procedure. 4-day-old females were intrathoracically injected with 69 nL of L15 media containing virus (5 or 50 pfu), using a nano-injector Nanoject III (Drummond Scientific Company). Mosquitoes were harvested at different days post injection for RNA extraction. Tissues or mosquitoes were ground in TRIzol (Invitrogen) using glass beads. Total RNA was extracted from individual mosquitoes or individual tissues according to the manufacturer’s protocol.

RESULTS

DENV and ZIKV Trigger Accumulation of Hemocytes in the Mosquito Midgut

Hemocytes play an important role in mosquito immunity but their function in the antiviral response against arboviruses remains unclear. Here, we first analyzed whether hemocytes would respond to the presence of arboviruses, DENV and ZIKV, in the blood meal (**Figure 1A**). Others have observed that blood feeding induces an increase in the numbers of hemocytes in mosquitoes (40, 41). Here we observed that there is also an increase in the number of hemocytes associated with the midgut compared to mosquitoes that were kept on sugar at 4

and 8 days post feeding (Figures 1B, C). At the earlier time point, there was no significant difference between the number of hemocytes associated with the midgut of mosquitoes fed with blood or blood and virus (Figure 1B). However, at 8 days post feeding, numbers of midgut-associated hemocytes were significantly higher in the presence of DENV or ZIKV when compared to a control blood meal (Figure 1C). Notably, these hemocytes do not seem to be recruited to sites of viral replication. We observed that hemocytes were often found dispersed throughout the midgut and not necessarily concentrated around regions with staining of the viral E protein as an indication of infection (Figures 1D–G). These results suggest that the presence of virus particles in the blood meal increases the number of hemocytes associated to midgut possibly by providing signals for increased recruitment or longer retention of these cells in the organ. The delayed effect at later times post infection also suggests that the accumulation of hemocytes may require prolonged stimuli.

Phagocytosis by Hemocytes Does Not Contribute to the Control of DENV and ZIKV in the Midgut

Increased numbers of hemocytes in the midgut in response to arboviruses in the blood meal suggests that these cells may play a role in the antiviral defense. Phagocytosis is a major function of hemocytes. Indeed, blocking phagocytosis by hemocytes or complete genetic ablation of these cells leads to decreased resistance to viruses in *Drosophila* (20–22). Here, we decided to use injection of latex beads into mosquitoes, which is often used as a strategy to over-load hemocytes and inhibit their phagocytic capacity (20, 21, 42). In our experiments, we observed that injection of beads seemed to decrease the number of circulating hemocytes in the mosquito but that was not significant (Supplementary Figure 2A). The number of hemocytes was estimated in a fraction of the hemolymph obtained by perfusion of mosquitoes with a low volume of buffer. Although this strategy recovered smaller numbers of hemocytes compared to other methods (39, 40), it still allowed us to compare numbers of cells between two conditions, which was our objective. Using the same strategy, we observed that phagocytosis by hemocytes was significantly inhibited by latex beads 2 days after their injection into *Ae. aegypti* mosquitoes (Supplementary Figure 2B). We next analyzed the effect of latex beads in mosquitoes that were given a blood meal containing DENV or ZIKV 2 days later, during the time when phagocytosis is inhibited (Figure 2A). Blocking phagocytosis did not affect significantly the area of infection by DENV or ZIKV in the midgut at 4 days post feeding (Figures 2B, C). In contrast, at 8 days post feeding, we observed that midgut of mosquitoes injected with latex beads had a significantly decreased area of infection by DENV and ZIKV compared to controls (Figures 2B–G). Importantly, injection of latex beads did not significantly change the total size of the midgut at the same time point but affected the absolute infection area suggesting that the kinetics of viral replication itself was affected (Supplementary Figure 3). At 4 days post feeding, injection of beads caused a reduction in viral RNA levels in DENV and ZIKV infected mosquitoes, although it

was only significant for the latter (Supplementary Figure 4). At 8 days post feeding, DENV and ZIKV RNA levels were also significantly decreased in midguts from mosquitoes injected with latex beads compared to controls (Figures 2H, I). These results suggest that blocking the phagocytic activity of hemocytes using latex beads led to decreased viral replication in the midgut of mosquitoes. Notably, we consistently observed that latex beads increased the number of midgut-associated hemocytes in sugar and blood fed mosquitoes, independent of virus infection (Supplementary Figure 5). Latex beads also increased numbers of hemocytes in the midgut of DENV and ZIKV infected mosquitoes at 4 and 8 days post feeding (Figures 2J, K). During viral infection, latex beads had a less striking effect on hemocyte numbers at later time points since infection itself led to accumulation of hemocytes in the midgut (Figure 1C). This increased accumulation of hemocytes in the midgut induced by latex beads preceded the reduction in viral levels. Thus, we cannot rule out that increased accumulation of hemocytes in the midgut induced by beads is helping control viral infection but this would have to occur independently of their phagocytic activity.

Phagocytosis by Hemocytes Is Required for Systemic Control of DENV and ZIKV

The above results suggest that phagocytosis is not involved in the control of viral infection in the midgut of *Aedes* mosquitoes. This contrasts with the well-known roles of phagocytosis by hemocytes in insect immunity especially in the antiviral defense of *Drosophila*. However, these cells have also been shown to host replication of arboviruses such as DENV, Sindbis and O'nyong'nyong virus, which could help explain a proviral function (36–38). We confirmed that hemocytes could be directly infected by ZIKV as indicated staining for the viral E protein (Supplementary Figure 6). Thus, phagocytosis of viral particles by hemocytes could help promote viral replication in mosquitoes. In order to look further into this possibility, we analyzed dissemination of DENV and ZIKV infection from the midgut to the carcass in mosquitoes injected with latex beads (Figure 3A). Although the midgut infection rate was significantly reduced when phagocytosis was inhibited (Figures 2B–I), this did not significantly affect the prevalence of mosquitoes with disseminated infection (Figures 3B, C). Nevertheless, we observed a significant increase in viral RNA levels in the carcass of mosquitoes infected with DENV and ZIKV when phagocytosis by hemocytes was inhibited (Figures 3D, E). Here we note that mosquitoes fed on viremic mice show over 80% prevalence of infection. Therefore, to further analyze a possible effect of latex beads on the dissemination, we decided to analyze a model of artificial blood feeding where virus concentrations could be more easily controlled to reach closer to 50% prevalence (Figure 3F). In this model, injection of beads into mosquitoes prior to blood feeding containing DENV or ZIKV led to a significant decrease in the prevalence of infection (Figures 3G, H). At the same time, viral loads were not significantly different for DENV and were increased in ZIKV infected individuals when phagocytosis was inhibited (Figures 3I, J). This reinforces the idea that blocking phagocytosis by hemocytes leads to decreased midgut replication that results in lower systemic dissemination. In order to bypass the midgut and directly analyze the role of




FIGURE 1 | Hemocyte accumulation in the midgut of *Ae. aegypti* mosquitoes in response to DENV and ZIKV. **(A)** Mosquitoes fed with sugar, blood or blood and virus were dissected at different times and their midgut was analyzed by confocal microscopy. Virus-infected mice were used as a source of blood. **(B, C)** Quantification of the number of midgut-associated hemocytes between mosquitoes fed with sugar, blood or blood and virus at 4 **(B)** and 8 **(C)** days post feeding. DENV and ZIKV were analyzed together. 2 independent experiments for each virus were pooled. Each dot represents an individual midgut. Total number of midguts tested is indicated below each box plot. Upper, middle and lower bars in the boxplot represent the 75th percentile, the median and the 25th percentile, respectively. Statistical analyses were performed using the Kruskal-Wallis test followed by Dunn’s test to correct for multiple comparisons. ns, non-significant. **(D–G)** Representative confocal microscopy images of mosquito midguts showing CM-DiI stained hemocytes in magenta, DNA in yellow, viral E proteins in green and actin in blue. Midguts from mosquitoes fed with sugar **(D)**, blood **(E)**, blood + DENV **(F)** and blood + ZIKV **(G)** are shown at 8 days post feeding.




FIGURE 2 | Phagocytosis by hemocytes is not required to control DENV and ZIKV infection in the midgut of *Ae. aegypti* mosquitoes. **(A)** Mosquitoes injected with latex beads were fed 2 days later with blood + virus and dissected at different times to be analyzed. Virus-infected mice were used as a source of blood. **(B, C)** Percentage of total midgut infection area that shows staining for the viral protein at 4 and 8 days post infection was determined by immunofluorescence. Total number of midguts tested is indicated below each box plot. DENV **(B)** and ZIKV **(C)** were analyzed separately. 2 independent experiments for each virus were pooled. Each dot represents an individual midgut. **(D–G)** Representative confocal microscopy images of mosquito midguts showing CM-DiI stained hemocytes in magenta, DNA in yellow, viral E proteins in green and actin in blue. **(D, E)** Midguts from mosquitoes fed on blood + DENV. **(F, G)** Midguts from mosquitoes fed on blood + ZIKV. **(D, F)** Midguts from control mosquitoes injected with buffer; **(E, G)** Midguts from mosquitoes injected with latex beads. **(H)** DENV and **(I)** ZIKV RNA levels measured by RT-qPCR at 8 days post feeding. The number of positive midguts over the total tested is indicated below each boxplot. **(J, K)** Number of midgut-associated hemocytes in individual midguts from control and virus infected mosquitoes at 4 and 8 days post feeding. DENV **(J)** and ZIKV **(K)** were analyzed separately. Total number of midguts tested is indicated below each box plot. 2 independent experiments for each virus were pooled. **(B, C, H–K)** Each dot represents an individual midgut. Upper, middle and lower bars in the boxplot represent the 75th percentile, the median and the 25th percentile, respectively. Statistical analyses were performed using the Mann-Whitney-Wilcoxon test.




FIGURE 3 | Systemic dissemination of ZIKV and DENV infection is controlled by hemocyte phagocytosis. **(A)** Mosquitoes injected with latex beads were fed 2 days later with blood and virus and dissected at 8 days post feeding to be analyzed. Virus-infected mice were used as a source of blood. **(B, C)** Prevalence of DENV **(B)** and ZIKV **(C)** infection in the mosquito carcass at 8 days post feeding. The number of positive mosquitoes over the total tested is indicated below each column. One representative experiment is shown. This experiment was repeated 3 times for DENV and once for ZIKV. Statistical analyses were performed using two-tailed Fishers exact test. **(D, E)** Viral RNA levels at 8 days post feeding for DENV **(D)** and ZIKV **(E)**. One representative experiment is shown. This experiment was repeated 3 times for DENV and once for ZIKV. Each dot represents an individual mosquito. **(F)** Mosquitoes injected with latex beads were given an artificial blood meal with virus 2 days later and analyzed at 8 days post feeding. **(G, H)** Prevalence of DENV **(G)** and ZIKV **(H)** infection in mosquitoes injected with buffer or beads. The number of positive mosquitoes over the total tested is indicated below each column. One representative experiment is shown. This experiment was repeated twice for DENV and once for ZIKV. Statistical analyses were performed using two-tailed Fishers exact test. **(I, J)** DENV **(I)** and ZIKV **(J)** RNA levels in mosquitoes injected with buffer or beads. Each dot represents an individual mosquito. The number of positive mosquitoes over the total tested is indicated below each boxplot. One representative experiment is shown. This experiment was repeated twice for DENV and once for ZIKV. **(D, E, I, J)** Upper, middle and lower bars in the boxplot represent the 75th percentile, the median and the 25th percentile, respectively. Statistical analyses were performed using the Mann-Whitney-Wilcoxon test.


hemocytes during systemic viral replication, we used a model of intrathoracic injection of the virus (**Figure 4A**). Consistent with results using the oral infection models, we observed that inhibition of the phagocytic activity of hemocytes led to a clear increase in systemic viral replication after injection of DENV and ZIKV (**Figures 4B, C**). This effect was highly significant and did not depend on the dose of virus used or the kinetics of infection. Together, our data indicate that phagocytosis by hemocytes is essential to control systemic viral replication, which is consistent with their important roles in cellular immunity.

DISCUSSION

Here we have studied the role of phagocytosis by insect macrophage-like cells in the control of DENV and ZIKV in *Ae. aegypti* mosquitoes. These macrophage-like cells, known as hemocytes, are important components of the mosquito immune system (25). We and others have previously shown that phagocytosis by these cells plays an important function in the antiviral defense of *Drosophila* (20–22) but their role during viral infections in mosquitoes remain unclear.

Our results show that hemocytes accumulate in the midgut of *Ae. aegypti* mosquitoes in response to the presence of ZIKV and

DENV in the blood meal. Interestingly, increased numbers of hemocytes in the midgut are not observed at 4 days post infection but only later at 8 days, suggesting it either requires continuous stimulation or is triggered only after certain levels of viral replication. Since the infection did not significantly change the number of circulating hemocytes, these results suggest that these cells were recruited or retained more efficiently in the midgut. Increased numbers of hemocytes in the midgut suggests an important role for these cells in the response to viral infection. However, our results were less clear regarding their possible function in the midgut. We observed that blocking phagocytosis by hemocytes using latex beads led to decreased virus replication in the midgut after 8 days post infection when these cells accumulate significantly. We show that phagocytosis is inhibited at 2 days post injection of latex beads at the time of viral infection in the midgut. Although it is unclear how long this inhibition lasts, these results suggest that phagocytosis by hemocytes has a proviral function during the early stages of DENV and ZIKV infection in the midgut. However, when phagocytosis was blocked by latex beads, we also observed increased numbers of hemocytes in the midgut of mosquitoes as early as 4 days post infection. This effect was independent of viral replication or blood feeding and could be related to lower motility of hemocytes after phagocytosis since we do observe a



tendency of decreased numbers of circulating hemocytes after injection of beads. Nevertheless, increased numbers of hemocytes in the midgut precede and could be responsible for the inhibition of viral replication independent of phagocytosis. Phagocytosis of latex beads does not seem to prime hemocytes for wound healing in *Drosophila* (26), which would suggest that these hemocytes in the mosquito midgut are not activated but rather inert. Our current data do not allow us to rule out that there are other antiviral functions by hemocytes triggered by latex beads (e.g. production of antiviral cytokines) but it is clear that phagocytosis is not required to control viral infection in the midgut of mosquitoes. Paradoxically, when the infection disseminates from the midgut, phagocytosis by hemocytes has an important role controlling systemic viral replication. Together, our data point to a dual role for phagocytosis by hemocytes in the antiviral response of *Ae. aegypti* mosquitoes against DENV and ZIKV. Phagocytosis does not affect virus replication in the midgut but is essential to control systemic infection. Notably, work by our groups and others have pointed to important differences in the requirements to control viral replication in the midgut compared to systemic infection in mosquitoes (14, 36). For example, RNA interference plays a major role during systemic infection but has little contribution to the control of viral replication in the midgut (14). Similarly, apoptosis, which is coupled to phagocytosis by hemocytes to restrict viral infection in *Drosophila* (20, 21), may not be efficiently induced in response to virus infection in the midgut epithelium of *Aedes* mosquitoes.

The reason for contrasting roles of hemocytes during infection of the midgut compared to systemic dissemination of DENV and ZIKV suggests a complex scenario. It is possible that hemocytes might carry out immune functions that have opposing impacts over viral infection whether in the midgut or systemically. For example, phagocytosis might be important to clear viruses from the circulation but, in the midgut, could help virus dissemination. However, recent single cell analyses have indicated that hemocytes are composed of many subgroups that likely have distinct functions in immunity (43–46). Based on these data, it is possible that epithelial and systemic responses to viral infections mobilize different subtypes of hemocytes. Upon blood feeding, there is extensive damage to the basal lamina of the midgut and this is further exacerbated by infection by chikungunya virus (47, 48). ZIKV causes similar damage to the basal lamina (49) and that is likely true for other arboviruses. Damage to the basal lamina presumably leads to the recruitment of certain subtypes of hemocytes to these damaged regions with high concentration of the virus (47, 49). It is possible that hemocytes that are recruited to repair this damage become infected and help amplify local viral replication. Alternatively, these hemocytes could promote enterocyte survival or intestinal stem cell proliferation (50–53) and thus favor viral replication in the midgut. In contrast, during systemic dissemination of ZIKV and DENV, other subtypes of hemocytes would then play a more classical antiviral role by clearing particles and infected cells (20). Hemocytes may also participate in a systemic antiviral RNA interference of mosquitoes, as proposed in *Drosophila* (22), and

this may not be functional in the midgut. Notably, recent work in *Anopheles* mosquitoes has suggested that subtypes of hemocytes may have different roles during specific stages of *Plasmodium* infection (34). These are pressing questions that we are currently investigating to elucidate the mechanism by which hemocytes contribute to the antiviral defense. Alternative methods for hemocyte depletion (34, 54) or genetic approaches to ablate or interfere with cell function in mosquitoes will be important tools for the field going forward. These studies will help understand how vector mosquitoes recognize and fight viral infections that could lead to novel strategies to control transmission of arboviruses.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**. Further inquiries can be directed to the corresponding author.

ETHICS STATEMENT

The animal study was reviewed and approved by Comissão de Ética no uso de animais - CEUA da UFMG (337/2016).

AUTHOR CONTRIBUTIONS

TL, AF, J-LI, and JM designed the experiments. TL and AF carried out the experiments and statistical analysis. TL, AF, J-LI, and JTM analyzed results. TL, AF, J-LI, and JM wrote the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was funded by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Fundação de Amparo a Pesquisa do Estado de Minas Gerais (FAPEMIG), Rede Mineira de Imunobiológicos (grant # REDE-00140-16), Instituto Nacional de Ciência e Tecnologia de Vacinas (INCTV), the Institute for Advanced Studies of the University of Strasbourg (USIAS fellowship 2019) and the Investissement d'Avenir Programs (ANR-10-LABX-0036 and ANR-11-EQPX-0022). This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

ACKNOWLEDGMENTS

We would like to thank members of the Marques laboratory and research units CNRS UPR9022, Inserm U1257, in particular Dominique Ferrandon, for helpful discussions and suggestions.

Images in this work were obtained with the help of Gustavo Menezes and Hortência Oliveira from the Center for Gastrointestinal Biology at UFMG and also using equipment in the Centro de Aquisição e Processamento de Imagens from UFMG (CAPI- ICB/UFMG).

REFERENCES

- Shaw WR, Catteruccia F. Vector Biology Meets Disease Control: Using Basic Research to Fight Vector-Borne Diseases. *Nat Microbiol* (2019) 4:20–34. doi: 10.1038/s41564-018-0214-7
- Marques JT, Imler J-L. The Diversity of Insect Antiviral Immunity: Insights From Viruses. *Curr Opin Microbiol* (2016) 32:71–6. doi: 10.1016/j.mib.2016.05.002
- Deddouche S, Matt N, Budd A, Mueller S, Kemp C, Galiana-Arnoux D, et al. The DEXD/H-box Helicase Dicer-2 Mediates the Induction of Antiviral Activity in *Drosophila*. *Nat Immunol* (2008) 9:1425–32. doi: 10.1038/ni.1664
- Dostert C, Jouanguy E, Irving P, Troxler L, Galiana-Arnoux D, Hetru C, et al. The Jak-STAT Signaling Pathway is Required But Not Sufficient for the Antiviral Response of *Drosophila*. *Nat Immunol* (2005) 6:946–53. doi: 10.1038/ni1237
- Goto A, Okado K, Martins N, Cai H, Barbier V, Lamiabile O, et al. The Kinase Ikk β Regulates a STING- and NF- κ B-Dependent Antiviral Response Pathway in *Drosophila*. *Immunity* (2018) 49:225–34. doi: 10.1016/j.immuni.2018.07.013
- Marques JT, Wang J-P, Wang X, de Oliveira KPV, Gao C, Aguiar ERGR, et al. Functional Specialization of the Small Interfering RNA Pathway in Response to Virus Infection. *PLoS Pathog* (2013) 9:e1003579. doi: 10.1371/journal.ppat.1003579
- Cai H, Holleufer A, Simonsen B, Schneider J, Lemoine A, Gad HH, et al. 2'3'-cGAMP Triggers a STING- and NF- κ B-Dependent Broad Antiviral Response in *Drosophila*. *Sci Signal* (2020) 13:eabc4537. doi: 10.1126/scisignal.abc4537
- Kemp C, Mueller S, Goto A, Barbier V, Paro S, Bonnay F, et al. Broad RNA Interference-Mediated Antiviral Immunity and Virus-Specific Inducible Responses in *Drosophila*. *J Immunol* (2013) 190:650–8. doi: 10.4049/jimmunol.1102486
- West C, Silverman N. p38b and JAK-STAT Signaling Protect Against Invertebrate Iridescent Virus 6 Infection in *Drosophila*. *PLoS Pathog* (2018) 14:e1007020. doi: 10.1371/journal.ppat.1007020
- van Rij RP, Saleh M-C, Berry B, Foo C, Houk A, Antoniewski C, et al. The RNA Silencing Endonuclease Argonaute 2 Mediates Specific Antiviral Immunity in *Drosophila Melanogaster*. *Genes Dev* (2006) 20:2985–95. doi: 10.1101/gad.1482006
- Merkling SH, Bronkhorst AW, Kramer JM, Overheul GJ, Schenck A, Van Rij RP. The Epigenetic Regulator G9a Mediates Tolerance to RNA Virus Infection in *Drosophila*. *PLoS Pathog* (2015) 11:e1004692. doi: 10.1371/journal.ppat.1004692
- Wang X-H, Aliyari R, Li W-X, Li H-W, Kim K, Carthew R, et al. RNA Interference Directs Innate Immunity Against Viruses in Adult *Drosophila*. *Science* (2006) 312:452–4. doi: 10.1126/science.1125694
- Souza-Neto JA, Sim S, Dimopoulos G. An Evolutionary Conserved Function of the JAK-STAT Pathway in Anti-Dengue Defense. *Proc Natl Acad Sci U S A* (2009) 106:17841–6. doi: 10.1073/pnas.0905006106
- Olmo RP, Ferreira AGA, Izidoro-Toledo TC, Aguiar ERGR, de Faria IJS, de Souza KPR, et al. Control of Dengue Virus in the Midgut of *Aedes Aegypti* by Ectopic Expression of the dsRNA-binding Protein Loqs2. *Nat Microbiol* (2018) 3:1385–93. doi: 10.1038/s41564-018-0268-6
- Sánchez-Vargas I, Scott JC, Poole-Smith BK, Franz AWE, Barbosa-Solomieu V, Wilusz J, et al. Dengue Virus Type 2 Infections of *Aedes Aegypti* Are Modulated by the Mosquito's RNA Interference Pathway. *PLoS Pathog* (2009) 5:e1000299. doi: 10.1371/journal.ppat.1000299
- Morazzani EM, Wiley MR, Murreddu MG, Adelman ZN, Myles KM. Production of Virus-Derived Ping-Pong-Dependent piRNA-like Small RNAs in the Mosquito Soma. *PLoS Pathog* (2012) 8:e1002470. doi: 10.1371/journal.ppat.1002470
- Khoo CC, Piper J, Sanchez-Vargas I, Olson KE, Franz AW. The RNA Interference Pathway Affects Midgut Infection- and Escape Barriers for Sindbis Virus in *Aedes Aegypti*. *BMC Microbiol* (2010) 10:130. doi: 10.1186/1471-2180-10-130
- Khoo CCH, Doty JB, Heersink MS, Olson KE, Franz AWE. Transgene-Mediated Suppression of the RNA Interference Pathway in *Aedes Aegypti* Interferes With Gene Silencing and Enhances Sindbis Virus and Dengue Virus Type 2 Replication. *Insect Mol Biol* (2013) 22:104–14. doi: 10.1111/imb.12008
- Wu X, Wu F-H, Wang X, Wang L, Siedow JN, Zhang W, et al. Molecular Evolutionary and Structural Analysis of the Cytosolic DNA Sensor cGAS and STING. *Nucleic Acids Res* (2014) 42:8243–57. doi: 10.1093/nar/gku569
- Lamiabile O, Arnold J, de Faria IJ da S, Olmo RP, Bergami F, Meignin C, et al. Analysis of the Contribution of Hemocytes and Autophagy to *Drosophila* Antiviral Immunity. *J Virol* (2016) 90:5415–26. doi: 10.1128/JVI.00238-16
- Nainu F, Tanaka Y, Shiratsuchi A, Nakanishi Y. Protection of Insects Against Viral Infection by Apoptosis-Dependent Phagocytosis. *J Immunol* (2015) 195:5696–706. doi: 10.4049/jimmunol.1500613
- Tassetto M, Kunitomi M, Andino R. Circulating Immune Cells Mediate a Systemic RNAi-Based Adaptive Antiviral Response in *Drosophila*. *Cell* (2017) 169:314–25.e13. doi: 10.1016/j.cell.2017.03.033
- Hillyer J. The Antibacterial Innate Immune Response by the Mosquito *Aedes Aegypti* is Mediated by Hemocytes and Independent of Gram Type and Pathogenicity. *Microbes Infect* (2004) 6:448–59. doi: 10.1016/j.micinf.2004.01.005
- Hillyer JF, Schmidt SL, Christensen BM. Rapid Phagocytosis and Melanization of Bacteria and *Plasmodium* Sporozoites by Hemocytes of the Mosquito *Aedes Aegypti*. *J Parasitol* (2003) 89:62–9. doi: 10.1645/0022-3395
- Hillyer JF, Strand MR. Mosquito Hemocyte-Mediated Immune Responses. *Curr Opin Insect Sci* (2014) 3:14–21. doi: 10.1016/j.cois.2014.07.002
- Weavers H, Evans IR, Martin P, Wood W. Corpse Engulfment Generates a Molecular Memory That Primes the Macrophage Inflammatory Response. *Cell* (2016) 165:1658–71. doi: 10.1016/j.cell.2016.04.049
- Sanchez Bosch P, Makhijani K, Herboso L, Gold KS, Baginsky R, Woodcock KJ, et al. Adult *Drosophila* Lack Hematopoiesis But Rely on a Blood Cell Reservoir At the Respiratory Epithelia to Relay Infection Signals to Surrounding Tissues. *Dev Cell* (2019) 51:787–803.e5. doi: 10.1016/j.devcel.2019.10.017
- King JG, Hillyer JF. Spatial and Temporal In Vivo Analysis of Circulating and Sessile Immune Cells in Mosquitoes: Hemocyte Mitosis Following Infection. *BMC Biol* (2013) 11:55. doi: 10.1186/1741-7007-11-55
- King JG, Hillyer JF. Infection-Induced Interaction Between the Mosquito Circulatory and Immune Systems. *PLoS Pathog* (2012) 8:e1003058. doi: 10.1371/journal.ppat.1003058
- Barletta ABF, Trisnadi N, Ramirez JL, Barillas-Mury C. Mosquito Midgut Prostaglandin Release Establishes Systemic Immune Priming. *iScience* (2019) 19:54–62. doi: 10.1016/j.isci.2019.07.012
- Rodríguez J, Brayner FA, Alves LC, Dixit R, Barillas-Mury C. Hemocyte Differentiation Mediates Innate Immune Memory in *Anopheles gambiae* Mosquitoes. *Science* (2010) 329:1353–5. doi: 10.1126/science.1190689
- Volohonsky G, Paul-Gilloteaux P, Štáfková J, Soichot J, Salamero J, Levashina EA. Kinetics of *Plasmodium* Midgut Invasion in *Anopheles* Mosquitoes. *PLoS Pathog* (2020) 16:e1008739. doi: 10.1371/journal.ppat.1008739
- Castillo JC, Ferreira ABB, Trisnadi N, Barillas-Mury C. Activation of Mosquito Complement Antiplasmodial Response Requires Cellular Immunity. *Sci Immunol* (2017) 2:eaa11505. doi: 10.1126/sciimmunol.aal1505
- Kwon H, Smith RC. Chemical Depletion of Phagocytic Immune Cells in *Anopheles Gambiae* Reveals Dual Roles of Mosquito Hemocytes in anti-*Plasmodium* Immunity. *Proc Natl Acad Sci U S A* (2019) 116:14119–28. doi: 10.1073/pnas.1900147116
- Franz A, Kantor A, Passarelli A, Clem R. Tissue Barriers to Arbovirus Infection in Mosquitoes. *Viruses* (2015) 7:3741–67. doi: 10.3390/v7072795
- Carissimo G, Pondeville E, McFarlane M, Dietrich I, Mitri C, Bischoff E, et al. Antiviral Immunity of *Anopheles Gambiae* is Highly Compartmentalized,

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2021.660873/full#supplementary-material>

- With Distinct Roles for RNA Interference and Gut Microbiota. *Proc Natl Acad Sci U S A* (2015) 112:E176–85. doi: 10.1073/pnas.1412984112
37. Salazar MI, Richardson JH, Sánchez-Vargas I, Olson KE, Beaty BJ. Dengue Virus Type 2 Replication and Tropisms in Orally Infected *Aedes Aegypti* Mosquitoes. *BMC Microbiol* (2007) 7:9. doi: 10.1186/1471-2180-7-9
 38. Parikh GR, Oliver JD, Bartholomay LC. A Hemocyte Tropism for an Arbovirus. *J Gen Virol* (2009) 90:292–6. doi: 10.1099/vir.0.005116-0
 39. Coggins SA, Estévez-Lao TY, Hillyer JF. Increased Survivorship Following Bacterial Infection by the Mosquito *Aedes Aegypti* as Compared to *Anopheles Gambiae* Correlates With Increased Transcriptional Induction of Antimicrobial Peptides. *Dev Comp Immunol* (2012) 37:390–401. doi: 10.1016/j.dci.2012.01.005
 40. Castillo J, Brown MR, Strand MR. Blood Feeding and Insulin-like Peptide 3 Stimulate Proliferation of Hemocytes in the Mosquito *Aedes Aegypti*. *PLoS Pathog* (2011) 7:e1002274. doi: 10.1371/journal.ppat.1002274
 41. Bryant WB, Michel K. Blood Feeding Induces Hemocyte Proliferation and Activation in the African Malaria Mosquito, *Anopheles Gambiae* Giles. *J Exp Biol* (2014) 217:1238–45. doi: 10.1242/jeb.094573
 42. Nehme NT, Liégeois S, Kele B, Giammarinaro P, Pradel E, Hoffmann JA, et al. A Model of Bacterial Intestinal Infections in *Drosophila Melanogaster*. *PLoS Pathog* (2007) 3:e173. doi: 10.1371/journal.ppat.0030173
 43. Raddi G, Barletta ABF, Efremova M, Ramirez JL, Cantera R, Teichmann SA, et al. Mosquito Cellular Immunity At Single-Cell Resolution. *Science* (2020) 369:1128–32. doi: 10.1126/science.abc0322
 44. Severo MS, Landry JJM, Lindquist RL, Goosmann C, Brinkmann V, Collier P, et al. Unbiased Classification of Mosquito Blood Cells by Single-Cell Genomics and High-Content Imaging. *Proc Natl Acad Sci U S A* (2018) 115:E7568–77. doi: 10.1073/pnas.1803062115
 45. Cattenoz PB, Sakr R, Pavlidaki A, Delaporte C, Riba A, Molina N, et al. Temporal Specificity and Heterogeneity of *Drosophila* Immune Cells. *EMBO J* (2020) 39:e104486. doi: 10.15252/embj.2020104486
 46. Tattikota SG, Cho B, Liu Y, Hu Y, Barrera V, Steinbaugh MJ, et al. A Single-Cell Survey of *Drosophila* Blood. *Elife* (2020) 9:e54818. doi: 10.7554/eLife.54818
 47. Dong S, Balaraman V, Kantor AM, Lin J, Grant DG, Held NL, et al. Chikungunya Virus Dissemination From the Midgut of *Aedes Aegypti* is Associated With Temporal Basal Lamina Degradation During Bloodmeal Digestion. *PLoS Negl Trop Dis* (2017) 11:e0005976. doi: 10.1371/journal.pntd.0005976
 48. Kantor AM, Grant DG, Balaraman V, White TA, Franz AWE. Ultrastructural Analysis of Chikungunya Virus Dissemination From the Midgut of the Yellow Fever Mosquito, *Aedes Aegypti*. *Viruses* (2018) 10:571. doi: 10.3390/v10100571
 49. Cui Y, Grant DG, Lin J, Yu X, Franz AWE. Zika Virus Dissemination From the Midgut of *Aedes Aegypti* is Facilitated by Bloodmeal-Mediated Structural Modification of the Midgut Basal Lamina. *Viruses* (2019) 11:1056. doi: 10.3390/v11111056
 50. Amcheslavsky A, Lindblad JL, Bergmann A. Transiently “Undead” Enterocytes Mediate Homeostatic Tissue Turnover in the Adult *Drosophila* Midgut. *Cell Rep* (2020) 33:108408. doi: 10.1016/j.celrep.2020.108408
 51. Haller S, Franchet A, Hakkim A, Chen J, Drenkard E, Yu S, et al. Quorum-Sensing Regulator RhlR But Not its Autoinducer RhlI Enables *Pseudomonas* to Evade Opsonization. *EMBO Rep* (2018) 19:e44880. doi: 10.15252/embr.201744880
 52. Ayyaz A, Li H, Jasper H. Hemocytes Control Stem Cell Activity in the *Drosophila* Intestine. *Nat Cell Biol* (2015) 17:736–48. doi: 10.1038/ncb3174
 53. Chakrabarti S, Dudzic JP, Li X, Collas EJ, Boquete J-P, Lemaitre B. Remote Control of Intestinal Stem Cell Activity by Hemocytes in *Drosophila*. *PLoS Genet* (2016) 12:e1006089. doi: 10.1371/journal.pgen.1006089
 54. Ramesh Kumar J, Smith JP, Kwon H, Smith RC. Use of Clodronate Liposomes to Deplete Phagocytic Immune Cells in *Drosophila Melanogaster* and *Aedes Aegypti*. *Front Cell Dev Biol* (2021) 9:627976. doi: 10.3389/fcell.2021.627976

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Leite, Ferreira, Imler and Marques. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.