

# *Moringa oleifera* extract promotes apoptosis-like death in *Toxoplasma gondii* tachyzoites *in vitro*

## Research Article

**Cite this article:** Nishi L *et al* (2021). *Moringa oleifera* extract promotes apoptosis-like death in *Toxoplasma gondii* tachyzoites *in vitro*. *Parasitology* **148**, 1447–1457. <https://doi.org/10.1017/S0031182021001086>

Received: 12 March 2021  
Revised: 29 April 2021  
Accepted: 15 June 2021  
First published online: 30 June 2021

### Key words:

HeLa cells; mechanism of action; plant extract; toxoplasmosis

### Author for correspondence:

Ana Lúcia Falavigna-Guilherme,  
E-mail: [alfguilherme@uem.br](mailto:alfguilherme@uem.br)

Letícia Nishi<sup>1</sup> , Raquel Arruda da Silva Sanfelice<sup>2</sup> ,  
Bruna Taciane da Silva Bortoleti<sup>2,3</sup>, Fernanda Tomiotto-Pellissier<sup>2,3</sup> ,  
Taylon Felipe Silva<sup>2</sup>, Fernanda Ferreira Evangelista<sup>1</sup> , Danielle Lazarin-Bidóia<sup>2</sup>,  
Idessania Nazareth Costa<sup>2</sup> , Wander Rogério Pavanelli<sup>2</sup>, Ivete Conchon Costa<sup>2</sup>,  
Aline Takaoka Alves Baptista<sup>4</sup>, Rosângela Bergamasco<sup>5</sup>  
and Ana Lúcia Falavigna-Guilherme<sup>1</sup>

<sup>1</sup>Graduate Program in Health Science, State University of Maringá, Colombo Avenue, 5790, Zip Code 87020-900, Maringá, Paraná, Brazil; <sup>2</sup>Department of Pathological Sciences, Laboratory of Immunoparasitology of Neglected Diseases and Cancer – LIDNC, Center of Biological Sciences, State University of Londrina, Rodovia Celso Garcia Cid Campus, Zip Code 86057-970, Postal box 10.011, Londrina, Paraná, Brazil; <sup>3</sup>Biosciences and Biotechnology Postgraduate Program, Carlos Chagas Institute (ICC/Fiocruz-PR), Curitiba, Paraná, Brazil; <sup>4</sup>Departament of Food and Chemical Engineering, Federal University of Technology – Paraná – UTFPR, Câmpus Campo Mourão, Via Rosalina Maria Dos Santos, 1233, Zip Code 87301-899, Campo Mourão, Paraná, Brazil and <sup>5</sup>Department of Chemical Engineering, State University of Maringá, Colombo Avenue, 5790, Zip Code 87020-900, Maringá, Paraná, Brazil

### Abstract

*Toxoplasma gondii* is the causative agent of toxoplasmosis, and an important problem of public health. The current treatment for toxoplasmosis is the combination of pyrimethamine and sulphadiazine, which do not act in the chronic phase of toxoplasmosis and have several side-effects. This study evaluated the anti-*T. gondii* activity and potential mechanism of *Moringa oleifera* seeds' aqueous extract *in vitro*. The concentration of *M. oleifera* extract in HeLa cells was determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide cell viability assays. The presence of *T. gondii* was assessed by quantitative polymerase chain reaction and toluidine blue staining. Pyrimethamine and sulphadiazine were used as drug controls. Modifications in *T. gondii* morphology and ultrastructure were observed by electron microscopy. *In vitro*, the *M. oleifera* extract had no toxic effect on HeLa cells at concentrations below 50 µg mL<sup>-1</sup>. *Moringa oleifera* extract inhibits *T. gondii* invasion and intracellular proliferation with similar results for sulphadiazine + pyrimethamine, and also shows cellular nitric oxide production at a concentration of 30 µg mL<sup>-1</sup>. Electron microscopy analyses indicated structural and ultrastructural modifications in tachyzoites after treatment. We also observed an increase in reactive oxygen species production and a loss of mitochondrial membrane integrity. Nile Red staining assays demonstrated a lipid accumulation. Annexin V–fluorescein isothiocyanate and propidium iodide staining demonstrated that the main action of *M. oleifera* extract in *T. gondii* tachyzoites was compatible with late apoptosis. In conclusion, *M. oleifera* extract has anti-*T. gondii* activity *in vitro* and might be a promising substance for the development of a new anti-*T. gondii* drug.

### Introduction

*Toxoplasma gondii* is an obligate intracellular protozoan parasite that infects a wide range of host mammals and birds, including humans (Su e Dubey, 2020). Toxoplasmosis affects one-third of the world population; in most cases, the disease is asymptomatic, but immunocompromised individuals may have severe symptoms, such as toxoplasmic encephalitis. Vertical transmission during pregnancy may lead to abortion, or the birth of children with mental retardation, hydrocephalus, psychomotor impairment and chorioretinitis, among others (Robert-Gangneux *et al.*, 2018).

Pyrimethamine associated with sulphadiazine has been the treatment available and indicated for severe cases for a long period (Silva *et al.*, 2019). Both drugs inhibit folate metabolism in the parasite, acting directly on tachyzoites (acute phase of infection); however, this does not eradicate cystic forms (chronic phase of infection) (Konstantinovic *et al.*, 2019). The initial treatment of choice must be carried out for several days (up to months) and the medication can cause several side-effects such as haematological abnormalities, increased creatinine and serum liver enzymes and hypersensitivity reactions (Konstantinovic *et al.*, 2019).

The investigation of new substances that are capable of eliminating *T. gondii* tachyzoites and cysts is important for reducing side-effects and serious outcomes, and possibly curing toxoplasmosis (Zwicker *et al.*, 2020). Research with compounds (extracts and molecules) from medicinal plants, which have therapeutic potential, have been used as alternative

treatments for some parasitic diseases; this is because, in addition to presenting microbicidal effects, they also confer lower toxicity compared to synthetic drugs (Mirzaalizadeh *et al.*, 2018).

*Moringa oleifera* is a plant belonging to the Moringaceae family; studies have shown that parts of the plant contained compounds with bactericidal, fungicidal, anti-tumour, larvicidal and trypanosomicidal properties (Dhakad *et al.*, 2019; Turner *et al.*, 2019; Arruda *et al.*, 2020; Coriolano *et al.*, 2020; Pagar *et al.*, 2020). In addition, *M. oleifera* has activity against the species of the protozoan *Plasmodium* (Somsak *et al.*, 2016) which, as *T. gondii*, belongs to the phylum Apicomplexa.

Considering the properties of *M. oleifera* and the fact that, in relation to *T. gondii*, there have been no studies on this plant, the aim of the current research was to evaluate the activity of *M. oleifera* seed extract on HeLa cells infected with *T. gondii* tachyzoites (RH strain) and directly on tachyzoites forms of *T. gondii*.

## Materials and methods

### HeLa cell culture maintenance

The cultivation of HeLa cells was performed in 75 cm<sup>2</sup> culture flasks (Ciencor Scientific, Brasil) containing Dulbecco's modified Eagle medium (DMEM) (Gibco, Invitrogen, New York, USA) supplemented with 10% inactivated fetal bovine serum (Sigma-Aldrich, St. Louis, Missouri, EUA), 1% antibiotic (10 000 U mL<sup>-1</sup> of penicillin and 10 mg mL<sup>-1</sup> streptomycin) (Cultilab, Brazil), L-glutamine, sodium pyruvate and 2β mercaptoethanol. The cells were maintained at 37°C with 5% CO<sub>2</sub>. For the tests of experimental infection and the maintenance of *T. gondii* strains *in vitro*, HeLa cells were used up to the 21st passage.

### Toxoplasma gondii RH strain

Tachyzoites from the *T. gondii* RH strain were kindly provided by Professor João Luís Garcia from the Laboratory of Zoonoses and Public Health at the State University of Londrina, where tachyzoites were obtained from the peritoneal lavage of previously infected Swiss Webster mice. The peritoneal exudates were passed through a 26G needle three times, and washed twice with phosphate-buffered saline (PBS) (pH 7.5) by centrifugation. The sediment was resuspended in PBS and the tachyzoites were counted in a Neubauer chamber.

Part of the tachyzoites was kept in HeLa cells for 'Anti-*T. gondii* activity assay *in vitro*' (intracellular tachyzoites) and another part was used to evaluate the direct action of *M. oleifera* extract on *T. gondii* (extracellular) tachyzoites.

All procedures involving animals in this study were approved by the Animal Experimentation Ethics Committee of the State University of Londrina no. 88/2017/CEUA; 82.862.016.60.

### Moringa oleifera seeds extract

*Moringa oleifera* seeds were collected between December 2018 and March 2019 in the area of Federal University of Sergipe, Aracaju, Sergipe State, located between the geographic coordinates 10°55'56" south latitude and 37°04'23" west longitude. Aracaju has a semiarid climate with low pluviometric index (Almeida *et al.*, 2017). The *M. oleifera* species was deposited in the herbarium of Federal University of Sergipe, no. ASE8288 (MOTA *et al.*, 2012).

For the preparation of aqueous extracts, the seeds were peeled, weighed (5 g) and processed by turbolysis for 3 min using 100 mL distilled water. Then, the solution was subjected to magnetic stirring (IKA, Germany) for 30 min, and, after this process, subjected to two consecutive filtrations, using qualitative filter paper and a

0.9 μm glass fibre membrane by vacuum filtration (Madrona *et al.*, 2010). The aqueous extract obtained was lyophilized (Freeze Dryer Christ ALPHA 1-2/LD Plus) for 48 h at -45°C (Baptista *et al.*, 2017). The material obtained was stored at 4°C until the moment of use. For experiments, the lyophilized material was diluted in DMEM at different concentrations and the solution obtained was called *M. oleifera* extract.

The extract was characterized by Fourier transform infrared spectroscopy (FTIR) in the form of potassium bromide (KBr) tablets containing approximately 1% of the sample using an FTIR-BOMEN 100 spectrometer with 21 scans per min and a resolution of 4 cm<sup>-1</sup>. The spectra of FTIR were obtained in the range from 4000 to 400 cm<sup>-1</sup> in absorbance mode.

### HeLa cell viability by MTT assay

The viability of HeLa cells after treatment with *M. oleifera* was evaluated based on mitochondrial oxidation, using the tetrazolium salt colorimetric test [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide or MTT] (Sigma-Aldrich) (Mosmann, 1983). The cells were grown in 96-well plates (3 × 10<sup>4</sup> cells per well per 200 μL), for 24 h in DMEM at 37°C and 5% CO<sub>2</sub>. After this period, *M. oleifera* extract was added separately at concentrations of 10, 30, 50, 100 and 200 μg mL<sup>-1</sup> and maintained for 24 h under the same conditions stated before. As a negative control, cells cultured with DMEM were used, and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) 0.06% was used as a positive control. Then, the supernatants were removed and the cells received MTT solution (5 mg mL<sup>-1</sup>) and incubated for 3 h at 37°C and 5% CO<sub>2</sub>. The formazan crystals were diluted with 100 μL of dimethyl sulphoxide (Sigma-Aldrich), and the absorbance was measured after 30 min at 570 nm in a microplate spectrophotometer (Thermo Plate – TP-Reader). The results were expressed as a relative percentage of MTT reduction in the treated groups compared to the control group, using the following formula:

$$\% \text{Viable cells} = \left( \frac{\text{OD treated cells}}{\text{OD untreated cells}} \right) \times 100$$

where OD is the optical density.

### Anti-T. gondii activity assay in vitro

To evaluate the effect of the *M. oleifera* extract on the invasion and proliferation of *T. gondii* in HeLa cells (1 × 10<sup>5</sup>), 24-well plates were used containing round coverslips of 13 mm (Ciencor Scientific, Brazil), and 5 × 10<sup>5</sup> *T. gondii* tachyzoites (1:5 cells/tachyzoites) were added to each one. After 3 h of infection, the supernatant was removed and the cells were washed once with PBS, the plate was manually and slightly shaken and the PBS was discarded. Then, the cells with *T. gondii* tachyzoites were treated with the *M. oleifera* extract, at concentrations of 10 and 30 μg mL<sup>-1</sup>. The negative controls were untreated infected cells (DMEM only), and infected cells treated with a combination of sulphadiazine and pyrimethamine (SDZ + PYR) (50 and 25 μg mL<sup>-1</sup>) were used as positive controls. They were maintained for 24 h at 37°C in 5% CO<sub>2</sub>. Afterwards, the supernatant was separated to quantify nitric oxide (NO) and the cells were fixed with 4% paraformaldehyde in PBS for 30 min and stained with 1% toluidine blue (Sigma-Aldrich).

The HeLa cells were analysed by light microscopy (e100, Nikon – led) at 1000× magnification to check the parameters of infection index (number of infected cells per 200 cells examined) and intracellular proliferation of the parasite (total number of parasites per 200 cells examined). Representative images were

captured using the conditions described above. The percentages of infection and intracellular proliferation inhibition rates of *T. gondii* were calculated as: average of the infection or intracellular proliferation rate analysed in untreated cells, corresponding to 100% infection or intracellular proliferation rate. The percentages of inhibition of these parameters submitted to treatments with *M. oleifera* were calculated by subtracting the percentage values obtained for treated cells from those obtained for untreated cells (Sanfelice *et al.*, 2017).

#### Quantification of NO produced by HeLa cells infected by *T. gondii* tachyzoites

The supernatants in the item above were separated and used to quantify NO, as described by Tomiotto-Pellissier *et al.* (2018). Briefly, the NO was determined through the Griess method in culture supernatant aliquots (60  $\mu$ L) of *T. gondii* invasion and proliferation assay. First of all, the aliquots were centrifuged at  $2370 \times g$  for 2 min. A 50  $\mu$ L aliquot of the supernatant was treated with 50  $\mu$ L of Griess reagent [1% sulphanilamide and 0.1% of *N*-(1-naphthyl) ethylenediamine in orthophosphoric acid ( $H_3PO_4$ ) 5%]. After a 10-min incubation at room temperature, the samples were placed in 96-well microplates. A calibration curve was made using dilutions of  $NaNO_2$ , and the absorbance was determined at 550 nm on a microplate reader (Thermo Scientific, Multiskan GO).

#### Quantification of parasitic load by qPCR

A second 24-well plate was prepared in the same way as in the Section 'Anti-*T. gondii* activity assay *in vitro*' to evaluate the reduction of intracellular tachyzoites observed in light microscopy for the *M. oleifera* extract of  $30 \mu g mL^{-1}$ . The prepared plate was used to assess the parasitic load of *T. gondii* contained in HeLa cells using quantitative real-time polymerase chain reaction (qPCR). DNA extraction was performed using a commercial kit (PureLink™ genomic DNA Mini Kit, Thermo Fisher Scientific®, EUA) following the manufacturer's recommendations. DNA concentration was measured by spectrophotometry (NanoDrop, Thermo Fisher Scientific®, EUA) and qPCR was performed in duplicate with 50 ng of DNA using the TaqMan system, with a standard curve with concentrations from  $6.3 \times 10^7$  to  $6.3 \times 10^3$  parasites per mL. Primers Tox-9F (5'-AGGAGAGATATCAGGA CTGTAG-3') and Tox11-R (5'-GCGTCGTCTCGTCTAGATC G-3') and probe Tox-TP1 (5'-CCGGCTTGGCTGCTTTTCC T-3') were used to amplify a 115 bp fragment from the repetitive 529 bp region of *T. gondii*, as previously described by Opsteegh *et al.* (2010). Each reaction consisted of 1 $\times$  master mix (Taqman™ Universal PCR Master Mix, Thermo Fisher Scientific®, EUA), 0.7  $\mu M$  of each primer (Tox-9F and Tox-11R), 0.1  $\mu M$  of probe (Tox-TP-1) and sterile ultrapure water, making the volume up to 20  $\mu$ L. Amplification was performed on the StepOne™ Plus Real Time PCR System (Thermo Fisher Scientific®, EUA). The thermocycling conditions used were: initial incubation of 95°C for 10 min, denaturation cycles 95°C for 10 s, annealing 58°C for 20 s and extension 72°C for 20 s, followed by a final stage of 40°C for 5 s. The fluorescence signal at 530 nm (Tox-TP1) was measured at the end of each extension step. Sterile ultrapure water without DNA was used as a negative control and was included in all reactions; HeLa cells infected with *T. gondii* tachyzoites without addition of the *M. oleifera* extract were used as a negative control of treatment. All samples were analysed in duplicate. The results were visualized and analysed using the program StepOne™ Software v.2.3 (Thermo Fisher Scientific®, EUA).

#### Scanning and transmission electron microscopy analyses

For scanning electron microscopy (SEM), *T. gondii* tachyzoites obtained from the peritoneal lavage of previously infected Swiss Webster mice were treated as described in the Section 'Toxoplasma gondii RH strain' and counted in the Neubauer chamber. Next, the tachyzoites ( $1.5 \times 10^6$ ) were treated with the *M. oleifera* extract ( $30 \mu g mL^{-1}$ ), for 1 h at 37°C. Then, they were centrifuged and washed once with PBS by pipetting up and down and centrifuged again. Next, the PBS was discarded and the tachyzoites were fixed with 2.5% glutaraldehyde in 0.1 M sodium cacodylate buffer, containing 1 mM  $CaCl_2$ . The tachyzoites were fixed on a glass support that was covered with poly-L-lysine, and dehydrated in an ascending series of ethanol (30–100%), before being submitted to critical-point dried with  $CO_2$ , coated with gold and observed on a high-resolution double beam electron microscope FEI SCIOS.


For transmission electron microscopy (TEM), *T. gondii* tachyzoites were treated in the same way as described for SEM. After incubation, the parasites were washed with PBS and fixed with 2.5% glutaraldehyde in 0.1 M sodium cacodylate buffer and post-fixed in a solution of 1%  $OsO_4$ , 0.8% potassium ferrocyanide and 10.0 mM  $CaCl_2$  in 0.10 M cacodylate buffer, at room temperature and protected from light. Then, the tachyzoites were washed again with 0.1 M sodium cacodylate buffer, dehydrated in an increasing acetone gradient (30–100%) and embedded in EPON resin and polymerized in an incubator at 60°C for 72 h. Ultrathin sections were obtained, deposited on a copper grid, and stained with uranyl acetate and lead citrate, for 20 and 10 min, respectively. Ultrastructural alterations were observed using a JEOL JEM 1400 transmission electron microscope.

#### Determination of reactive oxygen species (ROS) generation on *T. gondii* tachyzoites

ROS generation was evaluated in *T. gondii* tachyzoites obtained from the peritoneal lavage of previously infected Swiss Webster mice were treated as described above and counted in the Neubauer chamber. The tachyzoites ( $1.5 \times 10^6$ ) were subjected to treatment with the *M. oleifera* extract ( $30 \mu g mL^{-1}$ ) for 1 h at 37°C in 5%  $CO_2$  and analysed by probe  $H_2DCFDA$  (Bortoleti *et al.*, 2019). Tachyzoites in DMEM were used as a negative control, and tachyzoites in  $H_2O_2$  solution (0.4%) were used as a positive control. The micrographs were produced using the EVOS® Microscope FL Auto Cell Imaging System (Thermo Fisher) with a magnification of 200 $\times$ .

#### Determination of the mitochondrial membrane potential in *T. gondii* tachyzoites

For determination of the mitochondrial membrane potential of *T. gondii* tachyzoites (RH strain), we conducted tetramethylrhodamine ethyl ester (TMRE) staining (Sigma-Aldrich) (Tomiotto-Pellissier *et al.*, 2018). Briefly, parasites ( $1.5 \times 10^6$ ), obtained in the same way as described for ROS, were treated with the *M. oleifera* extract ( $30 \mu g mL^{-1}$ ), incubated for 1 h in an incubator at 37°C in 5%  $CO_2$ , and then washed once with PBS by pipetting up and down and centrifuged. Next, the PBS was discarded and the tachyzoites were incubated with 25 nM TMRE for 30 min at 37°C. Again, they were washed with PBS and analysed immediately in a fluorescence microplate reader (Victor X3, Perkin-Elmer) using the excitation and emission wavelengths of 480 and 580 nm, respectively. Carbonyl cyanide 3-chlorophenylhydrazone (CCCP), 100  $\mu M$ , was used as a positive control. The micrographs were obtained using the EVOS®



**Fig. 1.** FTIR spectra of *Moringa oleifera* extract.

Microscope FL Auto Cell Imaging System (Thermo Fisher) with a magnification of 200 $\times$ .

#### Lipid staining and fluorescence analysis in *T. gondii* tachyzoites

In this experiment, the concentration of  $1.5 \times 10^6$  *T. gondii* tachyzoites, obtained in the same way as described above, was treated with the *M. oleifera* extract ( $30 \mu\text{g mL}^{-1}$ ) and incubated for 1 h at 37°C. Then, the tachyzoites were collected, washed twice with PBS and left to incubate with  $10 \mu\text{g mL}^{-1}$  Nile Red (Sigma-Aldrich) for 30 min at 37°C in 5% CO<sub>2</sub>. PBS was used as a positive control. The cytoplasmic lipid bodies of the parasites were detected in a spectrofluorometer (Victor X3, Perkin-Elmer) at the excitation and emission wavelengths of 530 and 635 nm, respectively.

#### Evaluation of autophagic vacuoles

*Toxoplasma gondii* tachyzoites ( $1.5 \times 10^6$ ) obtained in the same way as described above were treated with *M. oleifera* extract ( $30 \mu\text{g mL}^{-1}$ ) for 1 h at 37°C in 5% CO<sub>2</sub>. Then, the autophagic vacuole formation was evaluated using monodansylcadaverine (MDC)  $50 \mu\text{M}$  labelling, and incubated for 1 h at 25°C in a dark room, according to Bortoleti *et al.* (2021). After this period, the tachyzoites were washed with PBS and analysed on the spectrofluorimeter (Victor X3, Perkin-Elmer), using excitation and emission wavelengths of 380 and 525 nm, respectively. Tachyzoites in PBS solution were used as a positive control (Bortoleti *et al.*, 2021).

#### Co-determination of annexin-V-fluorescein isothiocyanate (FITC) and propidium iodide (PI) in tachyzoites of *T. gondii*

*Toxoplasma gondii* tachyzoites ( $1.5 \times 10^6$ ) obtained in the same way as described above were treated with the *M. oleifera* extract ( $30 \mu\text{g mL}^{-1}$ ), for 1 h at 37°C in 5% CO<sub>2</sub>. Then, they were washed with PBS, and resuspended in 100  $\mu\text{L}$  of binding buffer (Santa Cruz Biotechnology) with the addition of a mixture of 1  $\mu\text{L}$  of annexin-V-FITC and 1  $\mu\text{L}$  of PI (Santa Cruz Biotechnology). The analysis was performed by flow cytometry using the BD Accuri C6 (BD Biosciences, San Jose, CA) (Bortoleti *et al.*, 2019). The negative controls were untreated tachyzoites.

#### Statistical analysis

All data represent the average and standard error of the mean of three independent experiments performed in triplicate. The differences between treatments and controls were assessed using analysis of variance (one-way ANOVA), followed by Tukey's test, and using GraphPad Prism 5.0 software (GraphPad Software, Inc., San Diego, CA, USA). Statistical significance was considered when  $P < 0.05$ .

#### Results

##### Characterization of *M. oleifera* extract by FTIR


From the FTIR spectrum of the aqueous extract of *M. oleifera* (Fig. 1), a band is observed from 3200 to 3500  $\text{cm}^{-1}$  which can be attributed to the presence of hydroxyl groups (OH) in phenolic compounds (Sharaf *et al.*, 2013) and also in proteins, fatty acids, carbohydrates and lignin (Adebisi *et al.*, 2014). The peak at 1645  $\text{cm}^{-1}$  could be related to the C=C stretching vibration of aromatic rings and to the vibration of N-H of amines, C=O of amides and carboxylic groups; also, this band could be related to flavonoids and amino acids (Oliveira *et al.*, 2016).

##### *Moringa oleifera* extract in low concentrations does not alter the viability of HeLa cells

When analysing the viability of HeLa cells treated with the *M. oleifera* extract using the MTT cytotoxicity assay, we observed that the lowest concentrations studied (10, 30 and  $50 \mu\text{g mL}^{-1}$ ) did not significantly reduce the viability of the investigated cells (Fig. 2). On the contrary, concentrations of 100 and  $200 \mu\text{g mL}^{-1}$  significantly altered the viability of HeLa cells ( $P < 0.001$ ). The concentrations of 10 and  $30 \mu\text{g mL}^{-1}$  of the *M. oleifera* extract were chosen for subsequent experiments.

##### *Moringa oleifera* extract reduces infection and proliferation in the face of increased NO and reduces the parasitic burden and replication of *T. gondii* in HeLa cells

To assess the effect of *M. oleifera* treatment during *in vitro* infection with *T. gondii* tachyzoites, we analysed the number of infected cells and the number of tachyzoites within these cells (Fig. 3A and B).



**Fig. 2.** MTT cell viability assay of the *M. oleifera* seed aqueous extract in HeLa cells. HeLa cells were treated with *M. oleifera* extract (10, 30, 50, 100 and 200 µg mL<sup>-1</sup>) for 24 h. Control: cells treated with DMEM. H<sub>2</sub>O<sub>2</sub> (positive control). \*\*\*Significant difference in relation to the negative control ( $P < 0.001$ ). Three independent experiments were carried out in triplicate (one-way ANOVA followed by Tukey's test).

A significant reduction was observed in the infection rate of HeLa cells by tachyzoites treated with the *M. oleifera* extract at a concentration of 30 µg mL<sup>-1</sup> (Fig. 3A) ( $P < 0.01$ ). This was similar to that found in infected cells treated with SDZ + PYR, with reductions of 20.50 and 20.25%, respectively. Both reductions were significant when compared to the control group, infected with untreated HeLa cells (Fig. 3A). The concentration of 10 µg mL<sup>-1</sup> of the *M. oleifera* extract was not able to significantly reduce the number of infected HeLa cells (9.25%), with results similar to that observed in the control without treatment (Fig. 3A) ( $P > 0.05$ ). However, for both concentrations of *M. oleifera*, 30 and 10 µg mL<sup>-1</sup> ( $P < 0.01$ ;  $P < 0.05$ , respectively), there was a significant reduction in the amount of tachyzoites observed inside HeLa cells, in the same way as in the treated control group (SDZ + PYR) ( $P < 0.05$ ) (Fig. 3B).

In view of the promising results of the concentration of 30 µg mL<sup>-1</sup> of *M. oleifera*, both in reducing the rate of infection of cells and in the intracellular proliferation of tachyzoites, this concentration was chosen for the next experiments.

Knowing that NO is the main microbicidal mediator responsible for the elimination of intracellular pathogens, we verified whether treatment with the extract could induce the production of this agent. It was observed that the concentration of 30 µg mL<sup>-1</sup> of the *M. oleifera* extract promoted a significant increase of NO in HeLa cells infected with *T. gondii* compared to untreated infected cells ( $P < 0.001$ ) (Fig. 3C). When we evaluated the parasitic load by qPCR, we observed that treatment with *M. oleifera* (30 µg mL<sup>-1</sup>) was able to reduce the parasitic load of HeLa cells infected with *T. gondii* (strain RH) by 65.8% compared to untreated cells ( $P < 0.001$ ) (Fig. 3D).

The optical microscopy images demonstrate that HeLa cells treated with 30 µg mL<sup>-1</sup> of the *M. oleifera* extract had a smaller number of tachyzoites inside their parasitophorous vacuoles when compared to the untreated control (Fig. 4).

#### Moringa oleifera extract changes the morphology and ultrastructure of *T. gondii* tachyzoites

In Fig. 5A–F, SEM images are presented, whereas Fig. 5G–L show TEM photomicrographs. In Figure 5A and B, we can see the untreated tachyzoites. They were half-moon-shaped, with an enlarged centre and pointed ends. In Fig. 5C–F, we can see the changes in morphology of membrane surface after treatment with the *M. oleifera* extract (30 µg mL<sup>-1</sup>), with the deformation

and rupture of the outer membrane with the leakage of the cytoplasmic contents.

In Fig. 5G and H, we can see that the internal structure of the untreated tachyzoites is preserved. After 1 h of treatment, we can see that the tachyzoites lose their normal morphology and become rounded. It is possible to verify swelling mitochondria, lipid bodies and plasma membrane damage.

#### Moringa oleifera increases levels of ROS, autophagic and lipid vacuoles and interferes with the mitochondrial integrity of *T. gondii* tachyzoites

With the knowledge that *M. oleifera* has an action on HeLa cells infected by *T. gondii*, causing increasing of NO synthesis, and also induces changes in tachyzoites morphology and ultrastructure, we aimed to understand the possible mechanism implicated in tachyzoites death. For that, tachyzoites were treated with the *M. oleifera* extract at the most effective concentration, 30 µg mL<sup>-1</sup>, and possible changes in organelles involved with cell death were investigated.

First, we investigated the synthesis of ROS, important microbicidal molecules. It was found that the treatment increased ROS levels by 23.5-fold when compared to the untreated control ( $P < 0.001$ ), with values similar to those produced by the positive control (H<sub>2</sub>O<sub>2</sub>) (Fig. 6A). In addition, we found that *M. oleifera* induced mitochondrial depolarization, reducing the total fluorescence intensity of the TMRE in 10% in relation to the untreated control group ( $P < 0.01$ ), without differing from the positive control (CCCP) (Fig. 6B and C).

Furthermore, we found that *T. gondii* tachyzoites treated with *M. oleifera* showed an increase of up to 19.2-fold in the formation of lipid bodies when compared to the untreated control ( $P < 0.0001$ ) (Fig. 7A), as well as increase of 185% in the intensity of the MDC fluorescence ( $P < 0.0001$ ), indicating the formation of autophagic vacuoles (Fig. 7B). In both assays, treatment with *M. oleifera* induced similar increases to the positive control (PBS).

#### Moringa oleifera induces death by apoptosis in *T. gondii* tachyzoites


To determine whether the mechanism of action induced by *M. oleifera* in tachyzoites leads to cell death by apoptosis, we evaluated the externalization of phosphatidylserine (PS), a phospholipid confined to the inner face of the plasma membrane and translocated to the cell surface in apoptotic cells, by staining with annexin-V and PI. Figure 8 shows that treatment with 30 µg mL<sup>-1</sup> of *M. oleifera* for 1 h induced a significant increase in annexin-V fluorescence intensity of 119% compared to the negative control, indicating apoptosis. In addition, 34% of parasites exhibited double labelling for annexin-V+/PI+, when compared to untreated parasites, demonstrating that *M. oleifera* also induced characteristic signs of the process of late apoptosis.

#### Discussion


The current study was undertaken to investigate, for the first time, the *in vitro* effect of the *M. oleifera* extract on *T. gondii* tachyzoites and the results indicate that it was able to interfere with both the invasion and intracellular replication of tachyzoites.

It was found that the most striking substances present in the extract were phenolic compounds and flavonoids, since the presence of aromatic rings, structures which are characteristic of these compounds, was observed in the tests performed (Tanase *et al.*, 2019).

*Moringa oleifera* extract at concentrations of 10, 30 and 50 µg mL<sup>-1</sup> presented low toxicity in HeLa cells. In general,



**Fig. 3.** HeLa cells experimentally infected with *Toxoplasma gondii* tachyzoites RH strain, treated with the *M. oleifera* extract (30 and 10 µg mL<sup>-1</sup>). (A) Percentage of cells infected with *T. gondii* tachyzoites. (B) Amount of tachyzoites inside the infected cells. (C) NO production by HeLa cells infected with *T. gondii* tachyzoites and treated with the aqueous extract of *M. oleifera* 30 and 10 µg mL<sup>-1</sup>, during 24 h of treatment. Griess method for nitrite levels in supernatant of culture cells. (D) Parasitic load by qPCR of HeLa cells infected with *T. gondii* tachyzoites and treated with *M. oleifera* (30 µg mL<sup>-1</sup>). The values represent the mean ± s.e.m. of three independent experiments performed in duplicate. The values represent the mean ± s.e.m. of three independent experiments. \*Significant difference compared to negative control ( $P < 0.05$ ), \*\*( $P < 0.01$ ), \*\*\*( $P < 0.001$ ).




**Fig. 4.** HeLa cells experimentally infected with tachyzoites from the *T. gondii* RH strain. (A) Negative control - cells without treatment. (B) Cells treated with aqueous *M. oleifera* seed extract of 30 µg mL<sup>-1</sup>. The arrows indicate the parasitophorous vacuoles with tachyzoites inside.

other studies that evaluated the *in vitro* and *in vivo* effects of different parts of *M. oleifera* plants concluded that this is a safe plant for medicinal and nutritional purposes (Jafarain *et al.*, 2014; Padayachee and Bajjnath, 2020).

When the microbicidal activity was analysed, it was quite evident that the *M. oleifera* extract is capable of reducing intracellular forms of *T. gondii* tachyzoites, as well as interfering with both invasion and intracellular replication. Possibly this anti-*T. gondii* activity may be due to flavonoids and phenolic compounds. Plants rich in flavonoids have shown an effect against protozoa, including anti-*T. gondii* action (Si *et al.*, 2018). For example, the flavonoid Lycopodium, isolated from the root of *Glycyrrhiza* species, belonging to the Fabaceae family, showed activity on *T. gondii* tachyzoites (*in vitro*) and cysts (*in vivo*) (Zhang *et al.*, 2016; Si *et al.*, 2018), in *Plasmodium falciparum* trophozoites (Chen *et al.*, 1994), and in *Leishmania major* and *Leishmania donovani* promastigote and amastigote forms (Chen *et al.*, 1993).

*Moringa oleifera* extract had a dose-dependent effect in relation to the cellular invasion of *T. gondii* tachyzoites, being more efficient at a dose of 30 µg mL<sup>-1</sup> when compared to the group without treatment. This concentration had results similar to those of infected HeLa cells and cells treated with SDZ + PYR. The reduction in parasitic load on HeLa cells was confirmed by qPCR. The decrease in intracellular invasion and proliferation of *T. gondii* tachyzoites has been reported with the use of other plant derivatives (Si *et al.*, 2018; Chen *et al.*, 2019). For example, resveratrol reduced the tachyzoites' population, probably due to a disturbance in the redox homeostasis of the protozoan, cellular stress (apoptosis) of infected cells and the elimination of intracellular tachyzoites (Chen *et al.*, 2019).

*Moringa oleifera* extract also interfered with the redox homeostasis of *T. gondii* tachyzoites and the consequent release of cellular stress substance NO, that is a highly reactive free radical and is synthesized by several cells of the immune system, involved



**Fig. 5.** Morphological and ultrastructural alterations on tachyzoites of *T. gondii* treated with  $30 \mu\text{g mL}^{-1}$  of *M. oleifera* extract for 1 h. SEM images: (A, B) untreated tachyzoites; (C–F) *M. oleifera* extract-treated parasites. TEM images: (G, H) untreated tachyzoites; (I–L) *M. oleifera* extract-treated parasites. a, apicoplast; am, amylopectin granule; c, conoid; dg, dense granule; Gc, Golgi complex; m, mitochondria; mi, microneme; n, nucleus; r, rhoptry; \*, autophagic vacuoles; ◆, swelling mitochondrial; ▲, lipid-storage bodies; black arrow and white arrow head, plasma membrane damage; white arrow, leakage of cytoplasmic contents. Scale bars =  $2 \mu\text{m}$  (A–E),  $1 \mu\text{m}$  (F),  $0.5 \mu\text{m}$  (G, I–L),  $0.2 \mu\text{m}$  (H).


in the intracellular inactivation of pathogens by cytotoxic mechanisms (Coleman, 2001); at high concentrations, it can directly destroy *T. gondii* tachyzoites (Pimenta *et al.*, 2018).

SEM and TEM analyses showed that the tachyzoites treated with the *M. oleifera* extract present rupture and membrane deformation, lost internal organization, becoming rounded and developing lipid bodies. Similar results were obtained by Si *et al.* (2018), who used another plant derived from *T. gondii* tachyzoites and observed alterations in morphology and ultrastructure of the parasite. The changes observed in SEM and TEM were also confirmed by the tests that were carried out to verify the mechanisms of tachyzoites' death.

The treatment with the *M. oleifera* extract increased ROS production by *T. gondii* tachyzoites. ROS act as microbicides and can lead to cell damage and parasite destruction (Bortoleti *et al.*, 2019), organelle rupture and cell death by necrosis (Zong and

Thompson, 2006). It is worth remembering that the increase in ROS production causes, directly or indirectly, severe damage to biological macromolecules such as lipids, proteins and DNA, and a loss of membrane integrity (Desoti *et al.*, 2012).

Another interesting observation was that *M. oleifera* extract directly affected the mitochondrial function of *T. gondii* tachyzoites detected by a reduction in the total fluorescence intensity (TMRE). In other studies, it has been reported that flavonoids inhibit tyrosine, protein kinases, topoisomerase activity, mitochondrion function and fatty acid type II synthesis; they also disrupt cytoplasmic and plasma membrane integrity (Abugri and Witola, 2020). By compromising the integrity of the only mitochondria existing in this protozoan, we suggest that the *M. oleifera* extract has important pharmacological properties in terms of inhibiting the signalling of cell survival or death factors (Szweczyk and Wojtczak, 2002). The observed changes can lead to apoptosis,



**Fig. 6.** *Toxoplasma gondii* tachyzoites subjected to treatment with 30 µg mL<sup>-1</sup> aqueous *M. oleifera* seed extract. The following methods were used for the respective assessments: (A) H<sub>2</sub>DCFDA probe for reactive species of oxygen measurement and (B) TMRE assay for fluorometric analysis of the mitochondrial membrane potential. (C) Fluorescence microscopy images of tachyzoites after H<sub>2</sub>DCFDA and TMRE labelling. Data represents the mean ± s.e.m. of three independent experiments performed in duplicate. \*\*Significant difference compared to control ( $P < 0.01$ ), \*\*\*\*( $P < 0.0001$ ).

ROS production and increased mitochondrial permeability (Belyaeva *et al.*, 2006). The loss of mitochondrial integrity of *T. gondii* tachyzoites with the use of other plant-derived compounds such as artemisine and other active compounds have also been reported by other authors (Giovati *et al.*, 2018; Rosenberg *et al.*, 2019).


Treatment with the *M. oleifera* extract caused vacuolization of the cytoplasm of *T. gondii* tachyzoites, these results may suggest that autophagy mechanisms are involved in the elimination of parasites. Many studies have evaluated flavonoids as potential autophagy-inducing agents in toxoplasmosis infection and other diseases (Prieto-Domínguez *et al.*, 2018; Lee *et al.*, 2020). Studies indicate that autophagy mechanisms are involved in the elimination of intracellular and extracellular *T. gondii* tachyzoites (Lavine and Arrizabalaga, 2012; Lee *et al.*, 2020).

Treatment with the *M. oleifera* extract may have influenced the metabolism of intracellular lipids and the formation of neutral lipid aggregates inside the tachyzoites, as evidenced by the increase in fluorescence after staining with Nile Red. The influence of plant-derived compounds on the *T. gondii* lipid metabolism was also observed by Si *et al.* (2018), who evaluated the flavonoid LicoA (4 µg mL<sup>-1</sup>) in *T. gondii* tachyzoites inside human foreskin fibroblast (HFF) cells, found an increase in intracellular lipids.


By staining with annexin-V/PI, it was possible to verify that the main action of the *M. oleifera* extract in *T. gondii* tachyzoites was compatible with late apoptosis, as indicated by the translocation of PS from the inner surface of the tachyzoites membrane to the outer surface (Shamseddin *et al.*, 2015). The apoptosis-like death mechanism was identified in other studies against *T. gondii*. Giovati *et al.* (2018), studying anti-*T. gondii* activity *in vitro* (tachyzoites RH strain) for a synthetic decapeptide (killer peptide KP), identified an apoptosis-like mechanism verified by annexin-V labelling, the modification of mitochondrial membrane potential, DNA fragmentation and morphological changes observed by TEM, similar to the results presented in this study.

### Conclusions

In view of the results obtained, it is possible to infer that *M. oleifera* extract, more precisely flavonoids and phenolic compounds, have important pharmacological properties. An increase in the production of NO by HeLa cells, leading to the reduction of intracellular forms of *T. gondii* tachyzoites and also acting directly on the parasite, inducing cell death with expressive alterations in *T. gondii* machinery components and their function, probably in order to induce mechanisms of late apoptosis, stimulate the production of ROS and interfere with lipid metabolism, promoting alterations to the parasite mitochondrial membrane.



**Fig. 7.** *Toxoplasma gondii* tachyzoites subjected to treatment with aqueous *M. oleifera* seed extract of  $30 \mu\text{g mL}^{-1}$ . The following methods were used for the respective assessments: (A) Nile Red and (B) MDC. Data represent the mean  $\pm$  s.e.m. of three independent experiments performed in duplicate. \*\*\*\*Significant difference compared to control ( $P < 0.0001$ ).



**Fig. 8.** *Moringa oleifera* extract induces PS exposure in *T. gondii* tachyzoites subjected to 1 h of treatment with  $30 \mu\text{g mL}^{-1}$  of the extract using annexin V/FITC and PI. (A) Fluorescence intensity of parasites stained with annexin V/FITC and PI. (B) Typical dot plots of at least three independent experiments are shown. Data represent the mean  $\pm$  s.e.m. of three independent experiments performed in duplicate. \*Significant difference in relation to control ( $P < 0.05$ ).

The results demonstrate that the *M. oleifera* extract exhibits potential *in vitro* activity against *T. gondii*, and that it is extremely important to conduct studies to continue these findings, in an attempt to seek treatment options for acute phase of toxoplasmosis and mainly in the chronic phase of infection, since the inefficacy of drugs in this phase is the main motivators for the studies with alternatives treatments for toxoplasmosis.

**Acknowledgements.** The authors thank Prof. Dr João Luís Garcia of the State University of Londrina for ceding the RH strain of *T. gondii*. The authors also acknowledge Complexo de Centrais de Apoio a Pesquisa of the State University of Maringá (COMCAP-UEM) for the support in electron microscopy analysis.

**Author contributions.** Conceived and designed the experiments: LN, RASS, INC, WRP and ALFG. Performed the experiments: LN, RASS, BTSB, FTP, TFS, FFE, DLB and ATAB. Analysed the data: LN, RASS, FTP, BTSB, TFS, DLB, ATAB, RB, INC, WRP and ALFG. Contributed reagents/materials/analysis tools: LN, RB, INC, WRP, ICC and ALFG. Wrote the paper: LN, RASS, DLB, INC and ALFG.

**Financial support.** This study was supported by Coordination for the Improvement of Higher Education Personnel Brazil (CAPES) – Financing Code 001.

**Conflict of interest.** The authors declare there are no conflicts of interest.

## References

Abugri DA and Witola WH (2020) Interaction of apigenin-7-O-glucoside with pyrimethamine against *Toxoplasma gondii* growth. *Journal of Parasitic Diseases* **44**, 221–229.

- Adebisi F, Adedayo A, Oluwaseye A, Adekunle F, Michael A, Olutayo O, Clement A, Ikkoh P, Rasheed K and Ademola A (2014) Instrumental and chemical characterization of *Moringa oleifera* Lam root starch as an industrial biomaterial. *Research in Pharmaceutical Biotechnology* **3**, 7–12.
- Almeida AQ, Souza RMS, Loureiro DC, Pereira DR, Cruz MAS and Vieira JS (2017) Modelling the spatial dependence of the rainfall erosivity index in the Brazilian semi-arid region. *Pesquisa Agropecuária Brasileira* **52**, 371–379.
- Arruda CCL, Freitas DV, Seabra MABL, Xavier-Júnior FH, Figueiredo RCBQ, Napoleão TH, Paiva PMG, Navarro DMAF and Navarro M (2020) CdTe-GSH as luminescent biomarker for labeling the larvicidal action of WSMoL lectin in *Aedes aegypti* larvae. *Colloids and Surfaces B: Biointerfaces* **187**, 110672.
- Baptista ATA, Silva MO, Gomes RG, Bergamasco R, Vieira MF and Vieira AMS (2017) Protein fractionation of seeds of *Moringa oleifera* Lam and its application in superficial water treatment. *Separation and Purification Technology* **180**, 114–124.
- Belyaeva EA, Dymkowska D, Wieckowski MR and Wojtczak L (2006) Reactive oxygen species produced by the mitochondrial respiratory chain are involved in Cd<sup>2+</sup>-induced injury of rat ascites hepatoma AS-30D cells. *Biochimica et Biophysica Acta* **1757**, 1568–1574.
- Bortoleti B, Tomiotto-Pellissier F, Gonçalves MD, Miranda-Sapla MM, Assolini JP, Carlotto AC, Lima DM, Silveira GF, Almeida RS, Costa IN, Conchon-Costa I and Pavanelli WR (2019) Caffeic acid has antipromastigote activity by apoptosis-like process; and anti-amastigote by TNF- $\alpha$ /ROS/NO production and decreased of iron availability. *Phytomedicine: International Journal of Phytotherapy and Phytopharmacology* **57**, 262–270.
- Bortoleti BTS, Gonçalves MD, Tomiotto-Pellissier F, Contato VM, Silva TF, Matos RLN, Detoni MB, Rodrigues ACJ, Carlotto AC, Lazarin DB, Arakawa NS, Costa IN, Conchon-Costa I, Miranda-Sapla MM, Wowk PF and Pavanelli WR (2021) Solidagenone acts on promastigotes of *L. amazonensis* by inducing apoptosis-like processes on intracellular

- amastigotes by IL-12p70/ROS/NO pathway activation. *Phytomedicine: International Journal of Phytotherapy and Phytopharmacology* **85**, 153536.
- Chen M, Christensen SB, Blom J, Lemmich E, Nadelmann L, Fich K, Theander TG and Kharazmi A** (1993) Licochalcone A, a novel antiparasitic agent with potent activity against human pathogenic protozoan species of *Leishmania*. *Antimicrobial Agents and Chemotherapy* **37**, 2550–2556.
- Chen M, Theander TG, Christensen SB, Hviid L, Zhai L and Kharazmi A** (1994) Licochalcone A, a new antimalarial agent, inhibits *in vitro* growth of the human malaria parasite *Plasmodium falciparum* and protects mice from *P. yoelii* infection. *Antimicrobial Agents and Chemotherapy* **38**, 1470–1475.
- Chen QW, Dong K, Qin HX, Yang YK, He JL, Li J, Zheng ZW, Chen DL and Chen JP** (2019) Direct and indirect inhibition effects of resveratrol against *Toxoplasma gondii* tachyzoites *in vitro*. *Antimicrobial Agents and Chemotherapy* **63**, 1–17. doi: 10.1128/aac.01233-18.
- Coleman JW** (2001) Nitric oxide in immunity and inflammation. *International Immunopharmacology* **1**, 1397–1406.
- Coriolano MC, Brito JS, Ferreira GRS, Moura MC, Melo CML, Soares AKA, Lorena VMB, Figueiredo RCBQ, Paiva PMG, Napoleão TH and Coelho LCBB** (2020) Antibacterial lectin from *Moringa oleifera* seeds (WSMoL) has differential action on growth, membrane permeability and protease secretory ability of Gram-positive and Gram-negative pathogens. *South African Journal of Botany* **129**, 198–205.
- Desoti VC, Lazarin-Bidóia D, Sudatti DB, Pereira RC, Alonso A, Ueda-Nakamura T, Dias Filho BP, Nakamura CV and Silva SO** (2012) Trypanocidal action of (–)-elatal involves an oxidative stress triggered by mitochondrial dysfunction. *Marine Drugs* **10**, 1631–1646.
- Dhakad AK, Ikram M, Sharma S, Khan S, Pandey VV and Singh A** (2019) Biological, nutritional, and therapeutic significance of *Moringa oleifera* Lam. *Phytotherapy Research* **33**, 2870–2903.
- Giovati L, Santinoli C, Mangia C, Vismarra A, Belletti S, D'Adda T, Fumarola C, Ciociola T, Bacci C, Magliani W, Polonelli L, Conti S and Kramer LH** (2018) Novel activity of a synthetic decapeptide against *Toxoplasma gondii* tachyzoites. *Frontiers in Microbiology* **9**, 753.
- Jafarain A, Asghari G and Ghassami E** (2014) Evaluation of cytotoxicity of *Moringa oleifera* Lam. callus and leaf extracts on HeLa cells. *Advanced Biomedical Research* **3**, 1–5.
- Konstantinovic N, Guegan H, Stājner T, Belaz S and Robert-Gangneux F** (2019) Treatment of toxoplasmosis: current options and future perspectives. *Food and Waterborne Parasitology* **15**, e00036.
- Lavine MD and Arrizabalaga G** (2012) Analysis of monensin sensitivity in *Toxoplasma gondii* reveals autophagy as a mechanism for drug induced death. *PLoS One* **7**, e42107.
- Lee J, Choi JW, Han HY, Kim WS, Song HY, Byun EB, Byun EH, Lee YH and Yuk JM** (2020) 4-Hydroxybenzaldehyde restricts the intracellular growth of *Toxoplasma gondii* by inducing SIRT1-mediated autophagy in macrophages. *The Korean Journal of Parasitology* **58**, 7–14.
- Madrona GS, Serpelloni GB, Salcedo Vieira AM, Nishi L, Cardoso KC and Bergamasco R** (2010) Study of the effect of saline solution on the extraction of the *Moringa oleifera* seed's active component for water treatment. *Water, Air, & Soil Pollution* **211**, 409–415.
- Mirzaalizadeh B, Sharif M, Daryani A, Ebrahimzadeh MA, Zargari M, Sarvi S, Mehrzadi S, Rahimi MT, Mirabediny Z, Golpour M and Montazeri M** (2018) Effects of *Aloe vera* and *Eucalyptus* methanolic extracts on experimental toxoplasmosis *in vitro* and *in vivo*. *Experimental Parasitology* **192**, 6–11.
- Mosmann T** (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* **65**, 55–63.
- Mota WM, Barros ML, Cunha PEL, Santana MVA, Stevam CS, Leopoldo PTG and Fernandes RPM** (2012) Avaliação da inibição da acetilcolinesterase por extratos de plantas medicinais. *Revista Brasileira de Plantas Mediciniais* **14**, 624–628.
- Oliveira RN, Mancini MC, Oliveira FCS, Passos TM, Quilty B, Thiré RMSM and McGuinness GB** (2016) FTIR Analysis and quantification of phenols and flavonoids of five commercially available plants extracts used in wound healing. *Matéria (Rio de Janeiro)* **21**, 767–779.
- Opsteegh M, Langelaar M, Sprong H, den Hartog L, De Craeye S, Bokken G, Aizenberg D, Kijlstra A and van der Giessen J** (2010) Direct detection and genotyping of *Toxoplasma gondii* in meat samples using magnetic capture and PCR. *International Journal of Food Microbiology* **139**, 193–201.
- Padayachee B and Baijnath H** (2020) An updated comprehensive review of the medicinal, phytochemical and pharmacological properties of *Moringa oleifera*. *South African Journal of Botany* **129**, 304–316.
- Pagar K, Ghotekar S, Pagar T, Nikam A, Pansambal S, Oza R, Sanap D and Dabhane H** (2020) Antifungal activity of biosynthesized CuO nanoparticles using leaves extract of *Moringa oleifera* and their structural characterizations. *Asian Journal of Nanosciences and Materials* **3**, 15–23.
- Pimenta TS, Chaves NF, Rodrigues APD, Diniz CWP, DaMatta RA and Diniz JAP** (2018) Granulocyte macrophage colony-stimulating factor alone reduces *Toxoplasma gondii* replication in microglial culture by superoxide and nitric oxide, without IFN-gamma production: a preliminary report. *Microbes and Infection* **20**, 385–390.
- Prieto-Domínguez N, Garcia-Mediavilla MV, Sanchez-Campos S, Mauriz JL and Gonzalez-Gallego J** (2018) Autophagy as a molecular target of flavonoids underlying their protective effects in human disease. *Current Medicinal Chemistry* **25**, 814–838.
- Robert-Gangneux F, Meroni V, Dupont D, Botterel F, Garcia JMA, Brenier-Pinchart MP, Accoceberry I, Akan H, Abbate I, Boggian K, Bruschi F, Carratalà J, David M, Držgona L, Djurković-Djaković O, Farinas MC, Genco F, Gkrania-Klotsas E, Groll AH, Guy E, Hirzel C, Khanna N, Kurt Ö, Junie LM, Lazzarotto T, Len O, Mueller NJ, Munoz P, Pana ZD, Roilides E, Stājner T, van Delden C, Villena I, Pelloux H and Manuel O** (2018) Toxoplasmosis in transplant recipients, Europe, 2010–2014. *Emerging Infectious Diseases* **24**, 1497–1504.
- Rosenberg A, Luth MR, Winzeler EA, Behnke M and Sibley LD** (2019) Evolution of resistance *in vitro* reveals mechanisms of artemisinin activity in *Toxoplasma gondii*. *Proceedings of the National Academy of Sciences* **116**, 26881–26891.
- Sanfelice RA, Machado LF, Bosqui LR, Miranda-Sapla MM, Tomiotto-Pellissier F, Alcântara Dalevedo G, Ioris D, Reis GF, Panagio LA, Navarro IT, Bordignon J, Conchon-Costa I, Pavanelli WR, Almeida RS and Costa IN** (2017) Activity of rosuvastatin in tachyzoites of *Toxoplasma gondii* (RH strain) in HeLa cells. *Experimental Parasitology* **181**, 75–81.
- Shamseddin J, Akhlaghi L, Razmjou E, Shojae S, Monavari SHR, Tajik N, Ebrahimi SA and Meamar AR** (2015) Conjugated linoleic acid stimulates apoptosis in RH and Tehran strains of *Toxoplasma gondii*, *in vitro*. *Iranian Journal of Parasitology* **10**, 238–244.
- Sharaf S, Higazy A and Hebeish A** (2013) Propolis induced antibacterial activity and other technical properties of cotton textiles. *International Journal of Biological Macromolecules* **59**, 408–416.
- Si H, Xu C, Zhang J, Zhang X, Li B and Zhou X** (2018) Licochalcone A: an effective and low-toxicity compound against *Toxoplasma gondii* *in vitro* and *in vivo*. *International Journal for Parasitology – Drugs and Drug Resistance* **8**, 238–245.
- Silva LA, Fernandes MD, Machado AS, Reis-Cunha JL, Bartholomeu DC and Almeida Vitor RW** (2019) Efficacy of sulfadiazine and pyrimetamine for treatment of experimental toxoplasmosis with strains obtained from human cases of congenital disease in Brazil. *Experimental Parasitology* **202**, 7–14.
- Somsak V, Borkaew P, Klubsri C, Dondee K, Bootprom P and Saiphet B** (2016) Antimalarial properties of aqueous crude extracts of *Gynostemma pentaphyllum* and *Moringa oleifera* leaves in combination with artesunate in *Plasmodium berghei*-infected mice. *Journal of Tropical Medicine* **2016**, 8031392.
- Su C and Dubey JP** (2020) Isolation and genotyping of *Toxoplasma gondii* strains. In Tonkin CJ (ed.), *Toxoplasma gondii: Methods and Protocols*. New York, USA: Springer Science + Business Media, LLC, pp. 49–80.
- Szewczyk A and Wojtczak L** (2002) Mitochondria as a pharmacological target. *Pharmacological Reviews* **54**, 101–127.
- Tanase C, Coşarcă S and Muntean DL** (2019) A critical review of phenolic compounds extracted from the bark of woody vascular plants and their potential biological activity. *Molecules (Basel, Switzerland)* **24**, 1182.
- Tomiotto-Pellissier F, Alves DR, Miranda-Sapla MM, Morais SM, Assolini JP, Silva Bortoleti BT, Gonçalves MD, Cataneo AHD, Kian D, Madeira TB, Yamauchi LM, Nixdorf SL, Costa IN, Conchon-Costa I and Pavanelli WR** (2018) *Caryocar coriaceum* extracts exert leishmanicidal effect acting in promastigote forms by apoptosis-like mechanism and intracellular amastigotes by Nrf2/HO-1/ferritin dependent response and iron depletion: Leishmanicidal effect of *Caryocar coriaceum* leaf extracts. *Biomedicine & Pharmacotherapy* **98**, 662–672.

- Turner DN, Just J, Dasari R, Smith JA, Bissember AC, Kornienko A and Rogelj S** (2019) Activity of natural and synthetic polygodial derivatives against *Trypanosoma cruzi* amastigotes, trypomastigotes and epimastigotes. *Natural Product Research* **35**, 792–795.
- Zhang X, Jin L, Cui Z, Zhang C, Wu X, Park H, Quan H and Jin C** (2016) Antiparasitic effects of oxymatrine and matrine against *Toxoplasma gondii* *in vitro* and *in vivo*. *Experimental Parasitology* **165**, 95–102.
- Zong WX and Thompson CB** (2006) Necrotic death as a cell fate. *Genes & Development* **20**, 1–15.
- Zwicker JD, Smith D, Guerra AJ, Hitchens JR, Haug N, Vander Roest S, Lee P, Wen B, Sun D, Wang L, Keep RF, Xiang J, Carruthers VB and Larsen SD** (2020) Discovery and optimization of triazine nitrile inhibitors of *Toxoplasma gondii* cathepsin L for the potential treatment of chronic toxoplasmosis in the CNS. *ACS Chemical Neuroscience* **11**, 2450–2463.