

Chemical constituents and leishmanicidal activity from leaves of *Kielmeyera variabilis*

Betania B. Cota,^{*,1} Ezequias P. Siqueira,¹ Djalma M. de Oliveira,³
Tânia M. A. Alves,¹ Marcos E. G. Sobral,⁴ Ana Rabello,² Carlos
L. Zani¹

¹Laboratório de Química de Produtos Naturais, Centro de Pesquisas René Rachou, Fiocruz, Minas Gerais, Brazil,

²Laboratório de Pesquisas Clínicas do CPqRR-Fiocruz, Minas Gerais; Brazil,

³Departamento de Química e Exatas, Universidade Estadual do Sudoeste da Bahia, Campus Jequié, Brazil,

⁴Departamento de Ciências Naturais, Universidade Federal de São João Del-Rei, Minas Gerais, Brazil.

Article

Received 16 Mar 2012

Accepted 17 May 2012

Available online 16 Aug 2012

Abstract: Many phenolic compounds such as xanthenes, quinones and coumarins have been isolated from *Kielmeyera* species; however the presence of flavonoids have been showed in other genera in the Calophylleae tribe as *Caraipa*, *Mesua* and *Calophyllum*. Six known glycosidic flavonoids: quercetin 3- β -*O*-galactopyranoside (**1**), quercetin 3- β -*O*-glucopyranoside (**2**), quercetin 3-*O*- α -rhamnoside (**3**), luteolin 6-*C*- β -glucopyranoside (**4**), isovitexin (**5**), kaempferol 3-*O*- α -rhamnoside (**6**) and one triterpene, lupenone (**7**) were isolated, for the first time, from organic crude extract of *Kielmeyera variabilis* Mart. & Zucc., Calophyllaceae, leaves. The crude organic extract from *K. variabilis* leaves exhibited 95% of leishmanicidal activity at 20 μ g/mL on amastigote-like form of *Leishmania (Leishmania) amazonensis* *in vitro* model and only compound **3** showed 40-45% of growth inhibition at concentration ranging from 0.78 to 20 μ g/mL. In addition, quercetin 3-*O*- α -rhamnoside (quercitrin) was found to be the major metabolite. Our results and previous reports suggest that synergistic effects of flavonoid glycosides are the cause of significant leishmanicidal activity of the crude organic extract from *K. variabilis* leaves.

Keywords:

Kielmeyera variabilis
phenolic compounds
C-flavonoid
glycosides
lupenone
leishmanicidal activity

ISSN 0102-695X

<http://dx.doi.org/10.1590/S0102-695X2012005000095>

Introduction

Kielmeyera variabilis Mart. & Zucc., Calophyllaceae, is a shrub or tree that reaches up to 6 m in height and is endemic to the Cerrado (Bittrich, 2010), the savanna-like vegetation of Central-Brazil. It is popularly known as “pau-santo”, “rosa-do-campo” or “folha-santa”. Methanol extract obtained from *K. variabilis* stems showed molluscicidal activity against *Biomphalaria glabrata* (12.5 ppm) (Pinheiro et al., 2003a) and activity against *Staphylococcus aureus* (3.0 μ g/mL) and *Bacillus subtilis* (1.95 μ g/mL). A phytochemical study of this species led to the isolation of three xanthenes (assiguxanthone-B, kielcorin and 1,3,5,6-tetrahydroxy-2-prenylxanthone) and an organic acid (2,5-dihydroxy benzoic acid), being assiguxanthone-B active against *B. subtilis* (25 μ g/mL) (Pinheiro et al., 2003b).

The occurrence of coumarins as phenylcoumarins (Gramacho et al., 1999), prenylated 4-*n*-propylcoumarins (Cruz et al., 1998a,b; Scio et al., 2003) and xanthenes (Garcia Cortez et al., 1998, 2002) are frequently reported

as isolated compounds from *Kielmeyera* genus. An osajaxanthone, isolated from a stem extract of *Kielmeyera coriacea* showed protection against infection by cercariae of *Schistosoma mansoni* (Lopes et al., 1977) and other ones have exhibited antifungal activity against the pathogenic plant fungi *Cladosporium cucumerinum* and *Candida albicans* (Garcia Cortez et al., 1998). Extracts of leaves from *K. coriacea* have demonstrated anxiolytic activity after oral treatment with 120 mg/kg body weight per day for thirty days (Audi et al., 2002). Ethanolic extract from stem of *K. coriacea* at a dose of 60 mg/kg body weight and its dichloromethane semi-pure fraction at dose of 6 mg/kg body weight showed similar antidepressant-like effects after 45 days of oral treatment in Male Wistar rats (Martins et al., 2004, 2006). Hexane and dichloromethane extracts from different parts of this species have anticancer activity against SF-295 (brain); HCT-8 (human colon carcinoma); MDA-MB-435 (melanome) and HL-60 (leukemia) with IC₅₀ values ranging from 3.0 to 34.9 μ g/mL (Mesquita et al., 2009). Recently, Sobral et al. (2009) demonstrated antibacterial activity against *Micrococcus luteus* (CIM 7.8 μ g/mL), *Bacillus subtilis*

(CIM 15.6 $\mu\text{g/mL}$), *Staphylococcus aureus* (CIM 31.2 $\mu\text{g/mL}$) and *Streptococcus mutans* (CIM 31.2 $\mu\text{g/mL}$) of the dichloromethane extract from trunk of *Kielmeyera cuspidata*.

In this work we describe for the first time the isolation of one triterpene and the occurrence of flavonoids, including *O*-flavonoids and *C*-flavonoids, from leaves extract of *K. variabilis*. In addition, we suggested that quercetin 3-*O*- α -rhamnoside is among the leishmanicidal compounds from crude extract.

Materials and Methods

Plant material

Leaves of *Kielmeyera variabilis* Mart. & Zucc., Calophyllaceae, were collected in Brumadinho, Minas Gerais, Brazil. A voucher specimen was identified and deposited by Dr. Marcos E. G. Sobral in the Herbarium BHCB, at Universidade Federal de Minas Gerais, Instituto de Ciências Biológicas (Stehmann, number 2646) in Belo Horizonte-MG.

General procedures

The solvents were removed using a SpeedVac[®] vacuum centrifuge (ThermoSavant SPD SC250, Holbrook, NY, USA). Thin-layer chromatographic (TLC) analyses were conducted on pre-coated commercial silica gel G-60/F₂₅₄ (0.25 mm, Merck, Darmstadt, Germany) plates. They were eluted in a pre-saturated chamber using solvents mixtures in different proportions of: a) chloroform:methanol:water (65:50:5); b) hexane:ethyl acetate (70:30) or c) ethyl acetate:methanol:formic acid:glacial acetic acid:water (100:11:11:26). The spots were visualized under visible, UV light at 254 nm and 360 nm, and after spraying the plate with mixture (1:1) of ethanolic solutions of vanillin (1%) and sulfuric acid (10%). Medium-pressure liquid chromatography (MPLC) was performed using an LC8 pump (Shimadzu, Kyoto, Japan), Büchi columns n^o 19676 and n^o 19674 for reverse (RP-18, 25-40 mesh) and normal (silica gel 60, 25-40 mesh) phases, respectively, at flow rates of 20 mL/min and detection at λ 220 nm and 254 nm. Semi-preparative purifications were performed using Shim-pack[®] C18 column (5 μm , 20x250 mm, *i.d.*), mixtures of MeOH:H₂O as eluents, at flow rates of 10 mL/min and detection at λ 220 and 254 nm using a Shimadzu chromatographic system (Shimadzu, Kyoto, Japan) equipped with an LC6AD pump and UV-visible dual-wavelength detector (SPD10A). Analytical RP-HPLC analyses were developed in a HPLC system (Shimadzu, Kyoto, Japan) equipped with a LC6AD pump and a SPD M-10A VP Diode Array Detector. Analyses were carried out by Shim-pack[®] C18 column (5 μm , 4.6x250 mm *i.d.*) eluted with gradient of

MeOH/H₂O from 10% to 100% of MeOH at a flow rate of 1 mL/min. Gel Permeation Chromatography (GPC) was carried out using glass columns filled with Sephadex LH-20TM (GE Healthcare, U.S.A.) gel and MeOH as mobile phase. NMR (nuclear magnetic resonance) spectra were obtained using a Bruker DRX 400 spectrometer at 400 MHz with the pulse programs provided by the manufacturer. The substances were dissolved in perdeuterated solvents containing 0.1% tetramethylsilane as the internal chemical shift standard. UV spectra (200-500 nm) were obtained by Shimadzu SPD84-10A VP Diode Array Detector from HPLC system (Shimadzu, Kyoto, Japan). Electrospray ionization mass spectrometry (ESI-MS) were performed using a Bruker maXis ETD (Bruker, DE). Solutions of the compounds at 200 $\mu\text{g/mL}$ in ACN:H₂O (1:1) were infused at 3 $\mu\text{L/min}$ with 0.1% formic acid in the negative mass spectra acquired over a range between *m/z* 50-1500 daltons. The instrument was operated under the following conditions: end plate offset, -500 V; capillary voltage, 4500 V; nebulizer pressure, 0.4 bar; dry gas (nitrogen) flow rate, 4.0 L/min; dry temperature, 180 °C; ISCID energy, 0-20 eV; hexapole RF, 600 Vpp; collision energy, 0-30 eV; collision RF, 600 Vpp; transfer time, 30-100 μs .

Extraction and isolation

Leaves (62 g) of *K. variabilis* were extracted with a dichloromethane and MeOH mixture (1:1) at room temperature. The solution was concentrated under vacuum in a rotary evaporator at temperatures below 45 °C, and the residual solvent was removed in a vacuum centrifuge at 40 °C. Organic crude extract (3 g) was suspended in MeOH:H₂O (1:1) and extracted with hexane. This procedure afforded a Hex soluble layer (Hex, 0.6 g) and an aqueous soluble layer (Aq, 1.8 g) fractions. Solvents from fractions were dried using a vacuum centrifuge. The Aq fraction was subjected to reverse-phase MPLC separation using increasing amounts of 10% aq. MeOH to produce 31 fractions. Fractions 15 (MeOH:H₂O; 1:1) and 17 (MeOH:H₂O 70:30) yielded a mixture of compounds **1** and **2** (15 mg) and a pure compound **3** (200 mg), respectively. Fraction 13 (MeOH:H₂O; 1:1) was purified on semi-preparative, using a gradient of MeOH:H₂O (MeOH, 10-100% in 40 min) to obtain **4** (3 mg). Fraction 14 (MeOH; 100%) (45 mg) was purified by GPC to afford **5** (7 mg). Fraction 18 (MeOH:H₂O; 60:40) (49 mg) was submitted to semi-preparative HPLC, using a Shim-pack[®] C18 column and eluted with mixtures of MeOH:H₂O (50:50→70:30 in 30 min; 50:40→60:40 in 2 min; 60:40 in 8 min) to obtain **6** (3 mg). The Hex fraction was subjected to a normal MPLC, developed with petroleum ether containing increasing amounts of EtOAc to yield seventeen fractions. Fraction 12 (210 mg) was purified over silica gel column, eluted with Hex-EtOAc (97:3), and then isolated by GPC to yield **7** (110 mg).

Quercetin 3- β -*O*-galactopyranoside or hyperoside (**1**) UV $\lambda_{\text{max}}^{\text{MeOH}}$ /nm: 256, 356; NMR ^1H (400 MHz, CD_3OD): δH 7.82 (1H, *d*, *J* 2.0 Hz H-2'), 7.62 (1H, *dd*, *J* 8.6 and 1.1 Hz, H-6''), 6.89 (1H, *dd*, *J* 8.6 and 2.0 Hz, H-5'), 6.43 (1H, *d*, *J* 2.0 Hz, H-8), 6.23 (1H, *d*, *J* 2.0 Hz, H-6), 5.26 (1H, *d*, *J* 7.6 Hz, H-1''), 3.86 (1H, *m*, H-4''), 3.81 (1H, *dd*, *J* 9.5 and 8.0 Hz, H-2''), 3.65 (2H, *dd*, *J* 11.2 and 6.0 Hz, H-6''), 3.56 (1H, *m*, H-3''), 3.49 (1H, *t*, *J* 6.0 Hz, H-5''). NMR ^{13}C (100 MHz, CD_3OD): δC 179.61 (C-4) 166.13 (C-7), 163.19 (C-5), 158.92 (C-2), 158.56 (C-9), 150.06 (C-4'), 146.12 (C-3'), 135.66 (C-3), 123.34 (C-1'), 123.16 (C-6'), 117.92 (C-2'), 116.39 (C-5'), 105.66 (C-10), 105.16 (C-1''), 100.07 (C-6), 94.91 (C-8), 77.37 (C-5''), 75.18 (C-3''), 73.25 (C-2''), 70.12 (C-4''), 62.10 (C-6''). ESI-MS *m/z* negative mode: 463.0788 [M-H]⁻; MS² [300.0216].

Quercetin 3- β -*O*-glucopyranoside or isoquercitrin (**2**) UV $\lambda_{\text{max}}^{\text{MeOH}}$ /nm: 256, 356; NMR ^1H (400 MHz, CD_3OD): δH 7.62 (1H, *dd*, *J* 8.6 and 1.1 Hz, H-6''), 7.73 (1H, *d*, *J* 2.0 Hz H-2'), 6.89 (1H, *dd*, *J* 8.6 and 1.8 Hz, H-5'), 6.43 (1H, *d*, *J* 2.0 Hz, H-8), 6.23 (1H, *d*, *J* 2.0 Hz, H-6), 5.35 (1H, *d*, *J* 7.2 Hz, H-1''), 3.72 (2H, *dd*, *J* 12.0 and 2.2 Hz, H-6''a), 3.58 (m, H-6''b), 3.47 (1H, *m*, H-2''), 3.43 (1H, *m*, H-5''), 3.35 (1H, *m*, H-4''), 3.24 (1H, *ddd*, *J* 9.0, 5.0 and 2.0 Hz, H-3''). NMR ^{13}C (100 MHz, CD_3OD): δC 179.57 (C-4), 166.07 (C-7), 163.19 (C-5), 158.79 (C-2), 158.53 (C-9), 149.97 (C-4'), 146.12 (C-3'), 135.65 (C-3), 123.33 (C-1'), 123.17 (C-6'), 117.76 (C-2'), 116.31 (C-5'), 105.66 (C-10), 104.10 (C-1''), 100.07 (C-6), 94.91 (C-8), 78.63 (C-5''), 78.26 (C-3''), 75.87 (C-2''), 71.41 (C-4''), 62.68 (C-6''). ESI-MS *m/z* negative mode: 463.0788 [M - H]⁻; MS² [300.0216].

Quercetin 3-*O*- α -rhamnoside or quercitrin (**3**) UV $\lambda_{\text{max}}^{\text{MeOH}}$ /nm: 256, 350; NMR ^1H (400 MHz, CD_3OD): δH 7.34 (1H, *dd*, *J* 8.3 and 2.0 Hz, H-2'), 7.30 (1H, *dd*, *J* 8.3 and 2.0 Hz, H-6'), 6.93 (1H, *d*, *J* 8.3 Hz, H-5'), 6.36 (1H, *d*, *J* 2.0, H-8), 6.19 (1H, *d*, *J* 2.0 Hz, H-6), 5.34 (1H, *d*, *J* 1.3 Hz, H-1''), 4.21 (1H, *dd*, *J* 3.0 and 1.5, H-5''), 3.74 (1H, *dd*, *J* 9.4 and 3.2 Hz, H-3''), 3.42 (1H, *dq* *J* 9.4 and 6.1 Hz, H-2''), 3.32 (1H, *dd*, *J* 9.3 and 2.0, 4''), 0.93 (3H, *d*, *J* 6.1 Hz, H-6''). NMR ^{13}C (100 MHz, CD_3OD): δC 179.65 (C-4), 165.95 (C-7), 163.21 (C-5), 159.32 (C-2), 158.54 (C-9), 149.80 (C-4'), 146.41 (C-3'), 136.24 (C-3), 123.00 (C-1'), 122.87 (C-6'), 116.96 (C-2'), 116.38 (C-5'), 105.89 (C-10), 103.55 (C-1''), 99.85 (C-6), 94.74 (C-8), 73.27 (C-4''), 72.14 (C-3''), 72.03 (C-2''), 71.90 (C-5''), 17.64 (C-6''). ESI-MS *m/z* negative mode: 447.0885 [M-H]⁻; MS² [300.0248].

Luteolin 6-*C*- β -*D*-glucopyranoside or isoorientin (**4**) UV $\lambda_{\text{max}}^{\text{MeOH}}$ /nm: 269, 339; NMR ^1H (400 MHz, CD_3OD): δH 7.39 (1H, *dd*, *J* 8.6 and 2.3 Hz, H-6'), 7.36 (1H, *s*, H-2'), 6.91 (1H, *d*, *J* 8.6 and 1.4 Hz, H-5'), 6.56

(1H, *s*, H-3), 6.50 (1H, *s*, H-8), 4.91 (1H, *d*, *J* 9.8 Hz H-1''), 4.16 (1H, *t* like, *J* 9.5 Hz, H-2''), 3.88 (1H, *dd*, *J* 12.0 and 2.0 Hz, H-6''), 3.74 (1H, *dd*, *J* 12.0 and 5.2 Hz, H-6''), 3.50 (2H, *m*, H-4''), 3.49 (1H, *m*, H-3''), 3.47 (1H, *m*, H-5''). NMR ^{13}C (100 MHz, CD_3OD): δC 184.20 (C-4), 166.46 (C-2), 165.10 (C-7), 162.19 (C-5), 158.89 (C-9), 151.22 (C-4'), 147.22 (C-3'), 123.74 (C-1'), 120.48 (C-6'), 116.96 (C-5'), 114.32 (C-2'), 109.36 (C-6), 105.37 (C-10), 104.10 (C-3), 95.35 (C-8), 82.78 (C-5''), 80.29 (C-3''), 75.47 (C-1''), 72.77 (C-2''), 71.94 (C-4''), 63.01 (C-6''). ESI-MS *m/z* negative mode: 431.0866 [M-H]⁻; MS² [311.0480].

Isovitexin (**5**) UV $\lambda_{\text{max}}^{\text{MeOH}}$ /nm: 270, 337; NMR ^1H (400 MHz, CD_3OD): δH 7.91 (1H, *dd*, *J* 8.8 Hz, H-2' and H-6'), 6.99 (1H, *d*, *J* 8.8 Hz, H-3' and H-5'), 6.68 (1H, *s*, H-3), 6.59 (1H, *s*, H-8), 4.92 (1H, *d*, *J* 10.0 Hz, H-1''), 4.15 (1H, *m*, H-2''), 3.90 (1H, *dd*, *J* 12.0 and 2.0 Hz, H-6''b), 3.76 (1H, *d*, *J* 12.0 and 7.0 Hz, H-6''a), 3.47 (2H, *m*, H-3'' and H-4''), 3.41 (1H, *m*, H-5''). NMR ^{13}C (100 MHz, CD_3OD): δC 184.05 (C-4), 166.13 (C-2), 165.23 (C-7), 162.91 (C-9), 162.15 (C-4'), 158.77 (C-5), 2x 129.67 (C-2' and C-6'), 123.27 (C-1'), 2x 117.27 (C-3' and C-5'), 109.58 (C-6), 105.27 (C-10), 104.17 (C-3), 95.48 (C-8), 82.78 (C-5''), 80.31 (C-3''), 75.38 (C-1''), 72.65 (C-2''), 71.92 (C-4''), 62.95 (C-6''). ESI-MS *m/z* negative mode: 431.0910 [M-H]⁻; MS₂ [341.0611], MS₂ [311.0511].

Kaempferol 3-*O*- α -rhamnoside (**6**) UV $\lambda_{\text{max}}^{\text{MeOH}}$ /nm: 264, 344; NMR ^1H (400 MHz, $\text{DMSO}-d_6$): δH 7.75 (1H, *d*, *J* 8.8 Hz, H-2' and H-6'), 6.96 (1H, *d*, *J* 8.8 Hz, H-3' and H-5'), 6.38 (1H, *d*, *J* 2.0, H-8), 6.27 (1H, *d*, *J* 2.0 Hz, H-6), 5.51 (1H, *br s*, H-1''), 4.26 (1H, *dd*, *J* 3.0 and 1.2 Hz, H-2''), 3.71 (1H, *dd*, *J* 9.0 and 3.2 Hz, H-3''), 3.36 (1H, *m*, Hz, 4''), 3.30 (1H, *m*, H-5''), 0.96 (3H, *d*, *J* 6.0 Hz, H-6''). NMR ^{13}C (100 MHz, CD_3OD): δC 179.00 (C-4), 164.38 (C-7), 162.10 (C-5), 160.22 (C-4), 157.71 (C-2), 157.15 (C-9), 134.88 (C-3), 130.75 (C-2' and C-6'), 121.42 (C-1'), 2x 115.82 (C-3' and C-5'), 105.16 (C-10), 101.72 (C-1''), 99.29 (C-6), 94.10 (C-8), 72.50 (C-4''), 71.66 (C-3''), 70.52 (C-2''), 70.47 (C-5''), 17.44 (C-6''). ESI-MS *m/z* negative mode: 431.0873 [M-H]⁻; MS₂, MS₂ [285.0331].

Lupenone (**7**) NMR ^1H (400 MHz, CDCl_3): δH 4.69 (*br s*, H-29), 4.57 (*br s*, H-29), 2.42 (*m*, H-2), 2.38 (*m*, H-19), 1.69, 1.89 (*m*, H-1), 1.68 (H-30 and H-13), 1.46 (*m*, H-6), 1.42 (*m*, H-7), 1.38, 1.48 (*m*, H-16), 1.37 (*m*, H-9 and H-18), 1.32, 1.91 (*m*, H-21), 1.31 (*m*, H-5), 1.26, 1.40 (*m*, H-11), 1.19, 1.38 (*m*, H-22), 1.07 (*s*, H-25 and H-23 and H-26), 1.07, 1.69 (*m*, H-12), 1.03, 1.69 (*m*, H-15), 1.01 (*s*, 3H, H-24), 0.95 (*s*, 3H, H-27), 0.93 (*s*, 3H, H-25), 0.79 (*s*, 3H, H-28). NMR ^{13}C (100 MHz, CDCl_3): δC 218.31 (C-3), 150.91 (C-20), 109.41


(C-29), 54.94 (C-5), 49.81 (C-9), 48.26 (C-18), 47.98 (C-19), 47.36 (C-4), 43.01 (C-14), 42.91 (C-17), 40.80 (C-), 39.99 (C-22), 39.63 (C-1), 38.19 (C-13), 36.90 (C-10), 35.54 (C-16), 34.18 (C-2), 33.58 (C-7), 29.73 (C-21), 27.45 (C-15), 26.66 (C-23), 25.17 (C-12), 21.49 (C-11), 21.05 (C-24), 19.70 (C-6), 19.33 (C-30), 18.03 (C-28), 15.99 (C-25), 15.80 (C-26), 14.50 (C-27).

Assays with *Leishmania (Leishmania) amazonensis*

Leishmanicidal activity of the isolated compounds was determined against amastigote-like forms as previously described by Cota et al. (2008a). Briefly: promastigotes of *Leishmania (L.) amazonensis* (strain IFLA/BR/196/PH-8) were obtained from lesions of infected hamsters. The parasites were grown at 26 °C in pH 7.2 Schneider's medium, and then stimulated to differentiate into the amastigote forms by rising the temperature (32 °C), and lowering the pH (6.0) of the Schneider's medium. After seven days under these conditions, 90% of the promastigotes were transformed into amastigote-like forms, verified by means of microscope, and then used in the bioassays. Amastigote density was adjusted to 1×10^8 parasites per mL, and 90 μ L added to each well of 96-well plates. Solutions at 200 μ g/mL in DMSO (1% in water) were performed for each fraction or compound and then ten microliters of the solution were added to each well of 96-well plates. The plates were incubated at 32 °C for 72 h, and then the cells viability was determined using the MTT (methyl thiazolyl tetrazolium) assay (Teixeira et al., 2002). The results were expressed as percent inhibition in relation to the controls without drug. Amphotericin B at 0.2 μ g/mL (Fungison[®] Bristol-Myers Squibb B, Brazil) was used as a positive drug control. All assays were performed in triplicate.

Results and Discussion

The structures of isolated compounds were




- 1 R₁=OH; R₂=O- -gal; R₃=H
- 2 R₁=OH; R₂=O- -glc; R₃=H
- 3 R₁=OH; R₂=O- -rham; R₃=H
- 4 R₁=OH; R₂=H; R₃=C- -glc
- 5 R₁=R₂=H; R₃=C- -glc
- 6 R₁=R₃=H; R₂=O- -rham

identified by a combination of spectroscopic methods (MS, 1 D NMR and 2D NMR) and comparisons with the literature data as a mixture of quercetin 3- β -O-galactopyranoside or hyperoside (**1**) and quercetin 3- β -O-glucopyranoside or isoquercitrin (1:1) (**2**) (Almeida et al., 1998; Costa et al., 2007; Ohguchi et al., 2010), quercetin 3-O- α -rhamnoside or quercitrin (**3**) (Zhong et al., 1997), luteolin 6-C- β -glucopyranoside or orientin (**4**) (Li et al., 2009), apigenin-6-C-glucoside or isovitexin (**5**) (Leong et al., 2010), kaempferol 3-O- α -rhamnoside (**6**) (Fossen et al., 1999) and lupenone (**7**) (Prachayasittikul et al., 2010), respectively. Characterization of the sugar moieties from the mixture of quercetin 3- β -galactopyranoside (**1**) and quercetin 3- β -glucopyranoside (**2**) was performed according Cota et al. 2008b.

In this work, we isolated six flavonoids (**1-6**) and one lupane triterpene (**7**) for the first time from *Kielmeyera variabilis* Mart. & Zucc., Calophyllaceae. Triterpenes with different carbon skeletons that include lupeol, friedelin, β -amyrin and the sterol β -sitosterol, were found in *K. rugosa* Choisy (Nogueira et al., 2008), *K. lathrophyton* Saddi (Cruz et al., 2001), *K. candidissima* (Ferreira et al., 1972), *K. speciosa* A. St.-Hil. (Gottlieb et al., 1970) and *K. rugosa* Choisy (Nogueira et al., 2008), respectively. Indeed, Garcia-Cortez et al. (1998) obtained a mixture of 27-O-*cis* and trans-*p*-coumaroylcyclicodiscic acid, lupeol derivated triterpenes from a dichloromethane extract of *K. coriaceae* leaves. Our chemical study on the aqueous-soluble fraction resulted in the isolation of glycosilated flavonoids, where two of them are C-flavonoids (**4-5**). The major compound from Aq soluble fraction in *K. variabilis* was found to be quercitrin (**3**) what suggests that this species is a good source of this flavonoid.

Many phenolic compounds such as xanthenes (Pinheiro et al., 2003a; Sobral et al., 2009), quinones (Corrêa et al., 1970) and coumarins (Cruz et al., 1998b) have been isolated from *Kielmeyera* species. Despite 4-alkyl and 4-phenyl coumarins having been identified as the major constituents of some species that occur mainly



7

in the cerrado (Gottlieb et al., 1969; Gottlieb & Stefani, 1970) and restinga (Cruz et al., 2002; Nogueira et al., 2008) and xanthenes from species collected in the atlantic forest (Garcia-Cortez et al., 1998; Pinheiro et al., 2003a), *K. variabilis* did not demonstrate the same pattern as observed by previous studies. The presence of flavonoids in this species is in agreement with previous reports from other genera in the Calophylleae tribe as *Caraipa* (Silveira et al., 2010), *Mesua* (Banerji & Chowdhury, 1993) and *Calophyllum* (Cechinel et al., 2009). It is known that temperature and UV radiation can up-regulate key genes (Christie & Jenkins, 1996) that give plants the ability to offset the excessive absorption of UV radiation by accumulating UV-filtering secondary metabolites such as flavonoids (Cuadra et al., 1997). The chemotaxonomic applicability of phenolic compounds from *Kielmeyera* genus as markers needs to be properly evaluated, considering more members of the genus, their distribution and environmental influence.

The crude organic extract from *K. variabilis* leaves exhibited 95% of leishmanicidal activity at 20 µg/mL on amastigote-like form of *Leishmania amazonensis*. From isolated compounds 1-7, only quercitrin was able to inhibit the growth amastigote-like form of *L. amazonensis*, but it was not dose-dependent in this assay (0.78-20 µg/mL, 40-45%). The major compound from Aq soluble fraction in *K. variabilis* was found to be quercitrin what suggests that this species is a good source of this flavonoid. The potent antileishmanial activity of quercitrin was demonstrated by *in vitro* bioguided fractionation of *Kalanchoe pinnata* (Muzitano et al., 2006a). In addition, previous study showed that oral administration of 16 mg/kg body weight of quercetin 3-*O*- α -L-arabinopyranosyl (1 \rightarrow 2)- α -L-rhamnopyranoside or quercitrin or quercetin for 30 days in BALB/c mice suppressed parasite burden by 65%, 57%, and 76% in the lesion from infected ear compared with the untreated mice and similar to intraperitoneal pentostam (62%) (Muzitano et al., 2006b). These biological results and our *in vitro* studies are good evidence that synergistic effects of flavonoid glycosides are the cause of significant leishmanicidal activity of crude organic extract from *K. variabilis* leaves.

Acknowledgment

The authors thank Fiocruz and Fapemig for financial support. We are also grateful to Daniela Nabak Bueno Maia and Luciana Alves Rodrigues dos Santos Lima for technical assistance.

References

- Almeida AP, Miranda MMFS, Simoni IC, Wigg MD, Lagrota MHC, Costa SS 1998. Flavonol monoglycosides isolated from the antiviral fractions of *Persea americana* (Lauraceae) leaf infusion. *Phytother Res* 12: 562-567.
- Audi EA, Otobone F, Martins JV, Cortez DA 2002. Preliminary evaluation of *Kielmeyera coriacea* leaves extract on the central nervous system. *Fitoterapia* 73: 517-519.
- Banerji R, Chowdhury AR 1993. *Mesua ferrea*: Chemical constituents and biological. *J Chem Soc Pakistan* 15: 207-211.
- Bittrich V 2010. Clusiaceae in lista de esp c ias da ora do Brasil. *Jardim Botânico do Rio de Janeiro, Rio de Janeiro, Brazil*. <http://floradobrasil.jbrj.gov.br/2010/FB000089>, Accessed 1 Feb 2011.
- Cechinel Filho V, Meyre-Silva C, Niero R 2009. Chemical and pharmacological aspects of the genus *Calophyllum*. *Chem Biodivers* 6: 313-327.
- Christie JM, Jenkins GI 1996. Distinct UV-B and UV-A/blue light signal transduction pathways induce chalcone synthase gene expression in *Arabidopsis* cells. *Plant Cell* 8: 1555-1567.
- Corrêa DB, Silva, LGF, Gottlie OR, Gonçalves SJ 1970. Quinone and xanthone constituents of *Kielmeyera rupestris*. *Phytochemistry* 9: 447-451.
- Costa DA, Silva DA, Cavalcanti AC, Medeiros MAA, Lima JT, Cavalcante JMS, Silva BA, Agra MF, Souza MFV 2007. Chemical constituents from *Bakeridesia pickelii* Monteiro (Malvaceae) and the relaxant activity of kaempferol-3-*O*- β -D-(6''-*E*-*p*-coumaroyl) glucopyranoside on guinea-pig ileum. *Quim Nova* 30: 901-903.
- Cota BB, Rosa LH, Caligiorne RB, Rabello ALT, Alves TMA, Rosa CA, Zani CL 2008a. Altenusin, a biphenyl isolated from the endophytic fungus *Alternaria* sp., inhibits trypanothione reductase from *Trypanosoma cruzi*. *FEMS Microbiol Lett* 285: 177-182.
- Cota BB, Magalhães A, Pimenta AMC, Siqueira EP, Alves TMA, Zani CL, 2008b. Chemical constituents of *Habenaria petalodes* Lindl. (Orchidaceae). *J Braz Chem Soc* 19: 1098-1104.
- Cuadra P, Harborne JB, Waterman PG 1997. Increases in surface flavonols and photosynthetic pigments in *Gnaphalium luteo-album* in response to UV-B radiation. *Phytochemistry* 45: 1377-1383.
- Cruz FG, David JM, Guedes MLS, Chávez JP, Moreira LM 1998a. Coumarins from *Kielmeyera reticulata*. *Phytochemistry* 47: 1363-1366.
- Cruz FG, David JM, Guedes MLS, Chávez JP, Santos NAS 1998b. Coumarins from *Kielmeyera argentea*. *Phytochemistry* 48: 703-706.
- Cruz FG, Silva-Neto JT, Guedes MLS 2001. Xanthenes and coumarins from *Kielmeyera lathrophyton*. *J Braz Chem Soc* 12: 117-122.
- Cruz FG, Moreira LM, Santos NAS, Guedes MLS 2002. Additional coumarins from *Kielmeyera reticulata*. *J Braz Chem Soc* 13: 704-707.
- Ferreira GAL, Gottlieb OR, Mesquita AAL 1972. Xanthenes from a *Kielmeyera* species *Phytochemistry* 11: 1512-

- 1513.
- Fossen T, Larsen A, Kiremire BT, Andersen M 1999. Flavonoids from blue flowers of *Nymphaea caerulea*. *Phytochemistry* 51: 1133-1137.
- Garcia Cortez DA, Young MCM, Marston A, Wolfender JL, Hostettmann K 1998. Xanthonas, triterpenes and a biphenyl from *Kielmeyera coriacea*. *Phytochemistry* 47: 1367-1374.
- Garcia Cortez DAG, Abreu Filho BA, Nakamura CV, Dias Filho BP, Marston A, Hostettmann K 2002. Antibacterial activity of a biphenyl and xanthonas from *Kielmeyera coriacea*. *Pharm Biol* 40: 485-489.
- Gottlieb OR, Lins Mesquita AA, Martins da Silva E, Teixeira de Melo M 1969. Xanthonas of *Kielmeyera ferruginea*. *Phytochemistry* 8: 665-666.
- Gottlieb OR, Stefani GM 1970. Xanthonas from *Kielmeyera excelsa*. *Phytochemistry* 9: 453-454.
- Gottlieb OR, Mesquita AAL, Oliveira GG, Melo MT 1970. Xanthonas from *Kielmeyera speciosa*. *Phytochemistry* 9: 2537-2544.
- Gramacho RS, Nagem TJ, Oliveira TT, Queiroz MELR, Neves AA, Saddi N 1999. Phenylcoumarins from *Kielmeyera elata*. *Phytochemistry* 51: 579-581.
- Leong ACN, Kinjo Y, Tako M, Iwasaki H, Oku H, Tamaki H 2010. Flavonoid glycosides in the shoot system of Okinawa Taumu (*Colocasia esculenta* S.). *Food Chem* 119: 630-635.
- Li L, Henry GE, Seeram NP 2009. Identification and bioactivities of resveratrol oligomers and flavonoids from *Carex folliculata* seeds. *J Agric Food Chem* 57: 7282-7287.
- Lopes JLC, Lopes JNC, Gilbert B, Bonini SE 1977. Osajaxanthone from *Kielmeyera coriacea*. *Phytochemistry* 16: 1101-1101.
- Martins JVC, Otobone FJ, Sela VR, Obici S, Trombelli MA, Cortez DAG, Audi EA 2004. Activity of hydroethanolic extract from *Kielmeyera coriacea* stems on central nervous system in rats. *Acta Sci Health Sci* 26: 365-368.
- Martins JVC, Otobone FJ, Sela VR, Obici S, Trombelli MA, Cortez DAG, Audi EA 2006. Behavioral effects of *Kielmeyera coriacea* extract in rats. *Indian J Pharmacol* 38: 427-428.
- Mesquita ML, de Paula JE, Pessoa C, de Moraes MO, Costa-Lotufo LV, Grougnet R, Michel S, Tillequin F, Espindola LS 2009. Cytotoxic activity of Brazilian Cerrado plants used in traditional medicine against cancer cell lines. *J Ethnopharmacol* 123: 439-445.
- Muzitano MF, Cruz EA, Almeida AP, Silva SAG, Kaiser CR, Guette C, Rossi-Bergmann B, Costa SS 2006a. Quercitrin: An antileishmanial flavonoid glycoside from *Kalanchoe pinnata*. *Planta Med* 72: 81-83.
- Muzitano MF, Tinoco LW, Guette C, Kaiser CR, Rossi-Bergmann B, Costa SS 2006b. Assessment of antileishmanial activity of new and unusual flavonoids from *Kalanchoe pinnata*. *Phytochemistry* 67: 2071-2077.
- Nogueira PCL, Moacir SA, Larissa MA, Moraes VRS, Ribeiro AS, Bittrich V, Amaral MCE, Ferreira AG, Alcântara GB, Leão KV, Alves PB 2008. Chemical constituents from *Kielmeyera rugosa* Choisy (Clusiaceae). *Biochem Syst Ecol* 36: 921-924.
- Ohguchi K, Nakajima C, Oyama M, Inuma M, Itoh T, Akao Y, Nozawa Y, Ito M 2010. Inhibitory effects of flavonoid glycosides isolated from the peel of Japanese Persimmon (*Diospyros kaki* 'Fuyu') on melanin biosynthesis. *Biol Pharm Bull* 33: 122-124.
- Pinheiro L, Vidotti GJ, Young MCC, Ferreira AG, Cortez, Diogenes AG 2003a. Phytochemical study and evaluation of the molluscicidal activity of *Kielmeyera variabilis* Mart. (Clusiaceae). *Quim Nova* 26: 157-160.
- Pinheiro L, Nakamura CV, Dias Filho BP, Ferreira AG, Young MCM, Garcia Cortez DA 2003b. Antibacterial xanthonas from *Kielmeyera variabilis* Mart. (Clusiaceae). *Mem I Oswaldo Cruz* 98: 549-552.
- Prachayasittikul S, Saraban P, Cherdtrakulkiat R, Ruchirawat S, Prachayasittikul V 2010. New bioactive triterpenoids and antimalarial activity of *Diospyros rubra* Lec. *Excli J* 9: 1-10.
- Scio E, Ribeiro A, Alves TMA, Álvaro JR, Young GS, Cordell GA, Zani CL 2003. New bioactive coumarins from *Kielmeyera albopunctata*. *J Nat Prod* 66: 634-637.
- Silveira CV, Trevisan MTS, Rios JB, Erben G, Haubner R, Pfundstein B, Owen RW 2010. Secondary plant substances in various extracts of the leaves, fruits, stem and bark of *Caraipa densifolia* Mart. *Food Chem Toxicol* 48: 1597-1606.
- Sobral IS, Souza-Neta LC, Costa GAN, Guedes MLS, Martins D, Cruz FG 2009. Xanthonas, triterpenos e atividade antibacteriana do extrato em diclorometano de *Kielmeyera cuspidata* Saddi, Clusiaceae. *Rev Bras Farmacogn* 19: 686-689.
- Teixeira MCA, Santos RJ, Sampaio RB, Carvalho LP, Santos WLC 2002. A simple and reproducible method to obtain large numbers of axenic amastigotes of different *Leishmania* species. *Parasitol Res* 88: 963-968.
- Zhong XN, Otsuka H, Ide T, Hirata E, Takushi A, Takeda Y 1997. Three flavonol glycosides from leaves of *Myrsine seguinii*. *Phytochemistry* 46: 943-946.

*Correspondence

Betania Barros Cota
Laboratório de Química de Produtos Naturais, Centro de Pesquisas René Rachou, Fundação Oswaldo Cruz
Av. Augusto de Lima, 1715, 30190-002 Belo Horizonte-MG, Brazil
betania@cpqrr.fiocruz.br
Tel.: +55 31 3349 7845
Fax: +55 31 3295 3115