

















- RW 1993. Accessory cell function of human eosinophils. HLA-DR-dependent, MHC-restricted antigen-presentation and IL-1 alpha expression. *J Immunol* 150: 2554-2562.
- Woerly G, Decot V, Loiseau S, Loyens M, Chihara J, Ono N, Capron M 2004. CD28 and secretory immunoglobulin A-dependent activation of eosinophils: inhibition of mediator release by the anti-allergic drug, suplatast tosilate. *Clin Exp Allergy* 34: 1379-1387.
- Woerly G, Lacy P, Younes AB, Roger N, Loiseau S, Moqbel R, Capron M 2002. Human eosinophils express and release IL-13 following CD28-dependent activation. *J Leukoc Biol* 72: 769-779.
- Woerly G, Roger N, Loiseau S, Dombrowicz D, Capron A, Capron M 1999. Expression of CD28 and CD86 by human eosinophils and role in the secretion of type 1 cytokines (interleukin 2 and interferon gamma): inhibition by immunoglobulin A complexes. *J Exp Med* 190: 487-495.
- Ying S, Meng Q, Taborda-Barata L, Corrigan CJ, Barkans J, Assoufi B, Moqbel R, Durham SR, Kay AB 1996. Human eosinophils express messenger RNA encoding RANTES and store and release biologically active RANTES protein. *Eur J Immunol* 26: 70-76.
- Yu W, Bozza PT, Tzizik DM, Gray JP, Cassara J, Dvorak AM, Weller PF. 1998. Co-compartmentalization of MAP kinases and cytosolic phospholipase A2 at cytoplasmic arachidonate-rich lipid bodies. *Am J Pathol* 152: 759-769.

